

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number  
**WO 03/087351 A1**

(51) International Patent Classification<sup>7</sup>: C12N 7/00,  
15/01, 15/36, 15/51, C12Q 1/70

(21) International Application Number: PCT/AU03/00432

(22) International Filing Date: 11 April 2003 (11.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PS 1710 12 April 2002 (12.04.2002) AU  
PS 3224 26 June 2002 (26.06.2002) AU

(71) Applicants (for all designated States except US): MEL-  
BOURNE HEALTH [AU/AU]; 10th Floor, Connibere  
Building, Royal Melbourne Hospital, Flemington Road,  
Parkville, VIC 3050 (AU). AUSTIN & REPATRIATION  
MEDICAL CENTRE [AU/AU]; Studley Road, Heidel-  
berg, VIC 3081 (AU). SOUTHERN HEALTH [AU/AU];  
Monash Medical Centre, 246 Clayton Road, Clayton, VIC  
3168 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only):  
BARTHOLOMEUSZ, Angeline, Ingrid [AU/AU]; 64  
Miller Street, Carnegie, VIC 3613 (AU). LOCARNINI,  
Stephen, Alister [AU/AU]; 13 Carlisle Avenue, East  
St Kilda, VIC 3182 (AU). AYRES, Anna [AU/AU];  
4 Ferry Crescent, West Brunswick, VIC 3056 (AU).  
COLLEDGE, Danielle [AU/AU]; 12 Buttercup Road,  
Bundoora, VIC 3083 (AU). SASADEUSZ, Joseph  
[AU/AU]; c/- Melbourne Health, 10th Floor, Connibere

Street, Royal Melbourne Hospital, Flemington Road,  
Parkville, VIC3050 (AU). TILLMANN, Hans [DE/DE];  
Stolzestr 41, D-30171 Hannover (DE). ANGUS, Peter,  
William [AU/AU]; 45 Streeton Crescent, East Ivanhoe,  
VIC 3079 (AU). SIEVERT, William [AU/AU]; c/-  
Department of Medicine, Monash Medical Centre, 246  
Clayton Road, Clayton, VIC 3168 (AU).

(74) Agents: HUGHES, E, John, L et al.; Davies Collison  
Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU).

(81) Designated States (national): AE, AG, AL, AM, AT; AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,  
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 03/087351 A1

(54) Title: HEPATITIS B VIRAL VARIANTS WITH REDUCED SUSCEPTIBILITY TO NUCLEOSIDE ANALOGS AND USES THEREOF

(57) Abstract: The present invention relates generally to viral variants exhibiting reduced sensitivity to particular agents and/or reduced interactivity with immunological reagents. More particularly, the present invention is directed to hepatitis B virus (HBV) variants exhibiting complete or partial resistance to nucleoside or nucleotide analogs and/or reduced interactivity with antibodies to viral surface components including reduced sensitivity to these antibodies. The present invention further contemplates assays for detecting such viral variants, which assays are useful in monitoring anti-viral therapeutic regimens and in developing new or modified vaccines directed against viral agents and in particular HBV variants. The present invention also contemplates the use of the viral variants to screen for and/or develop or design agents capable of inhibiting infection, replication and/or release of the virus.

# PATENT COOPERATION TREATY

## PCT

### INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

REC'D 17 JUN 2003

WIPO PCT

Applicant's or agent's file reference <b>12175980/EJH</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/AU03/00432</b>	International filing date (day/month/year) <b>11 April 2003</b>	(Earliest) Priority Date (day/month/year) <b>12 April 2002</b>
Applicant <b>MELBOURNE HEALTH et al</b>		

This international search report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This international search report consists of a total of 7 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

**1. Basis of the report**

a. With regard to the language, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.  
☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

2. ☒ Certain claims were found unsearchable (See Box I).

3. ☒ Unity of invention is lacking (See Box II).

4. With regard to the title, ☐ the text is approved as submitted by the applicant.  
☒ the text has been established by this Authority to read as follows:

**HEPATITIS B VIRAL VARIANTS WITH REDUCED SUSCEPTIBILITY TO NUCLEOSIDE ANALOGS AND USES THEREOF**

5. With regard to the abstract, ☒ the text is approved as submitted by the applicant  
☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the drawings to be published with the abstract is Figure No. ☐ as suggested by the applicant. ☒ None of the figures  
☐ because the applicant failed to suggest a figure  
☐ because this figure better characterizes the invention

## VIRAL VARIANTS, DETECTION AND USE

### BACKGROUND OF THE INVENTION

#### 5 FIELD OF THE INVENTION

The present invention relates generally to viral variants exhibiting reduced sensitivity to particular agents and/or reduced interactivity with immunological reagents. More particularly, the present invention is directed to hepatitis B virus (HBV) variants exhibiting  
10 complete or partial resistance to nucleoside or nucleotide analogs and/or reduced interactivity with antibodies to viral surface components including reduced sensitivity to these antibodies. The present invention further contemplates assays for detecting such viral variants, which assays are useful in monitoring anti-viral therapeutic regimens and in developing new or modified vaccines directed against viral agents and in particular HBV  
15 variants. The present invention also contemplates the use of the viral variants to screen for and/or develop or design agents capable of inhibiting infection, replication and/or release of the virus.

#### DESCRIPTION OF THE PRIOR ART

20

Bibliographic details of the publications referred to in this specification are also collected at the end of the description.

The reference to any prior art in this specification is not, and should not be taken as, an  
25 acknowledgment or any form of suggestion that that prior art forms part of the common general knowledge in any country.

Hepatitis B virus (HBV) can cause debilitating disease conditions and can lead to acute liver failure. HBV is a DNA virus which replicates *via* an RNA intermediate and utilizes  
30 reverse transcription in its replication strategy (Summers and Mason, *Cell* 29: 403-415, 1982). The HBV genome is of a complex nature having a partially double-stranded DNA

- 2 -

structure with overlapping open reading frames encoding surface, core, polymerase and X genes. The complex nature of the HBV genome is represented in Figure 1. The polymerase consists of four functional regions, the terminal protein (TP), spacer, reverse transcriptase (rt) and ribonuclease (RNase).

5

The polymerase gene of HBV overlaps the envelope gene, mutations in the catalytic domain of the polymerase gene can also affect the nucleotide and the deduced amino acid sequence of the envelope protein and *vice versa*. In particular, the genetic sequence for the neutralization domain of HBV known as the 'a' determinant, which is found within the HBsAg and located between amino acids 99 and 169, actually overlaps the major catalytic regions of the viral polymerase protein and in particular domains A and B.

The presence of an HBV DNA polymerase has led to the proposition that nucleoside or nucleotide analogs could act as effective anti-viral agents. Examples of nucleoside analogs currently being tested are penciclovir and its oral form (FCV) [Vere Hodge, *Antiviral Chem Chemother* 4: 67-84, 1993; Boyd *et al.*, *Antiviral Chem Chemother*. 32: 358-363, 1987; Kruger *et al.*, *Hepatology* 22: 219A, 1994; Main *et al.*, *J. Viral Hepatitis* 3: 211-215, 1996], Lamivudine[(-)- $\beta$ -2'-deoxy-3'-thiacytidine]; (3TC or LMV) [Severini *et al.*, *Antimicrobial Agents Chemother*. 39: 430-435, 1995; Dienstag *et al.*, *New England J Med* 333: 1657-1661, 1995]. New nucleoside or nucleotide analogs which have already progressed to clinical trials include the pyrimidines Emtricitabine, ((-)- $\beta$ -L-2'-3'-dideoxy-5-fluoro-3'-thiacytidine; FTC), the 5-fluoro derivative of 3TC, and Clevudine (1-(2-fluoro-5-methyl- $\beta$ -L-arabino-furanosyl) uracil; L-FMAU), a thymidine analog. Like 3TC, these are pyrimidine derivatives with an unnatural "L"- configuration. Several purine derivatives have also progressed to clinical trials; they include Entecavir (BMS-200, 475; ETV), a carbocyclic deoxyguanosine analog, diaminopurine dioxolane (DAPD), an oral pro-drug for dioxolane guanine ((-)- $\beta$ -D-2-aminopurine dioxolane; DXG) and Adefovir dipivoxil, an oral prodrug for the acyclic deoxyadenosine monophosphate nucleoside analog Adefovir (9-[phosphonyl-methoxyethyl]-adenine; PMEA). Other drugs in pre-clinical and clinical trials include include FLG [Medivir], ACH-126,443 (L-d4C) [Archillion Pharmaceuticals], ICN 2001-3 (ICN) and Racivir (RCV) [Pharmasset].



- 3 -

Whilst these agents are highly effective in inhibiting HBV DNA synthesis, there is the potential for resistant mutants of HBV to emerge during long term antiviral chemotherapy. In patients on prolonged LMV therapy, key resistance mutations are selected in the rt domain within the polymerase at rtM204I/V +/- rtL180M as well as other mutations. The nomenclature used for the polymerase mutations is in accordance with that proposed by Stuyver *et al.*, 2001, *supra*. LMV is a nucleoside analog that has been approved for use against chronic HBV infection. LMV is a particularly potent inhibitor of HBV replication and reduces HBV DNA titres in the sera of chronically infected patients after orthotopic liver transplantation (OLT) by inhibiting viral DNA synthesis. LMV monotherapy seems unlikely to be able to control HBV replication in the longer term. This is because emergence of LMV-resistant strains of HBV seems almost inevitable during monotherapy.

Adefovir dipivoxil (ADV: formerly, bis-pom PMEA) is an orally available prodrug of the acyclic deoxyadenosine monophosphate analog adefovir (formerly, PMEA) (Figure 2). ADV is also a potent inhibitor of HBV replication and has recently been given FDA approval for use against chronic HBV infection. Adefovir dipivoxil differs from other agents in this class in that it is a nucleotide (vs. nucleoside) analog and as such bypasses the first phosphorylation reaction during drug activation. This step is often rate-limiting. Adefovir dipivoxil has demonstrated clinical activity against both wild-type and lamivudine-resistant strains of HBV and is currently in phase III clinical Testing (Gilson *et al.*, *J Viral Hepat* 6: 387-395, 1999; Perrillo *et al.*, *Hepatology* 32: 129-134, 2000; Peters *et al.*, *Transplantation* 68: 1912-1914, 1999; Benhamou *et al.*, *Lancet* 358: 718-723, 2001). During phase II studies a 30 mg daily dose of adefovir dipivoxil resulted in a mean 4 log<sub>10</sub> decrease in viremia over 12 weeks (Heathcote *et al.*, *Hepatology* 28: A620, 1998).

ADV is a substituted acyclic nucleoside phosphonate. This class of compounds also includes tenofovir disoproxil fumarate (also referred to as tenofovir DF, or tenofovir, or (TFV) or 9-R-(2-phosphonomethoxypropyl)adenine (PMPA) and is marketed as Viread by Gilead sciences).

- 4 -

TFV has antiviral activity against both HBV and HIV (Ying *et al.*, *J Viral Hepat.* 7(2): 161-165, 2000; Ying *et al.*, *J. Viral Hepat.* 7(1): 79-83, 2000; Suo *et al.*, *J Biol Chem.* 273(42): 27250-27258, 1998).

- 5    FTC has activity against HBV and HIV (Frick *et al.*, *Antimicrob Agents Chemother* 37: 2285-2292, 1993).

10    Nucleoside or nucleotide analog therapy may be administered as monotherapy or combination therapy where two or more nucleoside or nucleotide analogs may be administered. The nucleoside or nucleotide analogs may also be administered in combination with other antiviral agents such as interferon or hepatitis B immunoglobulin (HBIG).

15    There is a need to monitor for the emergence of nucleoside/nucleotide-analog- or antibody-resistant strains of HBV and to develop diagnostic protocols to detect these resistant viruses and/or to use them to screen for and/or develop or design agents having properties making them useful as anti-viral agents. Defective forms of these resistant strains or antigenic components therefrom are also proposed to be useful in the development of therapeutic vaccine compositions as are antibodies directed to viral surface components.

## SUMMARY OF THE INVENTION

Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the  
5 inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

Nucleotide and amino acid sequences are referred to by a sequence identifier number (SEQ ID NO:). The SEQ ID NOs: correspond numerically to the sequence identifiers <400>1  
10 (SEQ ID NO:1), <400>2 (SEQ ID NO:2), etc. A summary of the sequence identifiers is provided in Table 1. A sequence listing is provided after the claims.

Specific mutations in an amino acid sequence are represented herein as “Xaa<sub>1</sub>nXaa<sub>2</sub>” where Xaa<sub>1</sub> is the original amino acid residue before mutation, n is the residue number and  
15 Xaa<sub>2</sub> is the mutant amino acid. The abbreviation “Xaa” may be the three letter or single letter (i.e. “X”) code. An “rt” before “Xaa<sub>1</sub>nXaa<sub>2</sub>” means “reverse transcriptase”. An “s” means an envelope gene. The amino acid residues for HBV DNA polymerase are numbered with the residue methionine in the motif Tyr Met Asp Asp (YMDD) being residue number 204 (Stuyver *et al.*, *Hepatology* 33: 751-757, 2001). The amino acid  
20 residues for hepatitis B virus surface antigen are number according to Norder *et al.* (*J. Gen. Virol.* 74: 341-1348, 1993). Both single and three letter abbreviations are used to define amino acid residues and these are summarized in Table 2.

In accordance with the present invention, the selection of HBV variants is identified in  
25 patients (Patient A, C and D) with chronic HBV infection treated with ADV and liver transplant patients (Patients B and E) treated with both ADV and LMV post-OLT or ADV post-transplant. HBV variants from Patients F, G and H were also identified following similar treatments. Variants of HBV are identified during ADV or combination ADV and LMV treatment with mutations in the HBV DNA polymerase gene which reduce the  
30 sensitivity of HBV to this nucleoside analog. Consequently, HBV rt variants are contemplated which are resistant to, or which exhibit reduced sensitivity to, ADV,

- 6 -

LMV,TFV, FTC, ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combinations thereof . Corresponding mutations in the surface antigen also occur. The identification of these HBV variants is important for the development of assays to monitor ADV, LMV, FTC and/or TFV resistance and/or resistance to other nucleoside or nucleotide analogs or other anti-HBV agents or combinations thereof and to screen for agents which are useful as alternative therapeutic agents.

10

Reference herein to "anti-HBV agents" includes nucleoside and nucleotide analogs as well as immunological reagents (e.g. antibodies to HBV surface components) and chemical, proteinaceous and nucleic acid agents which inhibit or otherwise interfere with viral replication, maintenance, infection, assembly or release.

15

The detection of such HBV variants is particularly important in the management of therapeutic protocols including the selection of appropriate agents for treating HBV infection. The method of this aspect of the present invention is predicated in part on monitoring the development in a subject of an increased HBV load in the presence of a nucleoside or nucleotide analog or other anti-HBV agents or combinations thereof. The clinician is then able to modify an existing treatment protocol or select an appropriate treatment protocol accordingly.

Accordingly, one aspect of the present invention is directed to an isolated HBV variant comprising a nucleotide mutation in a gene encoding a DNA polymerase resulting in at least one amino acid addition, substitution and/or deletion to the DNA polymerase and which exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combinations thereof. The variant HBV

30

- 7 -

comprises a mutation in an overlapping open reading frame in its genome in a region defined by one or more of domains F and G and domain A through to E of HBV DNA polymerase.

- 5 Another aspect of the present invention provides an isolated HBV variant comprising a nucleotide mutation in the S gene resulting in at least one amino acid addition, substitution and/or deletion to the surface antigen and which exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV,  
 10 TFV and FTC and LMV, or ADV and FTC and LMV and TFV, ADV and LMV and FTC, and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combinations thereof.

Useful mutants in the rt region include, in one embodiment, rtS21A, rtL122F, rtN124H,  
 15 rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and  
 20 rtV214A; and in yet another embodiment, rtH90D and rtL/F108L; and in still a further embodiment, rtL157L/M, rtA181V and rtV207I and in yet a further embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; and in another embodiment, rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H or a combination thereof or an equivalent mutation.

25

Other HBV variants are also contemplated with mutations in rt at rtK32, rtN33, rtP34, rtH35 and rtT37 (these are upstream of the F domain of the DNA polymerase), rtP59, rtK60, rtF61, rtA62 and rtV63 (these are located between the F and A domains), rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91 (these are located within the A domain and  
 30 the region immediately prior to and following), rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184 (these are located in the B domain), rtM204 and rtY203 (these

- 8 -

are located in the C domain), rt235, rt236, rt237, rt238 and rt239 (these are located in the D domain) and rt247, rt248, rt249, rt250 and rt251 (these are located in the E domain) or a combination thereof or an equivalent mutation.

5 Useful mutants are provided below (see also Tables 16 and 17):

- K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;
- N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;
- P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;
- 10 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;
- T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;
- P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;
- K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;
- F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;
- 15 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;
- V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;
- D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;
- V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;
- S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;
- 20 A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;
- Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;
- H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;
- I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;
- P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;
- 25 F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;
- L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;
- L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;
- A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;
- Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;
- 30 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;
- T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;

- 9 -

Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 5 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 10 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 15 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion.

20

Reference above to "deletion" means that the first mentioned amino acid before the residue number has been deleted.

Useful mutations in the S gene include, in one embodiment, sP120T, sM125T and  
 25 sT127A; in another embodiment, T118R, sM133T, sF134V sI195M, sS207R and  
 sY225Y/C; in a further embodiment, sS126T, sM133L/M, sS143S/T, sD144A sG145A  
 and sW172Stop; in yet a further embodiment, sN40S, sC69 Stop, sM75I, sL88P, sT118A,  
 sW182stop, sW196L, sY206H and sY225F; and in yet another embodiment, sI81M and  
 sP214Q; and in still another embodiment, sF83S, sL173F and sW199L; and in still yet  
 30 another embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L  
 and sY221C; and in yet another embodiment, sC69Stop/C, sC76Y sI110V/I, sY134N,

- 10 -

sW172Stop/W, sW196Stop and sS207R or a combination thereof or an equivalent mutation.

The present invention further contemplates a method for determining the potential for an  
5 HBV to exhibit reduced sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV  
and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and  
LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and  
FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or  
nucleotide analogs or other anti-HBV agents or combination thereof by isolating DNA or  
10 corresponding mRNA from the HBV and screening for a mutation in the nucleotide  
sequence encoding HBV DNA polymerase resulting in at least one amino acid substitution,  
deletion and/or addition in any one or more of domains F and G and domains A through to  
E or a region proximal thereto of the DNA polymerase and associated with resistance or  
decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV,  
15 LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and  
TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or  
ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide  
analogs or other anti-HBV agents or combination thereof. The presence of such a mutation  
is an indication of the likelihood of resistance to ADV, LMV, TFV, or FTC, or ADV and  
20 LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or  
ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and  
LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or  
nucleotide analogs or other anti-HBV agents or combination thereof.

25 The present invention also provides a composition comprising a variant HBV resistant to  
ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and  
ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and  
TFV, TFV and FTC and LMV, or ADV and FTC and LMV and TFV, ADV and LMV and  
FTC, and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or  
30 combination thereof or an HBV surface antigen from the variant HBV or a recombinant or



- 11 -

derivative form thereof or its chemical equivalent and one or more pharmaceutically acceptable carriers and/or diluents.

- Yet another aspect of the present invention provides a use of the aforementioned composition or a variant HBV comprising a nucleotide mutation in a gene encoding a DNA polymerase resulting in at least one amino acid addition, substitution and/or deletion to the DNA polymerase and a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof in the manufacture of a medicament for the treatment and/or prophylaxis of hepatitis B virus infection.
- 15 The present invention also contemplates a method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside or nucleotide analog or other anti-HBV agents or by isolating DNA or corresponding mRNA from the HBV and screening for a mutation in the nucleotide sequence encoding the DNA polymerase wherein the presence of the following mutations in the rt region: in one embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment, rtH90D and rtL/F108L, in even yet another embodiment, rtL157L/M, rtA181V and rtV207I; in still yet another embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; in another embodiment, rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H; in a further embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37; in yet another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63; in still another embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91; in even yet another embodiment, rtP177,

- 12 -

rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in still yet another embodiment, rtM204 and rtY203; in another embodiment, rt235, rt236, rt237, rt238 and rt239; in a further embodiment, rt247, rt248, rt249, rt250 and rt251 or combinations thereof or an equivalent one or more other mutation is indicative of a variant which  
5 exhibits a decreased sensitivity to ADV, LMV,TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof.

10

Still a further methodology comprises screening for a mutation in the nucleotide sequence encoding the envelope genes (s) wherein the presence of the following mutations in the S gene: in one embodiment, sP120T, sM125T and sT127A; in another embodiment, sT118R, sM133T, SF134V, sI195M, sS207R and sY225Y/C; in a further embodiment, sS126T,  
15 sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop in yet another embodiment, sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182Stop, sW196L, sY206H and sY225F; in still yet another embodiment, sI81M and sP214Q; in another embodiment, sF83S, sL173F and sW199L; in a further aspect, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in a further embodiment, sC69Stop/C, sC76Y, sI110V/I, sY134N,  
20 sW172Stop/W, sW196Stop and sS207R or combinations thereof or an equivalent one or more other mutation is indicative of a variant which exhibits a decreased sensitivity to ADV, LMV,TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and  
25 TFV, and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof.

Preferably, the variants are in an isolated form such that they have undergone at least one purification step away from naturally occurring body fluid. Alternatively, the variants may  
30 be maintained in isolated body fluid or may be in DNA form. The present invention also contemplates infectious molecular clones comprising the genome or parts thereof from a

- 13 -

variant HBV. The detection of HBV or its components in cells, cell lysates, cultured supernatant fluid and bodily fluid may be by any convenient means including any nucleic acid-based detection means, for example, by nucleic acid hybridization techniques or *via* one or more polymerase chain reactions (PCRs). The term "bodily fluid" includes any fluid  
5 derived from the blood, lymph, tissue or organ systems including serum, whole blood, biopsy and biopsy fluid, organ explants and organ suspension such as liver suspensions.

Another aspect of the present invention is directed to a variant HBV comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution,  
10 addition and/or deletion or a truncation compared to a surface antigen from a reference or wild type HBV and wherein an antibody generated to the reference or wild type surface antigen exhibits an altered immunological profile relative to the HBV variant. One altered profile includes a reduced capacity for neutralizing the HBV. More particularly, the surface antigen of the variant HBV exhibits an altered immunological profile compared to  
15 a pre-treatment HBV where the variant HBV is selected for by a nucleoside or nucleotide analog or other anti-HBV agents of the HBV DNA polymerase. The variant HBV of this aspect of the invention may also comprise a nucleotide sequence comprising a single or multiple nucleotide substitution, addition and/or deletion compared to a pre-treatment HBV.

20

The present invention extends to an isolated HBsAg or a recombinant form thereof or derivative or chemical equivalent thereof corresponding to the variant HBV. Generally, the HBsAg or its recombinant or derivative form or its chemical equivalent comprises an amino acid sequence with a single or multiple amino acid substitution, addition and/or  
25 deletion or a truncation compared to an HBsAg from a reference HBV and wherein an antibody directed to a reference HBV exhibits an altered immunological profile to an HBV carrying said variant HBsAg. In one embodiment, the altered immunological profile comprises a reduction in the ability to neutralize the variant HBV.

30 Another aspect of the present invention contemplates a method for detecting an agent which exhibits inhibitory activity to an HBV by generating a genetic construct comprising

- 14 -

a replication competent-effective amount of the genome from the HBV contained in a plasmid vector and then transfecting said cells with said construct, contacting the cells, before, during and/or after transfection, with the agent to be tested, culturing the cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agents; and the subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of the agent. In a preferred embodiment, the plasmid vector in a baculovirus vector and the method comprises generating a genetic construct comprising a replication competent-effective amount of the genome from the HBV contained in or fused to an amount of a baculovirus genome effective to infect cells and then infecting said cells with said construct, contacting the cells, before, during and/or after infection, with the agent to be tested, culturing the cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agent and then subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of the agent.

20

In connection with these methods, the plasmid vector may include genes encoding part or all of other viral vectors such as baculovirus vectors or adenovirus vectors (see Ren and Nassal, *J. Virol.* 75(3): 1104-1116, 2001).

25 In an alternative embodiment, the method comprises generating a continuous cell line comprising an infectious copy of the genome of the HBV in a replication competent effective amount such that said infectious HBV genome is stably integrated into said continuous cell line such as but not limited to the 2.2.15 or AD cell line, contacting the cells with the agent to be tested, culturing the cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to the agent and then subjecting the cells,

30

- 15 -

cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of the agent.

- 5 In an alternative embodiment, the present invention also contemplates a method for detecting an agent which exhibits inhibitory activity to an HBV polymerase in an *in vitro* polymerase assay. The HBV polymerase activity can be examined using established assays (Gaillard *et al.*, *Antimicrob Agents Chemother.* 46(4): 1005-1013, 2002; Xiong *et al.*, *Hepatology.* 28(6): 1669-73, 1998). The HBV polymerase may be a wild-type or reference  
10 HBV polymerase or mutant HBV polymerase.

The identification of viral variants enables the production of vaccines comprising particular recombinant viral components such as polymerases or envelope genes PreS1, PreS2, S encoding for L, M, S proteins as well as therapeutic vaccines comprising  
15 defective HBV variants. Rational drug design may also be employed to identify or generate therapeutic molecules capable of interacting with a polymerase or or envelope genes PreS1, PreS2, S encoding for L, M, S proteins or other component of the HBV. Such drugs may also have diagnostic potential. In addition, defective HBV variants may also be used as therapeutic compositions to generate an immune response against the same, similar  
20 or homologous viruses. Alternatively, antibodies generated to the HBV variants or surface components thereof may be used in passive immunization of subjects against infection by HBV variants or similar or homologous viruses. Furthermore, agents such as nucleoside or nucleotide analogs, RNAi or siRNA molecules, antisense or sense oligonucleotides, chemical or proteinaceous molecules having an ability to down-regulate the activity of a  
25 component of HBV and inhibit replication, maintenance, infection, assembly or release are contemplated by the present invention.

A summary of the abbreviations used throughout the subject specification are provided in Table 3.

- 16 -

A summary of sequence identifiers used throughout the subject specification is provided in Table 1.

**TABLE 1***Summary of sequence identifiers*

5

SEQUENCE ID NO:	DESCRIPTION
1	Formula I
2	Formula II
3	OS1 primer
4	TTA3 primer
5	JM primer
6	TTA4 primer
7	OS2 primer
8	sense primer
9	antisense primer
10	internal regions primer
11	internal regions primer
12	PC1 forward primer
13	PC2 reverse primer
14	HBV-specific molecular beacon primer
15	ILA 1 F, A-E (Figure 4)
16	ILA 2 F, A-E (Figure 4)
17	ILA 3 F, A-E (Figure 4)
18	ILA 4 F, A-E (Figure 4)
19	Pol Trans Pre 1 (Figure 5)
20	Pol Trans 2 (Figure 5)
21	Pol Trans 3 (Figure 5)
22	Pol Trans 4 (Figure 5)
23	HBsAg Trans of Pre 1 (Figure 6)

- 17 -

SEQUENCE ID NO:	DESCRIPTION
24	HBsAg Trans of 2 (Figure 6)
25	HBsAg Trans of 3 (Figure 6)
26	HBsAg Trans of 4 (Figure 6)
27	S0 (Figure 7)
28	S6 (Figure 7)
29	S8 (Figure 7)
30	S12 (Figure 7)
31	S15 (Figure 7)
32	Pol Trans S0 (Figure 8)
33	Pol Trans S6 (Figure 8)
34	Pol Trans S8 (Figure 8)
35	Pol Trans S12 (Figure 8)
36	Pol Trans S15 (Figure 8)
37	HBsAg Trans of S0 (Figure 9)
38	HBsAg Trans of S6 (Figure 9)
39	HBsAg Trans of S8 (Figure 9)
40	HBsAg Trans of S12 (Figure 9)
41	HBsAg Trans of S15 (Figure 9)
42	Nucleotide sequence Patient C (Figure 10)
43	POL Trans of Patient C (Figure 11)
44	HBsAg Trans of Patient C (Figure 12)
45	Nucleotide sequence of Patient D (Figure 13)
46	Pol Trans of Patient D (Figure 14)
47	HBsAg Trans of Patient D (Figure 15)
48	Nucleotide sequence of Patient E (Figure 16)
49	Pol Trans of Patient E (Figure 17)
50	HBsAg Trans of Patient E (Figure 18)
51	Nucleotide sequence of Patient F (Figure 20)
52	Deduced sequence of DNA polymerase of Patient F (Figure 21)

- 18 -

SEQUENCE ID NO:	DESCRIPTION
53	HBsAg Trans of Patient F (Figure 22)
54	Nucleotide sequence of Patient G (Figure 23)
55	Deduced sequence of DNA polymerase of Patient G (Figure 24)
56	HBsAg Trans of Patient G (Figure 25)
57	Nucleotide sequence of Patient H (Figure 26)
58	Deduced sequence of DNA polymerase of Patient H (Figure 27)
59	HBsAg Trans of Patient H (Figure 28)



**TABLE 2**  
*Single and three letter amino acid abbreviations*

<b>Amino Acid</b>	<b>Three-letter Abbreviation</b>	<b>One-letter symbol</b>
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic acid	Asp	D
Cysteine	Cys	C
Glutamine	Gln	Q
Glutamic acid	Glu	E
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	The	T
Tryptophan	Trp	W
Tyrosine	-Tyr	Y
Valine	Val	V
Any residue	Xaa	X

- 20 -

A list of abbreviations used throughout the subject specification are provided in Table 3.

**TABLE 3**  
*Abbreviations*

5

ABBREVIATION	DESCRIPTION
3TC	(LMV); (-)- $\beta$ -2'-deoxy-3'-thiacytidine
ADV	adefovir dipivoxil
DAPD	diaminopurine dioxalone
DXG	dioxolane guanine
ETV	entecavir
FAM	famciclovir
FCV	famciclovir
FTC	emtricitabine
HBIG	hepatitis B immunoglobulin
HBsAg	hepatitis B surface antigen
HBV	hepatitis B virus
LMV	lamivudine
PMEA	9-[phosphonyl-methoxyethyl]-adenine; adefovir
PMPA	9-R-(2-phosphonomethoxypropyl)adenine
RNase	ribonuclease
rt ("rt" before "Xaa <sub>1</sub> nXaa <sub>2</sub> " means reverse transcriptase)	reverse transcriptase
s (as used in a mutation, e.g. sF134V)	envelope gene
TFV	tenofovir disoproxil fumarate
YMDD	Tyr Met Asp Asp-a motif in the polymerase protein; where the Met residue is designated residue number 204 of the reverse transcriptase

## BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a diagrammatic representation showing the partially double stranded DNA HBV genome showing the overlapping open reading frames encoding surface (S), core (C), polymerase (P) and X gene.

Figure 2 is a diagrammatic representation of the chemical structure of ADV.

Figure 3 is a diagrammatic representation of a computer system for determining the potency value ( $P_A$ ) of a variant HBV.

Figure 4 is a representation showing comparison of the HBV nucleotide sequence encoding the catalytic region of the polymerase gene in sequential samples from Patient A during ADV treatment.

Figure 5 is a representation showing comparison of the deduced amino acid sequence of the catalytic region of the polymerase gene in sequential samples from Patient A during ADV therapy.

Figure 6 is a representation showing comparison of the deduced amino acid sequence of the envelope gene in sequential samples from Patient A during ADV therapy.

Figure 7 is a representation showing comparison of the HBV nucleotide sequence encoding the catalytic region of the polymerase gene in sequential samples from Patient B during ADV and LMV treatment.

Figure 8 is a representation showing comparison of the deduced amino acid sequence of the catalytic region of the polymerase gene in sequential samples from Patient B during ADV and LMV therapy.

- 22 -

Figure 9 is a representation showing comparison of the deduced amino acid sequence of the envelope gene in sequential samples from Patient B during ADV and LMV therapy.

Figure 10 is a representation showing comparison of the HBV nucleotide sequence encoding the catalytic region of the polymerase gene in sequential samples from Patient C during ADV treatment.

Figure 11 is a representation showing comparison of the deduced amino acid sequence of the catalytic region of the polymerase gene in sequential samples from Patient C during ADV therapy.

Figure 12 is a representation showing comparison of the deduced amino acid sequence of the envelope gene in sequential samples from Patient C during ADV therapy.

Figure 13 is a representation showing comparison of the HBV nucleotide sequence encoding the catalytic region of the polymerase gene in sequential samples from Patient D during ADV treatment.

Figure 14 is a representation showing comparison of the deduced amino acid sequence of the catalytic region of the polymerase gene in sequential samples from Patient D during ADV therapy.

Figure 15 is a representation showing comparison of the deduced amino acid sequence of the envelope gene in sequential samples from Patient D during ADV therapy.

Figure 16 is a representation showing comparison of the HBV nucleotide sequence encoding the catalytic region of the polymerase gene in sequential samples from Patient E during ADV treatment.

- 23 -

**Figure 17** is a representation showing comparison of the deduced amino acid sequence of the catalytic region of the polymerase gene in sequential samples from Patient E during ADV therapy.

- 5 **Figure 18** is a representation showing comparison of the deduced amino acid sequence of the envelope gene in sequential samples from Patient E during ADV therapy.

**Figure 19** is a diagrammatic representation of a system used to carry out the instructions encoded by the storage medium.

10

**Figure 20** is a representation showing the nucleotide sequence of envelope/rt region of an HBV isolated from Patient F having ADV therapy.

- 15 **Figure 21** is a representation showing the deduced amino acid sequence of DNA polymerase encoded by the nucleotide sequence shown in Figure 20.

**Figure 22** is a representation showing the deduced amino acid sequence of HBsAg encoded by the nucleotide sequence shown in Figure 20.

- 20 **Figure 23** is a representation showing the nucleotide sequence of envelope/rt region of an HBV isolated from Patient G having ADV therapy.

**Figure 24** is a representation showing the deduced amino acid sequence of DNA polymerase encoded by the nucleotide sequence shown in Figure 23.

25

**Figure 25** is a representation showing the deduced amino acid sequence of HBsAg encoded by the nucleotide sequence shown in Figure 23.

- 30 **Figure 26** is a representation showing the nucleotide sequence of envelope/rt region of an HBV isolated from Patient H having ADV therapy.

- 24 -

**Figure 27** is a representation showing the deduced amino acid sequence of DNA polymerase encoded by the nucleotide sequence shown in Figure 26.

**Figure 28** is a representation showing the deduced amino acid sequence of HBsAg  
5 encoded by the nucleotide sequence shown in Figure 26.

- 25 -

## DETAILED DESCRIPTION OF THE INVENTION

The present invention is predicated in part on the identification and isolation of nucleoside or nucleotide analog-resistant variants of HBV following treatment of patients with either ADV or LMV or more particularly ADV and LMV, or optionally other nucleoside analogs or nucleotide analogs or other anti-HBV agents such as TFV or FTC. In particular, ADV or ADV and LMV treated patients gave rise to variants of HBV exhibiting decreased or reduced sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV. . Reference herein to "decreased" or "reduced" in relation to sensitivity to ADV and/or LMV and/or FTC and/or TFV includes and encompasses a complete or substantial resistance to the nucleoside or nucleotide analog or other anti-HBV agents as well as partial resistance and includes a replication rate or replication efficiency which is more than a wild-type in the presence of a nucleoside or nucleotide analog or other anti-HBV agents. In one aspect, this is conveniently measured by an increase in viral load during treatment, or alternatively, there is no substantial decrease in HBV DNA viral load from pre-treatment HBV DNA levels during treatment (i.e., non-response to treatment).

Before describing the present invention in detail, it is to be understood that unless otherwise indicated, the subject invention is not limited to specific formulations of components, manufacturing methods, dosage regimens, or the like, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

It must be noted that, as used in the subject specification, the singular forms "a", "an" and "the" include plural aspects unless the context clearly dictates otherwise. Thus, for example, reference to "a nucleoside or nucleotide analog" includes a single analog, as well as two or more analogs; reference to "an HBV variant" includes reference to two or more HBV variants; and so forth.

- 26 -

In describing and claiming the present invention, the following terminology is used in accordance with the definitions set forth below.

5 The terms "analog", "compound", "active agent", "pharmacologically active agent", "medicament", "active" and "drug" are used interchangeably herein to refer to a chemical compound that induces a desired effect such as inhibit viral replication, infection, maintenance, assembly and/or the function of an enzyme such as HBV DNA polymerase. The terms also encompass pharmaceutically acceptable and pharmacologically active  
10 ingredients of those active agents specifically mentioned herein including but not limited to salts, esters, amides, prodrugs, active metabolites, analogs and the like. When the terms "analog", "compound", "active agent", "pharmacologically active agent", "medicament", "active" and "drug" are used, then it is to be understood that this includes the active agent *per se* as well as pharmaceutically acceptable, pharmacologically active salts, esters,  
15 amides, prodrugs, metabolites, analogs, etc.

The present invention contemplates, therefore, compounds useful in inhibiting HBV replication, infection, maintenance, assembly and/or the function of an enzyme such as HBV DNA polymerase. Reference to an "analog", "compound", "active agent",  
20 "pharmacologically active agent", "medicament", "active" and "drug" such as ADV, LMV, FTC and/or TFV includes combinations of two or more actives such as ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV. A  
25 "combination" also includes a two-part or more such as a multi-part anti-HBV therapeutic composition where the agents are provided separately and given or dispensed separately or admixed together prior to dispensation.

The terms "effective amount" and "therapeutically effective amount" of an agent as used  
30 herein mean a sufficient amount of the agent to provide the desired therapeutic or physiological effect of inhibiting HBV replication, infection, maintenance, assembly



- 27 -

and/or the function of an enzyme such as HBV DNA polymerase. Furthermore, an "effective HBV-inhibiting amount" or "effective symptom-ameliorating amount" of an agent is a sufficient amount of the agent to directly or indirectly inhibit replication, infection, maintenance, assembly and/or the function of an enzyme such as HBV DNA polymerase. Undesirable effects, e.g. side effects, are sometimes manifested along with the desired therapeutic effect; hence, a practitioner balances the potential benefits against the potential risks in determining what is an appropriate "effective amount". The exact amount required will vary from subject to subject, depending on the species, age and general condition of the subject, mode of administration and the like. Thus, it may not be possible to specify an exact "effective amount". However, an appropriate "effective amount" in any individual case may be determined by one of ordinary skill in the art using only routine experimentation.

By "pharmaceutically acceptable" carrier, excipient or diluent is meant a pharmaceutical vehicle comprised of a material that is not biologically or otherwise undesirable, i.e. the material may be administered to a subject along with the selected active agent without causing any or a substantial adverse reaction. Carriers may include excipients and other additives such as diluents, detergents, coloring agents, wetting or emulsifying agents, pH buffering agents, preservatives, and the like.

Similarly, a "pharmacologically acceptable" salt, ester, emide, prodrug or derivative of a compound as provided herein is a salt, ester, amide, prodrug or derivative that this not biologically or otherwise undesirable.

The terms "treating" and "treatment" as used herein refer to reduction in severity and/or frequency of symptoms, elimination of symptoms and/or underlying cause, prevention of the occurrence of symptoms and/or their underlying cause, and improvement or remediation of damage in relation to HBV infection. Thus, for example, "treating" a patient involves prevention of HBV infection as well as treatment of a clinically HBV symptomatic individual by inhibiting HBV replication, infection, maintenance, assembly and/or the function of an enzyme such as HBV DNA polymerase. Thus, for example, the

- 28 -

present method of "treating" a patient with HBV infection or with a propensity for one to develop encompasses both prevention of HBV infection as well as treating HBV infection or symptoms thereof. In any event, the present invention contemplates the treatment or prophylaxis of HBV infection.

5

"Patient" as used herein refers to an animal, preferably a mammal and more preferably a primate including a lower primate and even more preferably, a human who can benefit from the formulations and methods of the present invention. A patient regardless of whether a human or non-human animal may be referred to as an individual, subject,  
10 animal, host or recipient. The compounds and methods of the present invention have applications in human medicine, veterinary medicine as well as in general, domestic or wild animal husbandry. For convenience, an "animal" includes an avian species such as a poultry bird (including ducks, chicken, turkeys and geese), an aviary bird or game bird. The condition in a non-human animal may not be a naturally occurring HBV infection but  
15 HBV-like infection may be induced.

As indicated above, the preferred animals are humans, non-human primates such as marmosets, baboons, orangatangs, lower primates such as tupia, livestock animals, laboratory test animals, companion animals or captive wild animals. A human is the most  
20 preferred target. However, non-human animal models may be used.

Examples of laboratory test animals include mice, rats, rabbits, guinea pigs and hamsters. Rabbits and rodent animals, such as rats and mice, provide a convenient test system or animal model as do primates and lower primates. Livestock animals include sheep, cows,  
25 pigs, goats, horses and donkeys. Non-mammalian animals such as avian species, zebrafish, amphibians (including cane toads) and *Drosophila* species such as *Drosophila melanogaster* are also contemplated. Instead of a live animal model, a test system may also comprise a tissue culture system.

30 Accordingly, one aspect of the present invention is directed to an isolated HBV variant wherein said variant comprises a nucleotide mutation in a gene encoding a DNA

- 29 -

polymerase resulting in at least one amino acid addition, substitution and/or deletion to said DNA polymerase and wherein said variant exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, 5 TFV and FTC and LMV, or ADV and LMV and FTC, ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof.

HBV is a member of the Hepdnaviridae that includes also avian hepatitis viruses such as 10 Duck hepatitis B virus (DHBV) and hepatitis viruses from mammals such as woodchuck hepatitis virus (WHV). These viruses have similarity to HBV and may be used in *in vitro* and *in vivo* or animal model systems to investigate the equivalent HBV mutants and anti-viral sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or 15 ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV,

An "anti-HBV agent" includes a nucleoside or nucleotide analog, protein, chemical compound, RNA or DNA or RNAi or siRNA oligonucleotide.

20

Preferably, the decreased sensitivity is in respect of ADV. Alternatively, the decreased sensitivity is in respect of LMV. Alternatively, the decreased sensitivity is in respect of TFV. Alternatively, the decreased sensitivity is in respect of FTC. Alternatively, the decreased sensitivity is in respect of ADV and LMV. Alternatively, the decreased 25 sensitivity is in respect of ADV and TFV. Alternatively, the decreased sensitivity is in respect of LMV and TFV. Alternatively, the decreased sensitivity is in respect of ADV and FTC. Alternatively, the decreased sensitivity is in respect to FTC and TFV. Alternatively, the decreased sensitivity is in respect of FTC and LMV. Alternatively, the decreased sensitivity is in respect of ADV and LMV and TFV. Alternatively, the decreased sensitivity 30 is in respect to ADV and TFV and FTC. Alternatively, the decreased sensitivity is in respect to LMV and TFV and FTC. Alternatively, the decrease sensitivity is in respect of ADV and

- 30 -

LMV and FTC. Alternatively, the decreased sensitivity is in respect of ADV and FTC and TFV and LMV.

Reference herein to "anti-HBV agents" includes nucleoside and nucleotide analogs as well  
5 as immunological reagents (e.g. antibodies to HBV surface components) and chemical, proteinaceous and nucleic acid agents which inhibit or otherwise interfere with viral replication, maintenance, infection, assembly or release. Reference herein to "nucleic acid" includes reference to a sense or antisense molecule, RNA or DNA, oligonucleotides and RNAi and siRNA molecules and complexes containing same.

10

In addition to a mutation in the gene encoding DNA polymerase, due to the overlapping nature of the HBV genome (Figure 1), a corresponding mutation may also occur in the gene encoding the S gene encoding the surface antigen (HBsAg) resulting in reduced interactivity of immunological reagents such as antibodies and immune cells to HBsAg.  
15 The reduction in the interactivity of immunological reagents to a viral surface component generally includes the absence of immunological memory to recognize or substantially recognize the viral surface component. The present invention extends, therefore, to an HBV variant exhibiting decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or  
20 ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, and/or ADV and FTC and LMV and TFV or a reduced interactivity of an immunological reagent to HBsAg wherein the variant is selected for following ADV and/or LMV combination or sequential treatment. The term "sequential" in this respect means ADV followed by LMV and/or TFV, and /or FTC; LMV followed by ADV and/or  
25 TFV, and /or FTC, or multiple sequential administrations of each of ADV, LMV and/or TFV, and /or FTC.

A viral variant may, therefore, carry mutation only in the DNA polymerase gene or both in the DNA polymerase gene and the S gene. The term "mutation" is to be read in its broadest  
30 context and includes multiple mutations.

- 31 -

The present invention extends to a mutation and any domain of the HBV DNA polymerase and in particular regions F and G, and domains A through to E provided said mutation leads to decreased sensitivity to ADV and/ or LMV and/or TFV or combinations thereof. Regions F and G of the HBV DNA polymerase is defined by the amino acid sequence set forth in Formula I below [SEQ ID NO:1]:

### FORMULA I

L, X<sub>1</sub>, X<sub>2</sub>, D, W, G, P, C, X<sub>3</sub>, X<sub>4</sub>, H, G, X<sub>5</sub>, H, X<sub>6</sub>, I, R, B<sub>7</sub>, P, R, T, P, X<sub>8</sub>, R, V, X<sub>9</sub>, G, G,  
 10 V, F, L, V, D, K, N, P, H, N, T, X<sub>10</sub>, E, S, X<sub>11</sub>, L, X<sub>12</sub>, V, D, F, S, Q, F, S, R, G, X<sub>13</sub>, X<sub>14</sub>,  
 X<sub>15</sub>, V, S, W, P, K, F, A, V, P, N, L, X<sub>16</sub>, S, L, T, N, L, L, S\*

wherein:

- 15 X<sub>1</sub> is L, or R, or I
- X<sub>2</sub> is E, or D
- X<sub>3</sub> is T, or D, or A, or N, or Y
- X<sub>4</sub> is E, or D
- X<sub>5</sub> is E, or K, or Q
- 20 X<sub>6</sub> is H, or R, or N,
- X<sub>7</sub> is I, or T
- X<sub>8</sub> is A, or S
- X<sub>9</sub> is T or R
- X<sub>10</sub> is A, or T, or S
- 25 X<sub>11</sub> is R, or T
- X<sub>12</sub> is V, or G
- X<sub>13</sub> is S, or L, or T, or N, or V
- X<sub>14</sub> is T, or S, or H, or Y
- X<sub>15</sub> is R, or H, or K, or Q
- 30 X<sub>16</sub> is Q, or P;

- 32 -

and wherein S\* is designated as amino acid 74.

In this specification, reference is particularly made to the conserved regions of the DNA polymerase as defined by domains A to E. Regions A to E are defined by the amino acid  
 5 sequence set forth in Formula II below [SEQ ID NO:2] (and in Australian Patent No. 734831):

### FORMULA II

10 S X<sub>1</sub> L S W L S L D V S A A F Y H X<sub>2</sub> P L H P A A M P H L L X<sub>3</sub> G S S G L X<sub>4</sub> R Y V  
 A R L S S X<sub>5</sub> S X<sub>6</sub> X<sub>7</sub> X N X<sub>8</sub> Q X<sub>9</sub> X<sub>10</sub> X X X X<sub>11</sub> L H X<sub>12</sub> X<sub>13</sub> C S R X<sub>14</sub> L Y V S L X<sub>15</sub>  
 L L Y X<sub>16</sub> T X<sub>17</sub> G X<sub>18</sub> K L H L X<sub>19</sub> X<sub>20</sub> H P I X<sub>21</sub> L G F R K X<sub>22</sub> P M G X<sub>23</sub> G L S P F L  
 L A Q F T S A I X<sub>24</sub> X<sub>25</sub> X<sub>26</sub> X<sub>27</sub> X<sub>28</sub> R A F X<sub>29</sub> H C X<sub>30</sub> X<sub>31</sub> F X<sub>32</sub> Y M\* D D X<sub>33</sub> V L G A  
 X<sub>34</sub> X<sub>35</sub> X<sub>36</sub> X<sub>37</sub> H X<sub>38</sub> E X<sub>39</sub> L X<sub>40</sub> X<sub>41</sub> X<sub>42</sub> X<sub>43</sub> X<sub>44</sub> X<sub>45</sub> X<sub>46</sub> L L X<sub>47</sub> X<sub>48</sub> G I H L N P X<sub>49</sub> K  
 15 T K R W G Y S L N F M G Y X<sub>50</sub> I G

wherein:

- X is any amino acid
- 20 X<sub>1</sub> is N or D;
- X<sub>2</sub> is I or P;
- X<sub>3</sub> is I or V;
- X<sub>4</sub> is S or D;
- X<sub>5</sub> is T or N;
- 25 X<sub>6</sub> is R or N;
- X<sub>7</sub> is N or I;
- X<sub>8</sub> is N or Y or H;
- X<sub>9</sub> is H or Y;
- X<sub>10</sub> is G or R;
- 30 X<sub>11</sub> is D or N;
- X<sub>12</sub> is D or N;

- 33 -

- X<sub>13</sub> is S or Y;  
X<sub>14</sub> is N or Q;  
X<sub>15</sub> is L or M;  
X<sub>16</sub> is K or Q;  
5 X<sub>17</sub> is Y or F;  
X<sub>18</sub> is R or W;  
X<sub>19</sub> is Y or L;  
X<sub>20</sub> is S or A;  
X<sub>21</sub> is I or V;  
10 X<sub>22</sub> is I or L;  
X<sub>23</sub> is V or G;  
X<sub>24</sub> is C or L;  
X<sub>25</sub> is A or S;  
X<sub>26</sub> is V or M;  
15 X<sub>27</sub> is V or T;  
X<sub>28</sub> is R or C;  
X<sub>29</sub> is F or P;  
X<sub>30</sub> is L or V;  
X<sub>31</sub> is A or V;  
20 X<sub>32</sub> is S or A;  
X<sub>33</sub> is V or L or M;  
X<sub>34</sub> is K or R;  
X<sub>35</sub> is S or T;  
X<sub>36</sub> is V or G;  
25 X<sub>37</sub> is Q or E;  
X<sub>38</sub> is L or S or R;  
X<sub>39</sub> is S or F;  
X<sub>40</sub> is F or Y;  
X<sub>41</sub> is T or A;  
30 X<sub>42</sub> is A or S;  
X<sub>43</sub> is V or I;

- 34 -

- X<sub>44</sub> is T or C;  
X<sub>45</sub> is N or S;  
X<sub>46</sub> is F or V;  
X<sub>47</sub> is S or D;  
5 X<sub>48</sub> is L or V;  
X<sub>49</sub> is N or Q;  
X<sub>50</sub> is V or I; and  
M\* is amino acid 204;

- 10 and wherein the first S is designated as amino acid 75.

Preferably, the mutation results in an altered amino acid sequence in any one or more of domains F and G, and domains A through to E or regions proximal thereto of the HBV DNA polymerase.

15

- Another aspect of the present invention provides an HBV variant comprising a mutation in an overlapping open reading frame in its genome wherein said mutation is in a region defined by one or more of domains F and G, and domains A through to E of HBV DNA polymerase and wherein said variant exhibits decreased sensitivity to ADV, LMV, TFV, or  
20 FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents.

- 25 In a related embodiment, there is provided an HBV variant comprising a mutation in the nucleotide sequence encoding a DNA polymerase resulting in an amino acid addition, substitution and/or deletion in said DNA polymerase in one or more amino acids as set forth in Formula I [SEQ ID NO:1] and/or Formula II [SEQ ID NO:2]:



- 35 -

**FORMULA I**

L, X<sub>1</sub>, X<sub>2</sub>, D, W, G, P, C, X<sub>3</sub>, X<sub>4</sub>, H, G, X<sub>5</sub>, H, X<sub>6</sub>, I, R, X<sub>7</sub>, P, R, T, P, X<sub>8</sub>, R, V, X<sub>9</sub>, G, G,  
 V, F, L, V, D, K, N, P, H, N, T, X<sub>10</sub>, E, S, X<sub>11</sub>, L, X<sub>12</sub>, V, D, F, S, Q, F, S, R, G, X<sub>13</sub>, X<sub>14</sub>,  
 5 X<sub>15</sub>, V, S, W, P, K, F, A, V, P, N, L, X<sub>16</sub>, S, L, T, N, L, L, S\*

wherein:

- X<sub>1</sub> is L, or R, or I  
 10 X<sub>2</sub> is E, or D  
 X<sub>3</sub> is T, or D, or A, or N, or Y  
 X<sub>4</sub> is E, or D  
 X<sub>5</sub> is E, or K, or Q  
 X<sub>6</sub> is H, or R, or N,  
 15 X<sub>7</sub> is I, or T  
 X<sub>8</sub> is A, or S  
 X<sub>9</sub> is T or R  
 X<sub>10</sub> is A, or T, or S  
 X<sub>11</sub> is R, or T  
 20 X<sub>12</sub> is V, or G  
 X<sub>13</sub> is S, or I, or T, or N, or V  
 X<sub>14</sub> is T, or S, or H, or Y  
 X<sub>15</sub> is R, or H, or K, or Q  
 X<sub>16</sub> is Q, or P;

25

and

**FORMULA II**

30 S X<sub>1</sub> L S W L S L D V S A A F Y H X<sub>2</sub> P L H P A A M P H L L X<sub>3</sub> G S S G L X<sub>4</sub> R Y V  
 A R L S S X<sub>5</sub> S X<sub>6</sub> X<sub>7</sub> X N X<sub>8</sub> Q X<sub>9</sub> X<sub>10</sub> X X X X<sub>11</sub> L H X<sub>12</sub> X<sub>13</sub> C S R X<sub>14</sub> L Y V S L X<sub>15</sub>

- 36 -

LLYX<sub>16</sub>TX<sub>17</sub>GX<sub>18</sub>KLHLX<sub>19</sub>X<sub>20</sub>HPIX<sub>21</sub>LGFRKX<sub>22</sub>PMGX<sub>23</sub>GLSPFL  
 LAQFTSAIX<sub>24</sub>X<sub>25</sub>X<sub>26</sub>X<sub>27</sub>X<sub>28</sub>RAFX<sub>29</sub>HGX<sub>30</sub>X<sub>31</sub>FX<sub>32</sub>YM<sup>\*</sup>DDX<sub>33</sub>VLGA  
 X<sub>34</sub>X<sub>35</sub>X<sub>36</sub>X<sub>37</sub>HX<sub>38</sub>EX<sub>39</sub>LX<sub>40</sub>X<sub>41</sub>X<sub>42</sub>X<sub>43</sub>X<sub>44</sub>X<sub>45</sub>X<sub>46</sub>LLX<sub>47</sub>X<sub>48</sub>GIHLNPX<sub>49</sub>K  
 TKRWGYSLNFMGYX<sub>50</sub>IG

5

wherein:

- X is any amino acid
- X<sub>1</sub> is N or D;
- 10 X<sub>2</sub> is I or P;
- X<sub>3</sub> is I or V;
- X<sub>4</sub> is S or D;
- X<sub>5</sub> is T or N;
- X<sub>6</sub> is R or N;
- 15 X<sub>7</sub> is N or I;
- X<sub>8</sub> is N or Y or H;
- X<sub>9</sub> is H or Y;
- X<sub>10</sub> is G or R;
- X<sub>11</sub> is D or N;
- 20 X<sub>12</sub> is D or N;
- X<sub>13</sub> is S or Y;
- X<sub>14</sub> is N or Q;
- X<sub>15</sub> is L or M;
- X<sub>16</sub> is K or Q;
- 25 X<sub>17</sub> is Y or F;
- X<sub>18</sub> is R or W;
- X<sub>19</sub> is Y or L;
- X<sub>20</sub> is S or A;
- X<sub>21</sub> is I or V;
- 30 X<sub>22</sub> is I or L;
- X<sub>23</sub> is V or G;

- 37 -

- X<sub>24</sub> is C or L;  
X<sub>25</sub> is A or S;  
X<sub>26</sub> is V or M;  
X<sub>27</sub> is V or T;  
5 X<sub>28</sub> is R or C;  
X<sub>29</sub> is F or P;  
X<sub>30</sub> is L or V;  
X<sub>31</sub> is A or V;  
X<sub>32</sub> is S or A;  
10 X<sub>33</sub> is V or L or M;  
X<sub>34</sub> is K or R;  
X<sub>35</sub> is S or T;  
X<sub>36</sub> is V or G;  
X<sub>37</sub> is Q or E;  
15 X<sub>38</sub> is L or S or R;  
X<sub>39</sub> is S or F;  
X<sub>40</sub> is F or Y;  
X<sub>41</sub> is T or A;  
X<sub>42</sub> is A or S;  
20 X<sub>43</sub> is V or I;  
X<sub>44</sub> is T or C;  
X<sub>45</sub> is N or S;  
X<sub>46</sub> is F or V;  
X<sub>47</sub> is S or D;  
25 X<sub>48</sub> is L or V;  
X<sub>49</sub> is N or Q;  
X<sub>50</sub> is V or I; and  
M\* is amino acid 204;

- 30 and wherein S\* in Formula I is designated as amino acid 74 and the first S in Formula II is designated as amino acid 75;

- 38 -

and wherein said variant exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, 5 ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof. Preferably, the decreased sensitivity is to ADV or to both ADV and LMV or to one or both of ADV and/or LMV and/or TFV and /or FTC.

10 Another preferred aspect of the present invention contemplates an HBV variant comprising a mutation in the nucleotide sequence encoding HBsAg resulting in an amino acid addition, substitution and/or deletion in said HBsAg in a region corresponding to the amino acid sequence set forth in Formulae I and II wherein said variant exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and 15 TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof.

20 More particularly, the present invention provides a variant HBV comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or a truncation compared to a surface antigen from a reference or wild-type HBV and wherein an antibody generated to the reference or wild-type surface antigen exhibits reduced capacity for neutralizing said HBV variant, said variant selected 25 by exposure of a subject to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof.

30

- 39 -

The term "combination therapy" means that both combinations of ADV, LMV, FTC and/or TFV are co-administered in the same composition or simultaneously in separate compositions. The term "sequential therapy" means that the two agents are administered within seconds, minutes, hours, days or weeks of each other and in either order. Sequential  
5 therapy also encompasses completing a therapeutic course with one or other of ADV, LMV, FTC or TFV and then completing a second or third or subsequent therapeutic courses with the other of ADV, LMV, FTC or TFV.

Accordingly, another aspect of the present invention contemplates an HBV variant  
10 comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV therapy or therapy by one or more other  
15 nucleoside or nucleotide analogs or other anti-HBV agents.

Another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the  
20 surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to LMV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Yet another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure  
25 of a subject to FTC therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.  
30

- 40 -

Still another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

10 Even yet another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV and LMV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Even still another aspect of of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

25 A further aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure

30

- 41 -

of a subject to LMV and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV and FTC therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Yet another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to TFV and FTC therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Still another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to FTC and LMV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Even yet another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile

- 42 -

compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV, LMV and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

- 5 Even still another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure  
10 of a subject to ADV, LMV and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

A further aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid  
15 substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV, LMV and FTC therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

20

Another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to  
25 the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to FTC, LMV and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Yet another aspect of the present invention contemplates an HBV variant comprising a  
30 surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and



- 43 -

wherein the surface antigen of the variant HBV exhibits an altered immunological profile compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV, FTC and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

5

Still yet another aspect of the present invention contemplates an HBV variant comprising a surface antigen having an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or truncation compared to the pretreatment HBV and wherein the surface antigen of the variant HBV exhibits an altered immunological profile  
10 compared to the pretreatment HBV where the said variant HBV is selected for by exposure of a subject to ADV, LMV, FTC and TFV therapy or therapy by one or more other nucleoside or nucleotide analogs or other anti-HBV agents.

Preferably, the variants are in isolated form such that they have undergone at least one  
15 purification step away from naturally occurring body fluid. Alternatively, the variants may be maintained in isolated body fluid or may be in DNA form. The present invention also contemplates infectious molecular clones comprising the genome or parts thereof from a variant HBV. Furthermore, the present invention provides isolated components from the variant HBVs such as but not limited to an isolated HBsAg. Accordingly, the present  
20 invention provides an isolated HBsAg or a recombinant form thereof or derivative or chemical equivalent thereof, said HBsAg being from a variant HBV selected by exposure of a subject to one or more of ADV, LMV, FTC and/or TFV or optionally one or more nucleoside or nucleotide analogs or other anti-HBV agents.

25 More particularly, yet another aspect of the present invention is directed to an isolated variant HBsAg or a recombinant or derivative form thereof or a chemical equivalent thereof wherein said HBsAg or its recombinant or derivative form or its chemical equivalent exhibits an altered immunological profile compared to an HBsAg from a reference HBV, said HBsAg being from a variant HBV selected by exposure of a subject  
30 to one or more of ADV, LMV, FTC and/or TFV or optionally one or more nucleoside or nucleotide analogs or other anti-HBV agents.

- 44 -

Even more particularly, the present invention provides an isolated variant HBsAg or a recombinant or derivative form thereof or a chemical equivalent thereof wherein said HBsAg or its recombinant or derivative form or its chemical equivalent comprises an amino acid sequence with a single or multiple amino acid substitution, addition and/or deletion or a truncation compared to an HBsAg from a reference HBV and wherein a neutralizing antibody directed to a reference HBV exhibits no or reduced neutralising activity to an HBV carrying said variant HBsAg, said HBsAg being from a variant HBV selected by exposure of a subject to one or more of ADV, LMV, FTC and/or TFV or optionally one or more nucleoside or nucleotide analogs or other anti-HBV agents.

Preferred mutations in the HBV DNA polymerase include variants selected from patients with HBV recurrence following ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV treatment. Nucleoside or nucleotide analog or other anti-HBV agents treatment may occur in relation to a transplantation procedure (e.g. bone marrow transplantation (BMT) or OLT) or following treatment of patients diagnosed with hepatitis. Following selection of variants, viral loads are obtainable at levels similar to pre-treatment levels or are increasing while on therapy.

Preferred mutations include, in one embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment, rtH90D, and rtL/F108L; in even yet another embodiment, rtL157L/M, rtA181V and rtV207I; in still yet another embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; in another embodiment, rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and

- 45 -

rtN238N/H; in a further embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37; in yet another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63; in still another embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtL91; in even yet another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in still yet another  
 5 embodiment, rtM204 and rtY203; in another embodiment, rt235, rt236, rt237, rt238 and rt239; in a further embodiment, rt247, rt248, rt249, rt250 and rt251; in yet another embodiment,

K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 10 P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 15 F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
 V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 20 S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
 Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
 25 P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
 30 Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;

- 46 -

- T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 5 N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 10 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 15 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E; and  
 20 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y.

Reference above to "deletion" means that the first mentioned amino acid before the residue number has been deleted.

- 25 Such HBV variants are proposed to exhibit a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination  
 30 thereof. It should be noted that the nomenclature system for amino acid positions is based on the methionine residues in the YMDD motif being designated codon rtM204. This

- 47 -

numbering system is different to that in Australian Patent No. 734831 where the methionine residue in the YMDD motif within the polymerase gene is designated codon 550. In this regard, rtL180M and rtM204V correspond to L526M and M550V, respectively, in Australian Patent No. 734831. Corresponding mutations may also occur in  
5 envelope genes such as in one or more of PreS1, PreS2 and S. The mutations in S gene encoding HBsAg at sT118R, sP120T, sS143S/T, sD144A or sI195M also result in mutation in the in the polymerase gene rtY126C, rtT128N, rtF151S/F or rtM204V respectively.

10 Another potential mode of action of ADV and other acyclic nucleoside phosphonates is that of immune stimulation (Calio *et al.*, *Antiviral Res.* 23: 77-89, 1994). A number of mutations resulted in changes in the envelope gene detected in HBV variants which may be associated with immune escape. These changes include sT118R, sP120T, sS126T, sM133T, sM133L/M, sF134V, sS143S/T, sD144A, sG145A and/or sW172STOP.

15 HBV encoding the mutation at codon sG145R is a well characterized vaccine escape mutant, although the envelope protein from HBV encoding the mutation at sG145A does not have the same antigen/antibody binding characteristics as the sG145R. This mutation was detected in HBV isolated from patient C in conjunction with mutations at codons 143  
20 and 144.

The identification of the variants of the present invention permits the generation of a range of assays to detect such variants. The detection of such variants may be important in identifying resistant variants to determine the appropriate form of chemotherapy and/or to  
25 monitor vaccination protocols, or develop new or modified vaccine preparations.

Still another aspect of the present invention contemplates a method for determining the potential for an HBV to exhibit reduced sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV,  
30 or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other

- 48 -

nucleoside or nucleotide analogs or other anti-HBV agents, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV DNA polymerase resulting in at least one amino acid substitution, deletion and/or addition in any one or more of domains F and G, and A domains through to E or a region proximal thereto of said DNA polymerase and associated with resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents wherein the presence of such a mutation is an indication of the likelihood of resistance to said ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents.

Preferably, the assay detects one or more of the following mutations: in one embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment, rtH90D and rtL/F108L; in even yet another embodiment, sP120T, sM125T and sT127A; in still yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in another embodiment, sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop; in a further embodiment, sN40S, sC69STOP, sM75I, sL88P, sT118A, sW182Stop, sW196L, sY206H and sY225F; in yet another embodiment, sI181M and sP214Q; in still another embodiment, sF83S, sL173F and sW199L; in yet another embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in still another embodiment, sC69Stop/C, sC76Y, sI110V/I, sY134N, sW172Stop/W, sW196Stop, sS207R; in even still

- 49 -

another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37); in another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63); in a further embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91); in yet another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in still another embodiment, rtM204 and  
5 rtY203; in even yet another embodiment, rt235, rt236, rt237, rt238 and rt239 and in even still another embodiment, rt247, rt248, rt249, rt250 and rt251 and in another embodiment, K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletionF;  
10 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
15 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
20 A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
25 F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
30 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;

- 50 -

- Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
5 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
10 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
15 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or combinations thereof or an  
20 equivalent one or more other mutation is indicative of a variant wherein said variant  
exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV  
and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and  
LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and  
FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or  
25 nucleotide analogs or other anti-HBV agents or combination thereof.

Accordingly, another aspect of the present invention produces a method for determining  
whether an HBV strain exhibits reduced sensitivity to a nucleoside or nucleotide analog or  
other anti-HBV agents, said method comprising isolating DNA or corresponding mRNA  
30 from said HBV and screening for a mutation in the nucleotide sequence encoding the DNA  
polymerase and/or a corresponding region of the S gene, wherein the presence of a



- 51 -

mutation selected from, in one embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y,

5 rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment, rtH90D and rtL/F108L; in even yet another embodiment, sP120T, sM125T and sT127A; in still yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in another embodiment, sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop;

10 in a further embodiment, sN40S, sC69STOP, sM75I, sL88P, sT118A, sW182Stop, sW196L, sY206H and sY225F; in yet another embodiment, sI81M and sP214Q; in still another embodiment, sF83S, sL173F and sW199L; in yet another embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in still another embodiment, sC69Stop/C, sC76Y, sI110V/I, sY134N, sW172Stop/W, sW196Stop,

15 sS207R; in even still another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37); in another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63); in a further embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91); in yet another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in still another embodiment, rtM204 and rtY203; in even yet another embodiment, rt235, rt236, rt237, rt238 and

20 rt239 and in even still another embodiment, rt247, rt248, rt249, rt250 and rt251; and in another embodiment,

K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;

N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;

P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;

25 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;

T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;

P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;

K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;

F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;

30 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;

V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;

- 52 -

- D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
5 Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
10 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
15 T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
20 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
25 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
30 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;

- 53 -

M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;

G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and

V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or combinations thereof or an equivalent one or more other mutation is indicative of a variant which exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof.

10

A further aspect of the present invention produces a method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside or nucleotide analog or other anti-HBV agents, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding the DNA polymerase and/or a corresponding region of the S gene, wherein the presence of a mutation selected from, in one embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment, rtH90D and rtL/F108L; in even yet another embodiment, sP120T, sM125T and sT127A; in still yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in another embodiment, sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop; in a further embodiment, sN40S, sC69STOP, sM75I, sL88P, sT118A, sW182Stop, sW196L, sY206H and sY225F; in yet another embodiment, sI81M and sP214Q; in still another embodiment, sF83S, sL173F and sW199L; in yet another embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in still another embodiment, sC69Stop/C, sC76Y, sI110V/I, sY134N, sW172Stop/W, sW196Stop, sS207R; in even still another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37); in another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63); in a further embodiment,

25  
30

- 54 -

rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91); in yet another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in still another embodiment, rtM204 and rtY203; in even yet another embodiment, rt235, rt236, rt237, rt238 and rt239 and in even still another embodiment, rt247, rt248, rt249, rt250 and rt251; and in  
 5 another embodiment,

K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 10 T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 15 V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
 V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 20 Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
 P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 25 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 30 T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;

- 55 -

- M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 5 P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 10 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 15 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or combinations thereof or an  
 equivalent one or more other mutation is indicative of a variant which exhibits a decreased  
 20 sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and  
 TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV  
 and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC  
 and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-  
 HBV agents or combination thereof.

25

The detection of HBV or its components in cells, cell lysates, cultured supernatant fluid  
 and bodily fluid may be by any convenient means including any nucleic acid-based  
 detection means, for example, by nucleic acid hybridization techniques or *via* one or more  
 polymerase chain reactions (PCRs). The term "bodily fluid" includes any fluid derived  
 30 from the blood, lymph, tissue or organ systems including serum, whole blood, biopsy and  
 biopsy fluid, organ explants and organ suspension such as liver suspensions. The invention

- 56 -

further encompasses the use of different assay formats of said nucleic acid-based detection means, including restriction fragment length polymorphism (RFLP), amplified fragment length polymorphism (AFLP), single-strand chain polymorphism (SSCP), amplification and mismatch detection (AMD), interspersed repetitive sequence polymerase chain reaction (IRS-PCR), inverse polymerase chain reaction (iPCR) and reverse transcription polymerase chain reaction (RT-PCR), amongst others. Other forms of detection include Northern blots, Southern blots, PCR sequencing, antibody procedures such as ELISA, Western blot and immunohistochemistry. A particularly useful assay includes the reagents and components required for immobilized oligonucleotide- or oligopeptide-mediated detection systems.

One particularly useful nucleic acid detection system is the reverse hybridization technique. In this technique, DNA from an HBV sample is amplified using a biotin or other ligand-labeled primer to generate a labeled amplifcon. Oligonucleotides immobilized to a solid support such as a nitrocellulose film are then used to capture amplified DNA by hybridization. Specific nucleic acid fragments are identified *via* biotin or the ligand. Generally, the labeled primer is specific for a particular nucleotide variation to be detected. Amplification occurs only if the variation to be detected is present. There are many forms of the reverse hybridization assay and all are encompassed by the present invention.

Detecting HBV replication in cell culture is particularly useful.

This and other aspects of the present invention is particularly amenable to microarray analysis such as to identify oligonucleotides including sense and antisense molecules, RNAi or siRNA molecules or DNA or RNA-binding molecules which down-regulate genomic sequences or transcripts of HBV. Microarray analysis may also be used to identify particular mutations in the HBV genome such as within the HBV DNA polymerase-coding region or the HBsAg-coding region.

- 57 -

Another aspect of the present invention contemplates a method for detecting an agent which exhibits inhibitory activity to an HBV by:

generating a genetic construct comprising a replication competent-effective  
5 amount of the genome from the HBV contained in a plasmid vector and then transfecting said cells with said construct;

contacting the cells, before, during and/or after transfection, with the agent to be tested;

10

culturing the cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agents; and

15 then subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of the agent.

20 In a preferred embodiment, the plasmid vector may include genes encoding part or all of other viral vectors such as baculovirus or adenovirus (Ren and Nassal, 2001, *supra*) and the method comprises:

generating a genetic construct comprising a replication competent-effective  
25 amount of the genome from the HBV contained in or fused to an amount of a baculovirus genome or adenovirus genome effective to infect cells and then infecting said cells with said construct;

contacting the cells, before, during and/or after infection, with the agent to be  
30 tested;

- 58 -

culturing the cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agent; and

5           then subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of the agent.

10   In an alternative embodiment, the method comprises:

generating a continuous cell line comprising an infectious copy of the genome of the HBV in a replication competent effective amount such that said infectious HBV genome is stably integrated into said continuous cell line such as but not limited to 2.2.15

15   or AD;

contacting the cells with the agent to be tested;

20           culturing the cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to the agent; and

25           then subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of the agent.

The above-mentioned methods are particularly useful in identifying or developing agents against HBV variants such as those carrying mutations, in one embodiment, rtS21A, 30   rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a



- 59 -

further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment rtH90D and rtL/F108L; in even yet

5 another embodiment, rtL157L/M, rtA181V and rtV207I; in even still another embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; in another embodiment, rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtL204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H; in a further embodiment, sP120T, sM125T and sT127A; in yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in still

10 another embodiment, sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop; in even yet another embodiment, sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182STOP, sW196L, sY206H and sY225F; in even still another embodiment, sI81M and sP214Q; in another embodiment, sF83S, sL173F and sW199L; in a further embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in yet another

15 embodiment, sC69Stop/C, sC76Y sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R; in still another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37; in even yet another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63; in even still another embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91; in another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in a

20 further embodiment, rtM204 and rtY203; in yet another embodiment, rt235, rt236, rt237, rt238 and rt239 in still another embodiment, rt247, rt248, rt249, rt250 and rt251; and in even yet another embodiment, K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion; P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;

25 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion; T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion; P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion; K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;

30 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion; V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;

- 60 -

- D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
5 Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
10 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
15 T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
20 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
25 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
30 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;

- 61 -

M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion.

- 5 Accordingly, another aspect of the present invention contemplates a method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside or nucleotide analog or other potential anti-HBV agent, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence of the envelope genes or DNA polymerase gene selected from, in one
- 10 embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M,
- 15 rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment rtH90D and rtL/F108L; in even yet another embodiment, rtL157L/M, rtA181V and rtV207I; in even still another embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; in another embodiment, rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H; in a further embodiment, sP120T, sM125T
- 20 and sT127A; in yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in still another embodiment, sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop; in even yet another embodiment, sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182STOP, sW196L, sY206H and sY225F; in even still another embodiment, sI181M and sP214Q; in another embodiment, sF83S, sL173F and sW199L; in a further
- 25 embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in yet another embodiment, sC69Stop/C, sC76Y sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R; in still another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37; in even yet another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63; in even still another embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91; in
- 30 another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in a further embodiment, rtM204 and rtY203; in yet another embodiment, rt235, rt236,

- 62 -

rt237, rt238 and rt239 in still another embodiment, rt247, rt248, rt249, rt250 and rt251; and in even yet another embodiment,

- K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 5 P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 10 F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
 V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 15 S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
 20 P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 25 Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 30 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;

- 63 -

- T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 5 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L;  
 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I;  
 10 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R;  
 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G;  
 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/QE; and  
 15 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or combinations thereof or an  
 equivalent one or more other mutation is indicative of a variant wherein said variant  
 exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV  
 and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and  
 LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and  
 20 FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or  
 nucleotide analogs or other anti-HBV agents or combination thereof.

The detection of amino acid variants of DNA polymerase is conveniently accomplished by  
 reference to the amino acid sequence shown in Formulae I and II. The polymorphisms  
 25 shown represent the variations shown in various databases for active pathogenic HBV  
 strains. Where an HBV variant comprises an amino acid different to what is represented,  
 then such an isolate is considered a putative HBV variant having an altered DNA  
 polymerase activity.

30 The present invention further contemplates agents which inhibit ADV, LMV, TFV, or  
 FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV,

- 64 -

FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV resistant HBV variants. Such agents are particularly useful if long term treatment by ADV, LMV, FTC and/or TFV and/or optionally other nucleoside analogs or nucleotide analogs such as TFV is contemplated by the clinician. The agents may be DNA or RNA or proteinaceous or non-proteinaceous chemical molecules. Natural product screening such as from plants, coral and microorganisms is also contemplated as a useful potential source of masking agents as is the screening of combinatorial or chemical libraries. The agents may be in isolated form or in the form of a pharmaceutical composition or formulation and may be administered in place of or sequentially or simultaneously with a nucleoside or nucleotide analog. Furthermore, rationale drug design is contemplated including solving the crystal or NMR structure of, for example, HBV DNA polymerase and designing agents which can bind to the enzyme's active site. This approach may also be adapted to other HBV components.

Accordingly, another aspect of the present invention contemplates a method for detecting an agent which exhibits inhibitory activity to an HBV which exhibits resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof, , said method comprising:

generating a genetic construct comprising a replication competent-effective amount of the genome from said HBV contained in a plasmid vector and then transfecting said cells with said construct;

contacting said cells, before, during and/or after transfection, with the agent to be tested;

30

- 65 -

culturing said cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agent; and

5           subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of said agent.

Still another aspect of the present invention provides a method for detecting an agent  
10   which exhibits inhibitory activity to an HBV which exhibits resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-  
15   HBV agents or combination thereof, , said method comprising:

generating a genetic construct comprising a replication competent-effective amount of the genome from said HBV contained in or fused to an amount of a baculovirus genome effective to infect cells and then infecting said cells with said construct;

20

contacting said cells, before, during and/or after infection, with the agent to be tested;

culturing said cells for a time and under conditions sufficient for the HBV to  
25   replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agent; and

subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed  
30   genetic material and/or assembled and/or been released in the presence of said agent.

- 66 -

Preferably, the HBV genome is stably integrated into the cells' genome.

Particularly useful cells are 2.2.15 cells (Price *et al.*, *Proc. Natl. Acad. Sci. USA* 86(21): 8541-8544, 1989 or AD cells (also known as HepAD32 cells or HepAD79 cells [Ying *et al.*, *Viral Hepat.* 7(2): 161-165, 2000.

Whilst the baculovirus vector is a particularly useful in the practice of the present invention, the subject invention extends to a range of other vectors such as but not limited to adenoviral vectors.

10

The present invention further extends to cell lines (e.g. 2.2.15 or AD cells) carrying genetic constructs comprising all or a portion of an HBV genome or a gene or part of a gene therefrom.

15 The present invention also provides for the use of the subject HBV variants to screen for anti-viral agents. These anti-viral agents inhibit the virus. The term "inhibit" includes antagonizing or otherwise preventing infection, replication, assembly and/or release or any intermediate step. Preferred anti-viral agents include nucleoside or nucleotide analogs or anti-HBV agents, however, the present invention extends to non-nucleoside molecules.

20

In addition, rational drug design is also contemplated to identify or generate chemical molecules which either mimic a nucleoside or which interact with a particular nucleotide sequence or a particular nucleotide. Combinatorial chemistry and two hybrid screening are some of a number of techniques which can be employed to identify potential therapeutic or

25

diagnostic agents.

In one example, the crystal structure or the NMR structure of polymerase or the surface antigen is used to rationally design small chemical molecules likely to interact with key regions of the molecule required for function and/or antigenicity. Such agents may be

30

useful as inhibitors of polymerase activity and/or may alter an epitope on the surface antigen.



- 67 -

Several models of the HBV polymerase have been prepared due to the similarity with reverse transcriptase from HIV (Das *et al.*, *J. Virol.* 75(10): 4771-4779, 2001; Bartholomeusz *et al.*, *Intervirology* 40(5-6): 337-342 1997; Allen *et al.*, *Hepatology* 27(6):  
5 1670-1677, 1998). The models of the HBV polymerase can be used for the rational drug design of new agents effective against HBV encoding the resistant mutations as well as wild type virus. The rational drug that is designed may be based on a modification of an existing antiviral agent such as the agent used in the selection of the HBV encoding the mutations associated with resistance. Viruses or clones expressing HBV genomic material  
10 encoding the mutations may also be used to screen for new antiviral agents.

In an alternative embodiment, the present invention also contemplates a method for detecting an agent which exhibits inhibitory activity to an HBV polymerase in an *in vitro* polymerase assay. The HBV polymerase activity can be examined using established assays  
15 (Gaillard *et al.*, *Antimicrob Agents Chemother.* 46(4): 1005-1013, 2002; Xiong *et al.*, *Hepatology* 28(6): 1669-1673, 1998).

As indicated above, microarray technology is also a useful means of identifying agents which are capable of interacting with defined HBV internal or external components. For  
20 example, arrays of HBV DNA polymerase or peptide fragments thereof carrying different amino acid variants may be used to screen for agents which are capable of binding or otherwise interacting with these molecules. This is a convenient way of determining the differential binding patterns of agents between HBV variants. Arrays of antibodies may also be used to screen for altered HBsAg molecules. Microarrays are also useful in  
25 proteomic analysis to identify molecules such as antibodies, interferons or cytokines which have an ability to interact with an HBV component. Microarrays of DNA and RNA molecules may also be employed to identify sense and antisense molecules for genetic regions on the HBV genome or transcripts thereof.

30 The above methods are particularly useful in identifying an inhibitor of an HBV resistant to or exhibiting reduced sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV

- 68 -

and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof. The present invention  
5 extends, therefore, to compositions of the inhibitors. The inhibitors may also be in the form of antibodies or genetic molecules such as ribozymes, antisense molecules and/or sense molecules for co-suppression or the induction of RNAi or may be other nucleoside or nucleotide analogs or other anti-HBV agents or derivatives of known analogs. Reference to RNAi includes reference to short, interfering RNAs (siRNA).

10

The term "composition" includes a "pharmaceutical composition" or a formulation.

The inhibitor is referred to below as an "active ingredient" or "active compound" and may be selected from the list of inhibitors given above.

15

The composition may include an antigenic component of the HBV, a defective HBV variant or an agent identified through natural product screening or rational drug design (including combinatorial chemistry).

20 Pharmaceutically acceptable carriers and/or diluents include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the therapeutic compositions is  
25 contemplated. Supplementary active ingredients can also be incorporated into the compositions.

The pharmaceutical composition may also comprise genetic molecules such as a vector capable of transfecting target cells where the vector carries a nucleic acid molecule capable  
30 of encoding an aspartyl protease inhibitor. The vector may, for example, be a viral vector.

Pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water soluble) and sterile powders for the extemporaneous preparation of sterile injectable solutions. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi.

5 The carrier can be a solvent or dilution medium comprising, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof and vegetable oils. The proper fluidity can be maintained, for example, by the use of surfactants. The preventions of the action of microorganisms can be brought about by various anti-bacterial and anti-fungal agents, for example,  
10 parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminium monostearate and gelatin.

15

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with the active ingredient and optionally other active ingredients as required, followed by filtered sterilization or other appropriate means of sterilization. In the case of sterile powders for the preparation of sterile injectable  
20 solutions, suitable methods of preparation include vacuum drying and the freeze-drying technique which yield a powder of active ingredient plus any additionally desired ingredient.

When the active ingredient is suitably protected, it may be orally administered, for  
25 example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsule, or it may be compressed into tablets. For oral therapeutic administration, the active ingredient may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers and the like. Such compositions and preparations should contain at least 1%  
30 by weight of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight

- 70 -

of the unit. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 0.1  $\mu$ g and 200 mg of active compound. Alternative dosage amounts  
5 include from about 1  $\mu$ g to about 1000 mg and from about 10  $\mu$ g to about 500 mg. These dosages may be per individual or per kg body weight. Administration may be per hour, day, week, month or year.

The tablets, troches, pills, capsules and the like may also contain the components as listed  
10 hereafter. A binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen or cherry flavouring. When the dosage unit form is a capsule, it may contain,  
15 in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavouring. Of course, any material used in  
20 preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound(s) may be incorporated into sustained-release preparations and formulations.

As stated above, the present invention further extends to an isolated HBsAg from the HBV  
25 variants herein described. More particularly, the present invention provides an HBsAg or a recombinant form thereof or derivative or chemical equivalent thereof. The isolated surface component and, more particularly, isolated surface antigen or its recombinant, derivative or chemical equivalents are useful in the development of biological compositions such as vaccine formulations.

- 71 -

Yet another aspect of the present invention provides a composition comprising a variant HBV resistant to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or an HBV surface antigen from said variant HBV or a recombinant or derivative form thereof or its chemical equivalent and one or more pharmaceutically acceptable carriers and/or diluents. Such a composition may be regarded as a therapeutic composition and is useful in generating an immune response including a humoral response.

10 Generally, the HBV variants are "defective" and in themselves are unable to cause a sustained infection in a subject.

As indicated above, antibodies may be generated to the mutant HBV agents and used for passive or direct vaccination against infection by these viruses. The antibodies may be generated in humans or non-human animals. In the case of the latter, the non-human antibodies may need to be deimmunized or more specifically humanized prior to use. Deimmunized may include, for example, grafting complementarity determining regions (CDRs) from the variable region of a murine or non-human animal anti-HBV antibody onto a human consensus fragment antibody binding (Fab) polypeptide. Alternatively,

20 amino acids defining epitopes in the variable region of the antibody may be mutated so that the epitopes are no longer recognized by the human MHC II complex.

Insofar as ribozyme, antisense or co-suppression (RNAi) or siRNA or complexes thereof repression is concerned, this is conveniently aimed at post-transcription gene silencing.

25 DNA or RNA may be administered or a complex comprising RNAi or a chemical analog thereof specific for HBV mRNA may be employed.

All such molecules may be incorporated into pharmaceutical compositions.

- 72 -

In another embodiment, the present invention provides a biological composition comprising a variant HBV or an HBsAg or L, M or S proteins from said variant HBV or a recombinant or derivative form thereof or its chemical equivalent.

- 5 Generally, if an HBV is used, it is first attenuated. The biological composition according to this aspect of the present invention generally further comprises one or more pharmaceutically acceptable carriers and/or diluents.

The biological composition may comprise HBsAg or like molecule from one HBV variant  
10 or the composition may be a cocktail of HbsAgs or L, M or S proteins or like molecules from a range of ADV- and/or LMV- and/or, FTC- and/or TFV-resistant HBV variants. Similar inclusions apply where the composition comprises an HBV.

The present invention is further directed to the use of defective HBV variants in the  
15 manufacture of therapeutic vaccines to vaccinate individuals against infection by HBV strains having a particular nucleotide sequence or encoding a particular polymerase or surface antigen or L, M or S proteins.

Examples of suitable vaccine candidates are defective forms of HBV variants comprising a  
20 mutation selected from, in one embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment,  
25 rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment rtH90D and rtL/F108L; in even yet another embodiment, rtL157L/M, rtA181V and rtV207I; in even still another embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; in another embodiment, rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H; in a further  
30 embodiment, sP120T, sM125T and sT127A; in yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in still another embodiment, sS126T,

- 73 -

sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop; in even yet another embodiment, sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182STOP, sW196L, sY206H and sY225F; in even still another embodiment, s181M and sP214Q; in another embodiment, sF83S, sL173F and sW199L; in a further embodiment, sI126T, sK160R, 5 sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in yet another embodiment, sC69Stop/C, sC76Y sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R; in still another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37; in even yet another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63; in even still another embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91; in another embodiment, rtP177, 10 rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in a further embodiment, rtM204 and rtY203; in yet another embodiment, rt235, rt236, rt237, rt238 and rt239 in still another embodiment, rt247, rt248, rt249, rt250 and rt251; and in even yet another embodiment,

K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 15 N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 20 K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
 25 V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
 S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 30 I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
 P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;

- 74 -

- F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 5 Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 10 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 15 H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 20 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 25 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E; and  
 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or a combination of two or  
 more mutations.

In one embodiment, for example, an HBV variant may be identified having a particular  
 30 mutation in its polymerase conferring resistance or decreased sensitivity to a nucleoside  
 analog. This variant may then be mutated to render it defective, i.e. attenuated or unable to



- 75 -

cause infection. Such a defective, nucleoside analog-resistant virus may then be used as a therapeutic vaccine against virulent viruses having the same mutation in its polymerase.

The subject invention extends to kits for assays for variant HBV resistant to ADV, LMV,  
5 TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC  
and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV  
and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV.. Such  
kits may, for example, contain the reagents from PCR or other nucleic acid hybridization  
technology or reagents for immunologically based detection techniques. A particularly  
10 useful assay includes the reagents and components required for immobilized  
oligonucleotide- or oligopeptide-mediated detection systems.

Still another aspect of the present invention contemplates a method for determining the  
potential for an HBV to exhibit reduced sensitivity to ADV, LMV, TFV, or FTC, or ADV  
15 and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV,  
or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV  
and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other  
nucleoside or nucleotide analogs or other anti-HBV agents or combination thereof, said  
method comprising isolating DNA or corresponding mRNA from said HBV and screening  
20 for a mutation in the nucleotide sequence encoding HBV DNA polymerase resulting in at  
least one amino acid substitution, deletion and/or addition in any one or more of domains F  
and G, and domains A through to E or a region proximal thereto of said DNA polymerase  
and associated with resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or  
ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and  
25 LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV,  
ADV and LMV and FTC, or ADV and FTC and LMV and TFV, wherein the presence of  
such a mutation is an indication of the likelihood of resistance to said ADV, LMV, TFV, or  
FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV,  
FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC  
30 and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV.

- 76 -

An assessment of a potential viral variant is important for selection of an appropriate therapeutic protocol. Such an assessment is suitably facilitated with the assistance of a computer programmed with software, which *inter alia* adds index values ( $I_{vs}$ ) for at least two features associated with the viral variants to provide a potency value ( $P_A$ )  
5 corresponding to the resistance or sensitivity of a viral variant to a particular chemical compound or immunological agent. The  $I_{vs}$  can be selected from (a) the ability to exhibit resistance for reduced sensitivity to a particular compound or immunological agent; (b) an altered DNA polymerase from wild-type HBV; (c) an altered surface antigen from wild-type HBV; or (d) morbidity or recovery potential of a patient. Thus, in accordance with the  
10 present invention,  $I_{vs}$  for such features are stored in a machine-readable storage medium, which is capable of processing the data to provide a  $P_A$  for a particular viral variant or a biological specimen comprising same.

Thus, in another aspect, the invention contemplates a computer program product for  
15 assessing the likely usefulness of a viral variant or biological sample comprising same for determining an appropriate therapeutic protocol in a subject, said product comprising:

- (1) code that receives as input  $I_{vs}$  for at least two features associated with said viral agents or biological sample comprising same, wherein said features are selected  
20 from:
  - (a) the ability to exhibit resistance for reduced sensitivity to a particular compound or immunological agent;
  - (b) an altered DNA polymerase from wild-type HBV;
  - (c) an altered surface antigen from wild-type HBV;
  - (d) morbidity or recovery potential of a patient; or
  - (e) altered replication capacity (increased or decreased);
- (2) code that adds said  $I_{vs}$  to provide a sum corresponding to a  $P_v$  for said viral  
25 variants or biological samples; and  
30

- 77 -

- (3) a computer readable medium that stores the codes.

In a related aspect, the invention extends to a computer for assessing the likely usefulness of a viral variant or biological sample comprising same in a subject, wherein said computer  
5 comprises:

- (1) a machine-readable data storage medium comprising a data storage material  
10 encoded with machine-readable data, wherein said machine-readable data comprise  
I<sub>v</sub>s for at least two features associated with said viral variant or biological sample;  
wherein said features are selected from:-
- (a) the ability to exhibit resistance for reduced sensitivity to a particular  
compound or immunological agent;
  - (b) an altered DNA polymerase from wild-type HBV;
  - 15 (c) an altered surface antigen from wild-type HBV;
  - (d) morbidity or recovery potential of a patient; or
  - (e) altered replication capacity (increased or decreased);
- (2) a working memory for storing instructions for processing said machine-readable  
20 data;
- (3) a central-processing unit coupled to said working memory and to said machine-  
readable data storage medium, for processing said machine readable data to provide  
a sum of said I<sub>v</sub>s corresponding to a P<sub>v</sub> for said compound(s); and
- 25 (4) an output hardware coupled to said central processing unit, for receiving said P<sub>v</sub>.

Any general or special purpose computer system is contemplated by the present invention  
and includes a processor in electrical communication with both a memory and at least one  
30 input/output device, such as a terminal. Figure 19 shows a generally suitable computer  
system. Such a system may include, but is not limited, to personal computers, workstations

- 78 -

or mainframes. The processor may be a general purpose processor or microprocessor or a specialized processor executing programs located in RAM memory. The programs may be placed in RAM from a storage device, such as a disk or pre-programmed ROM memory. The RAM memory in one embodiment is used both for data storage and program  
5 execution. The computer system also embraces systems where the processor and memory reside in different physical entities but which are in electrical communication by means of a network.

In an alternative embodiment, the program screens for a mutation selected from, in one  
10 embodiment, rtS21A, rtL122F, rtN124H, rtH126R, rtT28N, rtP130Q, rtD131N and rtY135C; in another embodiment, rtN/S/T/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M; in a further embodiment, rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D; in yet another embodiment, rtS78T, rtV84M,  
15 rtY126C, rtV191I, rtM204I and rtV214A; in still another embodiment rtH90D and rtL/F108L; in even yet another embodiment, rtL157L/M, rtA181V and rtV207I; in even still another embodiment, rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K; in another embodiment, rtS78S/T, rtN118N/S; rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H; in a further embodiment, sP120T, sM125T  
20 and sT127A; in yet another embodiment, sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C; in still another embodiment, sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop; in even yet another embodiment, sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182STOP, sW196L, sY206H and sY225F; in even still another embodiment, s181M and sP214Q; in another embodiment, sF83S, sL173F and sW199L; in a further  
25 embodiment, sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C; in yet another embodiment, sC69Stop/C, sC76Y sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R; in still another embodiment, rtK32, rtN33, rtP34, rtH35 and rtT37; in even yet another embodiment, rtP59, rtK60, rtF61, rtA62 and rtV63; in even still another embodiment, rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtL/L91; in  
30 another embodiment, rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184; in a further embodiment, rtM204 and rtY203; in yet another embodiment, rt235, rt236,

- 79 -

rt237, rt238 and rt239 in still another embodiment, rt247, rt248, rt249, rt250 and rt251; and  
in even yet another embodiment,

- K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
5 P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
10 F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
15 S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
20 P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
25 Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
30 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;

- 80 -

- T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
5 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
10 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E; and  
15 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or a combination of two or  
more mutations.

The present invention is further described by the following non-limiting Examples.

- 81 -

## EXAMPLE 1

### *Overlapping genome of HBV*

The overlapping genome of HBV is represented in Figure 1. The gene encoding DNA  
5 polymerase (P), overlaps the viral envelope genes, Pre-S1 and Pre-S2, and partially  
overlaps the X and core (C) genes. The HBV envelope comprises small, middle and large  
proteins HBV surface antigens. The large protein component is referred to as the HBV  
surface antigen (HBsAg) and is encoded by the S gene sequence. The Pre-S1 and Pre-S2  
gene sequences encode the other envelope components.

10

## EXAMPLE 2

### *Patients and Treatment*

Patient A, a 48 year old Lebanese woman was initially referred for evaluation of  
15 thrombocytopenia and hepatosplenomegaly. At this time the patient had abnormal LFT's  
(ALT 67 U/L, normal <55) and the HBV DNA was 61 pg/ml (231 days prior to the start of  
treatment). The patient was HBsAg and HBeAg positive. The ALT's fluctuated between  
50-70 IU/L from (-231 to -35 days pretreatment). ADV was commenced on Day 0 in a  
clinical trial on 30 mg/day. HBV DNA levels were reduced with ADV treatment. The  
20 ADV treatment was reduced to 10 mg /day (144 days post-treatment). There was a  
problem with the randomization treatment protocol. The patient was on antiviral treatment  
for 1 month only during the second year of the treatment period. The study was completed  
on Day 679 post ADV treatment. The patient was not on ADV treatment until the open  
label ADV was recommenced on Day 875 from the start of the initial ADV treatment. This  
25 second period of ADV treatment was given for 108 days (day 983 post initial ADV  
treatment). The HBV DNA levels remained at 7-10 pg/ml ( $1.96 \times 10^5$  to  $2.8 \times 10^5$   
copies/ml). At Day 983, ADV treatment was stopped and the patient was treated with  
LMV.

30 Patient B is a male liver transplant patient. The patient has been on both sequential and  
combination antiviral therapy including HBIG, FCV+HBIG, LMV+HBIG, LMV,

- 82 -

LMV+GCV, LMV+FCV+GCV, LMV+GCV and finally LMV+ADV. The patient has been on long term ADV+LMV treatment for over 795 days.

Patient C, is a 58 year old male. Prior to ADV treatment the patient had abnormal LFT's  
5 (ALT 240 IU/L, normal <55) and the HBV DNA was  $2 \times 10^7$  copies/ml. ADV was commenced on Day 0 in a clinical trial on 10 mg/day for two years. The average ALT during the two year clinical trial period was 114 IU/L. However, the ALT was rising and at 630 days after the start of ADV treatment the ALT remained high 407 IU/L. Open label ADV was commenced on Day 668 from the start of the initial ADV treatment. This second  
10 period of ADV treatment was given for 71 days. The HBV DNA levels remained high during open label ADV treatment ( $3.7 \times 10^6$  to  $1.5 \times 10^7$  copies/ml). The peak ALT during open label ADV treatment was 517 IU/L (Day 738). The next day (Day 739), ADV treatment was stopped and the patient was treated with LMV.

15

### EXAMPLE 3

#### *Detection of Viral Markers*

Hepatitis B surface antigen (HBsAg), hepatitis B e antigen (HBeAg), anti-HBe and hepatitis B core antigen (HBcAg) specific IgG and IgM were measured using  
20 commercially available immunoassays (Abbott Laboratories, North Chicago, IL, USA). Hepatitis B viral DNA levels were measured using a capture hybridization assay according to the manufacturer's directions (Digene Hybrid Capture II, Digene Diagnostics Inc., Beltsville, MD). The manufacturers stated cut-off for detecting HBV viremia in clinical specimens was  $0.7 \times 10^6$  copies/ml or 2.5 pg/ml, [Hendricks *et al.*, *Am J Clin Pathol* 104:  
25 537-46, 1995]. HBV DNA levels can also be quantitated using other commercial kits such as Cobas amplification HBV monitor kit (Roche).



- 83 -

**EXAMPLE 4*****Sequencing of HBV DNA***

HBV DNA was extracted from 100 µl of serum as described previously by Aye *et al.*, *J. Hepatol.* 26: 1148-1153, 1997. Oligonucleotides were synthesized by Geneworks, Adelaide, Australia. Amplification of the HBV polymerase gene has been described by Aye *et al.*, 1997, *supra*.

The specific amplified products were purified using PCR purification columns from MO BIO Laboratories Inc (La Jolla, CA) and directly sequenced using Big Dye terminator Cycle sequencing Ready Reaction Kit (Perkin Elmer, Cetus Norwalk, CT). The PCR primers were used as sequencing primers, OS1 5'-GCC TCA TTT TGT GGG TCA CCA TA-3' (nt 1408-1430) [SEQ ID NO:3], TTA3 5'-AAA TTC GCA GTC CCC AAA-3'(nt2128-2145) [SEQ ID NO:4], JM 5'-TTG GGG TGG AGC CCT CAG GCT - 3'(nt1676-1696) [SEQ ID NO:5], TTA4 5'-GAA AAT TGG TAA CAG CGG -3' (nt 2615-2632) [SEQ ID NO:6], OS2 5' TCT CTG ACA TAC TTT CCA AT 3' (nt 2798-2817) [SEQ ID NO:7], to sequence the internal regions of the PCR products.

**EXAMPLE 5*****Analysis of HBV DNA***

Patient A: During ADV treatment, unique HBV mutations were detected by sequencing (Tables 4 and 5) This includes the unique mutation at rtY135C in addition to the mutation at rtT128N that was present prior to ADV treatment. A number of other unique changes were also detected in the polymerase and in the overlapping envelope gene (Table 5, Figures 4, 5 and 6). The unique change in the HBsAg include sP120T. These unique changes were compared to reference sequences from each of the seven genotypes A-G as well as a consensus sequence from pretreatment samples to determine unique changes.

30

- 84 -

Patient B: The HBV mutations prior to ADV treatment and during ADV treatment are listed in Table 6 and 7 and Figures 7, 8, and 9. The unique changes in the rt region of the HBV DNA polymerase include rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M. The unique changes in the HBsAg include sT118R, sM133T, sF134V,  
5 sI195M, sS207R, sY225Y/C.

Patient C: The HBV mutations prior to ADV treatment and during ADV treatment are listed in Tables 8 and 9 and Figures 10, 11 and 12. The unique changes in the rt region of the HBV DNA polymerase include rtN53D, rtS116P, rtF151F/T, rtN236T and rtN238D.  
10 The unique changes in the HBsAg include sG145A and sW172stop.

Patient D: The HBV mutations during ADV treatment is listed in Table 10 and Figures 13, 14 and 15. The unique changes in the HBV DNA polymerase include rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A. The unique changes in the surface include  
15 sN40S and sC69 Stop. A number of unique changes were detected after the stop codon mutation at codon 69 of the S gene including sM75L, sL88P, sT118A, sW182stop, sW196L, sY206H and sY225F.

Patient E: The HBV mutations during ADV treatment is listed in Table 11 and Figures  
20 16, 17 and 18. The unique changes in the HBV DNA polymerase include rtH90D and rtL/F108L. The unique changes in the surface include sI81M and sP214Q. A six nucleotide insertion was also detected resulting in a two amino acid insertion in the HBV polymerase and envelope gene at codons rt131 and s122, respectively. This insertion was previously detected in pre-ADV samples.

25

## EXAMPLE 6

### *Adefovir Dipivoxil (ADV)*

ADV (formerly Bis-pom PMEA)) is a potent inhibitor of HBV replication. The structure of  
30 ADV is shown in Figure 2 and its synthesis is described by Benzaria *et al.*, *J Med Chem.* 39: 4958-4965, 1996).

- 85 -

**EXAMPLE 7*****HBV rt mutants***

The HBV polymerase has similarities to other polymerases including HIV. Thus, mutations associated with resistance to antiviral agents may occur within the polymerase in functionally important regions such as the nucleotide triphosphate binding pocket that may also include the interaction between the DNA primer and template strand, magnesium ions and nucleoside triphosphates or nucleoside/ nucleotide analogs (and there vaious phosphroylated forms). Codons which are proposed to be mutated during anti-viral selection pressure are rtK32, rt N33, rtP34, rtH35 and rtT37 (that are upstream from the F domain); rt P59, rtK60, rtF61, rtA62 and rtV63 (between the F and A domains), rtD83, rtV84, rtS85, rtA86, rt Y89, rt H90 and rtI/L91 (within the A domain and the region immediately prior to and after), rtP177, rtF178, rt L179, rtL180, rtA181, rtQ182, rtF183 and rtT184 (B domain); rtM204 and rtY203(C Domain), rtL235, rtN236, rtP/T237, rtN/H/A/S/Q238 and rtK239 (D Domain), rtL247, rtN/H248, rtF249, rtM250 and rtG251 (E Domain). The codons are defined in Table 12 and examples of various mutants are given in Tables 13 and 14.

**EXAMPLE 8*****Patient F***

The HBV mutations during ADV treatment of Patient F are listed in Table 15 and Figures 20, 21 and 22. The unique changes in the HBV DNA polymerase includes rtL157L/M, rtA181V, rtV207L, and rtN236T. The unique changes in the surface includes sF83S, sL173F and sW199L.

- 86 -

**EXAMPLE 9*****Patient G***

The HBV mutations during ADV treatment of Patient G are listed in Table 16 and Figures  
5 23, 24 and 25. The unique changes in the HBV DNA polymerase includes rtL80V,  
rtP109S, rtI163V, rtM204I, rtL229M and rtN/H/A/S/Q238K. The unique changes in the  
surface includes sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and  
sY221C.

10

**EXAMPLE 10*****Patient H***

The HBV mutations during ADV treatment in Patient H are listed in Table 17 and Figures  
26, 27 and 28. The unique changes in the HBV DNA polymerase includes rtS78S/T,  
15 rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E,  
and rtN238N/H. The unique changes in the surface include sC69Stop/C, sC76Y sI110V/I,  
sY134N, sW172Stop/W, sW196Stop and sS207R.

**EXAMPLE 11**

20

***In vitro analysis of ADV resistance***

The sensitivity/resistance profile of HBV mutants to ADV was examined *in vitro* using  
recombinant HBV/baculovirus. The procedure for analyzing the resistance profile is  
outlined in the following Examples 12-20.

25

**EXAMPLE 12*****Cell culture***

Sf21 insect cells were maintained in supplemented Grace's insect medium further  
30 supplemented with 10% v/v heat-inactivated fetal bovine serum (Gibco BRL,

- 87 -

Gaithersburg, MD) in humidified incubator at 28°C with CO<sub>2</sub>. HepG2 cells were maintained in minimal essential medium supplemented with 10% v/v heat-inactivated fetal bovine serum (MEM-FBS). HepG2 cells were grown in humidified 37°C incubators at 5% v/v CO<sub>2</sub>.

5

### EXAMPLE 13

#### *Preparation of HBV/baculovirus transfer vector with specific point mutations*

The recombinant HBV/baculovirus system used for antiviral testing has been previously described (Delaney *et al.*, *Antimicrob Agents Chemother* 45(6): 1705-1013, 2001). In brief, the recombinant transfer vector was created by excising a fragment containing the 1.3x HBV genome construct and cloning it into the multiple cloning region of a baculovirus vector pBlueBac4.5 (Invitrogen, Carlsbad, CA). Point mutations were created by site directed mutagenesis using the commercial kits according to the manufacturer's specifications (QuikChange, Stratagene). HBV/ baculovirus recombinant clones encoding the reverse transcriptase mutations rtA181T/N236T/N238D and rtN236T/N236D in combination with the precore mutation at G1896A (pcW28 stop) or wild-type with respect to codon pcW28, were prepared by site-directed mutagenesis. The nucleotide sequence of the plasmid and the point mutations generated by site directed mutagenesis were confirmed by sequencing using the ABI Prism Big Dye Terminator Cycle Sequencing Ready Reaction Kit according to the manufacturer's specifications (Perkin Elmer, Cetus Norwalk, CT).

20

### EXAMPLE 14

#### *Generation of recombinant baculoviruses containing the 1.3 HBV construct*

25

Purified recombinant transfer vector and linear AcMNPV baculovirus DNA were co-transfected into Sf21 cells using the BacNBlue transfection kit from Invitrogen (Carlsbad, CA); recombinant viruses were isolated by plaque assay according to the manufacturer's instructions. A series of recombinant viruses were amplified from isolated plaques by infecting 100-mm dishes of Sf21 cells. Viral DNA was extracted from amplified viruses

30

- 88 -

using standard procedures. Purified viral DNA was digested with restriction enzymes and then fractionated by electrophoresis in a 1% v/v agarose gel. Southern blotting was performed to determine which virus isolates contained the intact 1.3 HBV construct. A Boehringer Mannheim Random Prime DNA Labeling kit (Indianapolis, IN) was used to generate [ $P^{32}$ ]-radiolabeled probes. A full-length double-stranded HBV genome was used as a template for all radiolabeled probes. Viral DNA sequence was confirmed by PCR amplification of the polymerase catalytic region using the sense primer 5'-GCC TCA TTT TGT GGG TCA CCA TA-3' [SEQ ID NO:8], (nucleotide 1408 to 1430 according to HBV Genebank Accession number M38454) and the antisense primer 5'-TCT CTG ACA TAC TTT CCA AT-3' [SEQ ID NO:9] (nucleotides 2817 to 2798 according to HBV Genebank Accession number M38454). The following primers were utilized for the sequencing of internal regions 5'-TGC ACG ATT CCT GCT CAA-3' [SEQ ID NO:10] (nucleotides 2345-2362 according to HBV Genebank Accession number M38454) and 5'-TTT CTC AAA GGT GGA GAC AG-3' [SEQ ID NO:11] (nucleotides 1790-1810 according to HBV Genebank Accession number M38454).

## EXAMPLE 15

### *Preparative baculovirus amplification and purification*

Baculoviruses were amplified by infecting suspension cultures of Sf21 cells in log phase at a multiplicity of infection (moi) of 0.5 pfu/cell. Infections were allowed to proceed until a majority of the cells in the flasks showed visible signs of infection (four to five days). Virions were concentrated from infected Sf21 medium by centrifugation at 80,000 x g and purified through a 20-60% w/v sucrose gradient. Purified virus was titrated in quadruplicate in Sf21 cells by end-point dilution. An aliquot of each high titer stock was used for DNA extraction. The polymerase gene was amplified and sequenced to confirm the presence of the site-directed mutagenesis as in Example 14.

- 89 -

## EXAMPLE 16

### *Infection of HepG2 cells with recombinant HBV expressing baculovirus*

HepG2 cells were seeded at approximately 20-40% confluency and then were grown for  
5 16-24 hours before infection. On the day of infection, triplicate plates of cells were  
trypsinized, and viable cell number was determined with a hemocytometer using Trypan  
blue exclusion. Average cell counts were calculated and used to determine the volume of  
high-titer viral stock necessary to infect cells at the indicated moi. HepG2 cells were  
washed one time with serum-free MEM to remove traces of serum. Baculovirus was  
10 diluted into MEM without serum to achieve the appropriate moi using volumes of 1.0, 0.5,  
and 0.25 ml to infect 100-mm, 60 mm, and 35-mm dishes, respectively. Baculovirus was  
adsorbed to HepG2 cells for one hour at 37°C with gentle rocking every 15 minutes to  
ensure that the inoculum was evenly distributed. The inoculum was then aspirated and  
HepG2 cells were washed two times with phosphate-buffered saline and refed MEM-FBS  
15 with or without various concentrations of agents.

## EXAMPLE 17

### *Detection of intracellular replicative intermediates*

20 HBV core particles were isolated from the cytoplasmic fraction of HepG2 cells lysed in  
0.5% w/v NP-40. Cytoplasmic extracts were adjusted to 10 mmol/l McC12 and  
unprotected DNA was removed by an incubation to 500 g/ml Proteinase K for 1.5 hours at  
37°C. HBV DNA in the samples were then extracted using commercial DNA extraction  
kits such as Qiagen (DNA extraction) or in-house methods using sequential phenol and  
25 chloroform extractions, and the nucleic acids were recovered by ethanol precipitation.  
Nucleic acids were resuspended in 50 µl /l TE (10 mmol/l Tris, 1 mmol/l  
ethylenediaminetetraacetic acid), normalized by OD260, and digested with 100 g/ml  
RNase (Boehringer Mannheim, Indianapolis, IN) for one hour at 37°C before analysis by  
real-time PCR or electrophoresis and Southern blotting. After southern blot analysis a  
30 BioRad GS-670 imaging densitometer and the Molecular Analyst software (BioRad,  
Hercules California) was used to analyze suitable exposures of Southern blots.

- 90 -

Densitometry data was fitted to logistic dose response curves using the TableCurve 2D software package from Jandel Scientific. Logistic dose response equations were used to calculate IC<sub>50</sub> and IC<sub>90</sub> values and co-efficients of variation.

5

## EXAMPLE 18

### *Real-time PCR*

For the real-time PCR based assay for HBV, HBV DNA was extracted from 200 µl of serum using the QIAamp DNA Mini Kit according to the manufacturer's instructions  
10 (QIAGEN GmbH, Hildens, Germany). Primers and a molecular beacon were designed for conserved nucleic acid sequences within the precore domain of the HBV genome to amplify and detect a 216-nucleotide product. Amplification was performed in a 50-µl reaction mixture containing 1.0 Taqman buffer A (Applied Biosystems, Foster City, CA), 3.0 mM MgCl<sub>2</sub>, 0.4 pmol of each primer per µL, forward primer, PC1 (5'-  
15 GGGAGGAGATTAGGTAA-3' [SEQ ID NO:12]) and reverse primer, PC2 (5'-GGCAAAAACGAGAGTAACTC-3' [SEQ ID NO:13]), 0.4 pmol of the HBV-specific molecular beacon per µL, (5'-FAM-CGCGTCCTACTGTTCAAGCCTCCAAGCTGT  
GACGCG-DABCYL-3' [SEQ ID NO:14]; where FAM represents fluorophore 6-carboxyfluorescein and DABCYL, 4-dimethylaminophenylazobenzoic acid, a quenching  
20 chromophore) and 1.25U of AmpliTaq Gold DNA polymerase (Perkin-Elmer). PCR was performed using the ABI PRISM 7700 spectrofluorometric thermocycler (Applied Biosystems). The PCR program consisted of an initial cycle (95°C for 10 minutes) followed by 45 amplification cycles (94°C for 15 secs, 50°C for 30 secs, 72°C for 30 secs). The instrument detected and recorded the fluorescence spectrum of each reaction tube  
25 during the annealing phase.

An external standard was constructed by ligation of a 1.3 kB wild-type HBV plasmid (genotype D) into the pBlueBac plasmid vector (Hershey Medical Center, Hershey, PA). Quantification of the DNA concentration of the plasmid was determined by  
30 spectrophotometry. Duplicates of serial 10-fold dilutions of the plasmid ranging from 10<sup>8</sup> copies/ml to 100 copies/ml were included in each run in order to generate a standard curve.



- 91 -

The copy number in each experimental reaction was determined by interpolation of the derived threshold cycle ( $C_T$ ).

## EXAMPLE 19

5

### *ADV treatments*

ADV was resuspended in sterile water, aliquoted, and frozen at -20°C to avoid repeated freezing and thawing of the drug. Medium containing ADV was prepared daily as needed using fresh aliquots of 3TC. In experiments in which ADV treatment was initiated after viral infection, HepG2 cells were exposed to the indicated concentration of ADV immediately after infection with HBV baculovirus. In experiments utilizing pretreatment with ADV, cells were fed medium containing ADV 16 hours prior to HBV baculovirus infection, HBV baculovirus infection was also carried out in medium containing ADV, and cells were refed fresh medium containing ADV immediately after completion of the infection and washing procedures.

## EXAMPLE 20

### *Antiviral testing performed with wild-type and HBV/baculovirus encoding rtA181T/N236T/N238D and rtN236T/N236D*

20

The *in vitro* antiviral drug cross-resistance testing of the HBV mutants is shown in Table 18. The laboratory reference strain of HBV (genotype D subtype ayw) containing the introduced D domain mutations demonstrated increased  $IC_{50}$  values against ADV (Table 18). The rt N236T/N238D mutation was associated with a twenty-three fold increase in  $IC_{50}$  against ADV. This was reduced to a five-fold increase when the rtA181T was also present and this triple HBV polymerase mutant was resistant to LMV.

**TABLE 4** Clinical, virological and HBV sequencing data summary for Patient A while on open label ADV.

5

Days post-ADV treatment	HBV DNA copies/ml (pg/ml)	ALT IU/L	Treatment protocol	Key polymerase mutations detected by sequencing <sup>1</sup>
-230	1.7 10 <sup>6</sup> (61)	67 U/L	pre-therapy	rtT/N128T/N rtQ/H/R215Q/stop
875			ADV recommenced	
904	1.55 x 10 <sup>6</sup>			
932	2.97 x 10 <sup>6</sup>			
959	1.76 x 10 <sup>6</sup>			
983	1.64 x 10 <sup>6</sup>	65	end ADV	rtT128N rtY135C

<sup>1</sup> Nomenclature according to Stuyver *et al.*, 2001, *supra*

**TABLE 5** Summary of HBV mutations in patient A treated with ADV

Sample name	Days post-ADV treatment	Genotype	Polymerase*	Surface
ILA1	-230	D	rtA/S21A/S rtT/N128T/N** rtQ/H/R215Q/stop	sP120P/T sI208I/L
ILA2	904	D	rtA/S21S rtF122L rtR126H rtT/N128T/N rtQ130P rtN131D rtQstop/215Q rtH248N	sP/T120P sT125M sI/208I/L
ILA3	932	D	rtA/S21S rtF122L rtR126H rtT/N128T/N rtQ130P rtN131D rtQstop/215Q rtH248N	sP/T120P sT125M sI/208I/L
ILA4	983	D	rtS21A rtL122F rtN124H rtH126R rtT128N rtP130Q rtD131N rtY135C	sP120T sM125T sT127A

\* Nomenclature according to Stuyver *et al.*, 2001, *supra*.

5 \*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.

**TABLE 6** Clinical, virological and HBV sequencing data summary for Patient B while on open label ADV.

Days post-ADV treatment	HBV DN copies/ml (pg/ml)	ALT IU/L	Treatment protocol	Key polymerase mutations detected by sequencing <sup>1</sup>
-867(S0)	183	298	pre-therapy	rtN/S/T/I/V53D rtV153G rtQ/E215S rtN248H
-8(S6)	955	427	pre-ADV on LMV	rtI/L80L rtY126Q rtL180M rtS202G rtI204V
76(S8)	not detected	150	on ADV (20 mg) and LMV	rtN/S/T/I/V53D rtY126Q rtL180M rtS202G rtI204V
637(S12)	not detected	36	on ADV (5 mg) and LMV	rtN/S/T/I/V53D rtY126Q rtL180M rtS202G rtI204V
872(S15)	not detected	67	on ADV (5 mg) and LMV	rtN/S/T/I/V53D rtY126Q rtL180M rtS202G rtI204V rtI235I/M

5 <sup>1</sup> Nomenclature according to Stuyver *et al.*, 2001, *supra*

TABLE 7 Summary of HBV mutations in Patient B treated with ADV

Sample name	Days post-ADV treatment	Genotype	Polymerase*	Surface
S0	-867	D	rtN/S/T/I/V53D rtV153G rtQ/E215S rtN248H	sM/K/L133T sF134V sS207R sL21V/L
S6	-8	D	rtI/L80L rtY126Q rtL180M rtS202G rtI204V	sT118R sM133T sF134V sI195M sS207R
S8	76	D	rtN/S/T/I/V53D rtY126Q rtL180M rtS202G rtI204V	sT118R sM133T sF134V sI195M sS207R
S12	637	D	rtN/S/T/I/V53D rtY126Q rtL180M rtS202G I204V	sT118R sM133T sF134V sI195M sS207R
S15	872	D	rtN/S/T/I/V53D rtY126Q rtL180M rtS202G rtI204V rtI235I/M	sT118R sM133T sF134V sI195M sS207R sY225Y/C

5 \* Nomenclature according to Stuyver *et al.*, 2001, *supra*

\*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.

- 96 -

**TABLE 8** Clinical, virological and HBV sequencing data summary for Patient C while on open label ADV.

Days post-ADV treatment	HBV DNA copies/ml (pg/ml)	ALT IU/L	Treatment protocol	Key polymerase mutations detected by sequencing <sup>1</sup>
-26	$2 \times 10^7$		pre-therapy	rtN53D rtS116P rtD/N/S134V rtN238D
0		240	ADV commenced clinical trial	
29		160		
630		407		
668			Open label ADV	
701	$1.5 \times 10^7$	226		
730	$3.7 \times 10^6$	361		rtN53D rtS116P rtF151S/T rtA181T rtN236T rtN238D
738		517		
739			end ADV, start LMV	

5 <sup>1</sup> Nomenclature according to Stuyver *et al.*, 2001, *supra*

TABLE 9 Summary of HBV mutations in Patient C treated with ADV

Sample name	Days post-ADV treatment	Genotype	Polymerase*	Surface
DRJ1299	-26	D	<b>rtN53D**</b> rtY54H rtS57P rtL91I <b>rtS116P</b> rtF122L rtY124H <b>rtD/N/S134V</b> rtK212R rtL217R rtS219A <b>rtN238D</b>	T126S <b>S204G</b> L209V <b>S210R</b>
DRJ1	730	D	<b>rtN53D</b> rtY54H rtS57P rtL91I <b>rtS116P</b> rtF122L rtY124H rtV134D rtY141Y/F rtL145M <b>rtF151T/F</b> <b>rtA181T</b> rtK212R rtL217R rtS219A <b>rtN236T</b> <b>rtN238D</b>	sS126T sM133L/M sS143S/T sD144A sG145A <b>sW172Stop</b>

\* Nomenclature according to Stuyver *et al.*, 2001, *supra*.

5 \*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.

TABLE 10 Summary of HBV mutations in Patient D treated with ADV

Sample Name	Genotype	Polymerase*	Surface
02575908	D	rtS78T rtV84M rtY126C rtV191I rtM204I rtV214A	sN40S sC69stop sM75I sL88P sT118A sW182STOP sW196L sY206H sY225F

\* Nomenclature according to Stuyver *et al.*, 2001, *supra*.

- 5 \*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.

TABLE 11 Summary of HBV mutations in Patient E treated with ADV

Sample Name	Genotype	Polymerase*	Surface
8123/02	A	rtH90D rtL/F108L 6nt insertion/duplication after codon rt131(aaQ&N)	sI81M sY/S100Y 6nt insertion/ duplication after codon s122 (aaT & K) sP214Q

10 \*

Nomenclature according to Stuyver *et al.*, 2001, *supra*.

- \*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.



**TABLE 12** Codons where mutations occur following exposure to nucleoside or nucleotide analogs

Region/ Domain	Original amino acid in reverse transcriptase (rt) and codon position	Nucleotide				
prior to F	K32	AAG	AAA			
	N33	AAT				
	P34	CCT				
	H35	CAC				
	T37	ACC				
F TO A	P59	CCA				
	K60	AAA				
	F61	TTC				
	A62	GCA				
	V63	GTC				
A	D83	GAT				
	V84	GTG				
	S85	TCT				
	A86	GCG				
	Y89	TAT				
	H90	CAT				
	I/L91	ATT	CTT			
B	P177	CCG				
	F178	TTT				
	L179	CTC				
	L180	CTG				
	A181	TTG				
	Q182	CAG				
	F183	TTT				
	T184	ACT				
C	Y203	TAT				
	M204	ATG				
D	L235	TTG	TTA			
	N236	AAC	AAT			
	T237	ACT	ACC			
	P237	CCT	CCC			
	N238	AAT	AAC			
	H238	CAC				
	A238	GCT				
	S238	TCT				
	Q238	CAG				

- 100 -

Region/ Domain	Original amino acid in reverse transcriptase (rt) and codon position	Nucleotide				
	K239	AAA	AAG			
E	L247	CTT	TTA	CTA	CTC	CTG
	N248	AAC	AAT			
	H248	CAT	CAC			
	F249	TTC	TTT			
	M250	ATG				
	G251	GGT	GGA	GGC	GGG	
	V251	GTC				

**TABLE 13** Target amino acid sites in rt with codons and mutations leading to amino acid changes.



Title	Codon	Amino Acid	Codon	Amino Acid	Codon	Amino Acid
K32	AAG	Lys	AAG	Lys	GAG	Glu
N33	AAT	Asn	AAT	Asn	GAT	Asp
P34	CCT	Pro	ACT	Thr	GCT	Ala
H35	CAC	His	AAC	Asn	GAC	Asp
T37	ACC	Thr	ACC	Thr	GCC	Ala
P59	CCA	Pro	ACA	Thr	GCA	Ala
K60	AAA	Lys	AAA	Lys	GAA	Glu
F61	TTC	Phe	ATC	Ile	GTC	Val
A62	GCA	Ala	ACA	Thr	GCA	Ala
V63	GTC	Val	ATC	Ile	GTC	Val
D83	GAT	Asp	AAT	Asn	GAT	Asp
V84	GTG	Val	ATG	Met	GTG	Val
S85	TCT	Ser	ACT	Thr	GCT	Ala
A86	GCG	Ala	ACG	Thr	GCG	Ala
Y89	TAT	Tyr	AAT	Asn	GAT	Asp
H90	CAT	His	AAT	Asn	GAT	Asp
I/L91	ATT	Ile	ATT	Ile	GTT	Val
P177	CCG	Pro	ACG	Thr	GCG	Ala
F178	TTT	Phe	ATT	Ile	GTT	Val
L179	CTC	Leu	ATC	Ile	GTC	Val
L180	CTG	Leu	ATG	Met	GTG	Val
A181	TTG	Leu	ATG	Met	GTG	Val
Q183	CAG	Gln	AAG	Lys	GAG	Glu
F183	TTT	Phe	ATT	Ile	GTT	Val
T184	ACT	Thr	ACT	Thr	GCT	Ala
Y203	TAT	Tyr	AAT	Asn	GAT	Asp
M204	ATG	Met	ATG	Met	GTG	Val
L235	TTG	Leu	ATG	Met	GTG	Val
N236	AAC	Asn	AAC	Asn	GAC	Asp
T237	ACT	Thr	ACT	Thr	GCT	Ala
P237	CCT	Pro	ACT	Thr	GCT	Ala
N238	AAT	Asn	AAT	Asn	GAT	Asp
H238	CAC	His	AAC	Asn	GAC	Asp
A238	GCT	Ala	ACT	Thr	GCT	Ala
S239	TCT	Ser	ACT	Thr	GCT	Ala
Q238	CAG	Gln	AAG	Lys	GAG	Glu
K239	AAA	Lys	AAA	Lys	GAA	Glu
L247	CTT	Leu	ATT	Ile	GTT	Val
N248	AAC	Asn	AAC	Asn	GAC	Asp
H248	CAT	His	AAT	Asn	GAT	Asp
F249	TTC	Phe	ATC	Ile	GTC	Val
M250	ATG	Met	ATG	Met	GTG	Val
G251	GGT	Gly	AGT	Ser	GGT	Gly
V251	GTC	Val	ATC	Ile	GTC	Val

TABLE 13 continued (II)

Codon	Amino Acid	Codon	Amino Acid	Codon	Amino Acid
CAG	Gln	TAG	Stop	AAG	Lys
CAT	His	TAT	Tyr	AAT	Asn
CCT	Pro	TCT	Ser	CAT	His
CAC	His	TAC	Tyr	CAC	His
CCC	Pro	TCC	Ser	AAC	Asn
CCA	Pro	TCA	Ser	CAA	Gln
CAA	Gln	TAA	Stop	AAA	Lys
CTC	Leu	TTC	Phe	TAC	Tyr
CCA	Pro	TCA	Ser	GAA	Glu
CTC	Leu	TTC	Phe	GAC	Asp
CAT	His	TAT	Tyr	GAT	Asp
CTG	Leu	TTG	Leu	GAG	Glu
CCT	Pro	TCT	Ser	TAT	Tyr
CCG	Pro	TCG	Ser	GAG	Glu
CAT	His	TAT	Tyr	TAT	Tyr
CAT	His	TAT	Tyr	CAT	His
CTT	Leu	TTT	Phe	AAT	Asn
CCG	Pro	TCG	Ser	CAG	Gln
CTT	Leu	TTT	Phe	TAT	Tyr
CTC	Leu	TTC	Phe	CAC	His
CTG	Leu	TTG	Leu	CAG	Gln
CTG	Leu	TTG	Leu	TAG	Stop
CAG	Gln	TAG	Stop	CAG	Gln
CTT	Leu	TTT	Phe	TAT	Tyr
CCT	Pro	TCT	Ser	AAT	Asn
CAT	His	TAT	Tyr	TAT	Tyr
CTG	Leu	TTG	Leu	AAG	Lys
CTG	Leu	TTG	Leu	TAG	Stop
CAC	His	TAC	Tyr	AAC	Asn
CCT	Pro	TCT	Ser	AAT	Asn
CCT	Pro	TCT	Ser	CAT	His
CAT	His	TAT	Tyr	AAT	Asn
CAC	His	TAC	Tyr	CAC	His
CCT	Pro	TCT	Ser	GAT	Asp
CCT	Pro	TCT	Ser	TAT	Tyr
CAG	Gln	TAG	Stop	CAG	Gln
CAA	Gln	TAA	Stop	AAA	Lys
CTT	Leu	TTT	Phe	CAT	His
CAC	His	TAC	Tyr	AAC	Asn
CAT	His	TAT	Tyr	CAT	His
CTC	Leu	TTC	Phe	TAC	Tyr
CTG	Leu	TTG	Leu	AAG	Lys
CGT	Arg	TGT	Cys	GAT	Asp
CTC	Leu	TTC	Phe	GAC	Asp

TABLE 13 continued (III)

Codon	Amino Acid	Codon	Amino Acid	Codon	Amino Acid
AGG	Arg	ACG	Thr	ATG	Met
AGT	Ser	ACT	Thr	ATT	Ile
CGT	Arg	CCT	Pro	CTT	Leu
CGC	Arg	CCC	Pro	CTC	Leu
AGC	Ser	ACC	Thr	ATC	Ile
CGA	Arg	CCA	Pro	CTA	Leu
AGA	Arg	ACA	Thr	ATA	Ile
TGC	Cys	TCC	Ser	TTC	Phe
GGA	Gly	GCA	Ala	GTA	Val
GGC	Gly	GCC	Ala	GTC	Val
GGT	Gly	GCT	Ala	GTT	Val
GGG	Gly	GCG	Ala	GTG	Val
TGT	Cys	TCT	Ser	TTT	Phe
GGG	Gly	GCG	Ala	GTG	Val
TGT	Cys	TCT	Ser	TTT	Phe
CGT	Arg	CCT	Pro	CTT	Leu
AGT	Ser	ACT	Thr	ATT	Ile
CGG	Arg	CCG	Pro	CTG	Leu
TGT	Cys	TCT	Ser	TTT	Phe
CGC	Arg	CCC	Pro	CTC	Leu
CGG	Arg	CCG	Pro	CTG	Leu
TGG	Trp	TCG	Ser	TTG	Leu
CGG	Arg	CCG	Pro	CTG	Leu
TGT	Cys	TCT	Ser	TTT	Phe
AGT	Ser	ACT	Thr	ATT	Ile
TGT	Cys	TCT	Ser	TTT	Phe
AGG	Arg	ACG	Thr	ATG	Met
TGG	Trp	TCG	Ser	TTG	Leu
AGC	Ser	ACC	Thr	ATC	Ile
AGT	Ser	ACT	Thr	ATT	Ile
CGT	Arg	CCT	Pro	CTT	Leu
AGT	Ser	ACT	Thr	ATT	Ile
CGC	Arg	CCC	Pro	CTC	Leu
GGT	Gly	GCT	Ala	GTT	Val
TGT	Cys	TCT	Ser	TTT	Phe
CGG	Arg	CCG	Pro	CTG	Leu
AGA	Arg	ACA	Thr	ATA	Ile
CGT	Arg	CCT	Pro	CTT	Leu
AGC	Ser	ACC	Thr	ATC	Ile
CGT	Arg	CCT	Pro	CTT	Leu
TGC	Cys	TCC	Ser	TTC	Phe
AGG	Arg	ACG	Thr	ATG	Met
GGT	Gly	GCT	Ala	GTT	Val
GGC	Gly	GCC	Ala	GTC	Val

TABLE 13 continued (IV)

Codon	Amino Acid	Codon	Amino Acid	Codon	Amino Acid	Codon	Amino Acid
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
CCA	Pro	CCG	Pro	CCC	Pro	CCT	Pro
CAA	Gln	CAG	Gln	CAC	His	CAT	His
ACA	Thr	ACG	Thr	ACC	Thr	ACT	Thr
CCA	Pro	CCG	Pro	CCC	Pro	CCT	Pro
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
TTA	Leu	TTG	Leu	TTC	Phe	TTT	Phe
GCA	Ala	GCG	Ala	GCC	Ala	GCT	Ala
GTA	Val	GTG	Val	GTC	Val	GTT	Val
GAA	Glu	GAG	Glu	GAC	Asp	GAT	Asp
GTA	Val	GTG	Val	GTC	Val	GTT	Val
TCA	Ser	TCG	Ser	TCC	Ser	TCT	Ser
GCA	Ala	GCG	Ala	GCC	Ala	GCT	Ala
TAA	Stop	TAG	Stop	TAC	Tyr	TAT	Tyr
CAA	Gln	CAG	Gln	CAC	His	CAT	His
ATA	Ile	ATG	Met	ATC	Ile	ATT	Ile
CCA	Pro	CCG	Pro	CCC	Pro	CCT	Pro
TTA	Leu	TTG	Leu	TTC	Phe	TTT	Phe
CTA	Leu	CTG	Leu	CTC	Leu	CTT	Leu
CTA	Leu	CTG	Leu	CTC	Leu	CTT	Leu
TTA	Leu	TTG	Leu	TTC	Phe	TTT	Phe
CAA	Gln	CAG	Gln	CAC	His	CAT	His
TTA	Leu	TTG	Leu	TTC	Phe	TTT	Phe
ACA	Thr	ACG	Thr	ACC	Thr	ACT	Thr
TAA	Stop	TAG	Stop	TAC	Tyr	TAT	Tyr
ATA	Ile	ATG	Met	ATC	Ile	ATT	Ile
TTA	Leu	TTG	Leu	TTC	Phe	TTT	Phe
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
ACA	Thr	ACG	Thr	ACC	Thr	ACT	Thr
CCA	Pro	CCG	Pro	CCC	Pro	CCT	Pro
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
CAA	Gln	CAG	Gln	CAC	His	CAT	His
GCA	Ala	GCG	Ala	GCC	Ala	GCT	Ala
TCA	Ser	TCG	Ser	TCC	Ser	TCT	Ser
CAA	Gln	CAG	Gln	CAC	His	CAT	His
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
CTA	Leu	CTG	Leu	CTC	Leu	CTT	Leu
AAA	Lys	AAG	Lys	AAC	Asn	AAT	Asn
CAA	Gln	CAG	Gln	CAC	His	CAT	His
TTA	Leu	TTG	Leu	TTC	Phe	TTT	Phe
ATA	Ile	ATG	Met	ATC	Ile	ATT	Ile
GGA	Gly	GGG	Gly	GGC	Gly	GGT	Gly
GTA	Val	GTG	Val	GTC	Val	GTT	Val

TABLE 14 Amino acid mutations at target sites in rt

Target	Mutation
K32	M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L
N33	D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R
P34	S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F
H35	I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G
T37	W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S
P59	S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F
K60	M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L
F61	P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M
A62	R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V
V63	A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y
D83	C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N
V84	A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y
S85	T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P
A86	R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V
Y89	V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W
H90	I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G
I/L91	K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H
P177	S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F
F178	P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M
L179	K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I
L180	K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I
A181	R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V
Q183	E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C
F183	P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M
T184	W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S
Y203	V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W
M204	F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K
L235	K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I
N236	D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R
T237	W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S
P237	S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F
N238	D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R
H238	I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G
A238	R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V
S239	T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P
Q238	E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C
K239	M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L
L247	K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I
N248	D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R
H248	I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G

- 106 -

Target	Mutation
F249	P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M
M250	F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K
G251	H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E
V251	A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y

TABLE 15 Summary of HBV mutations in Patient F treated with ADV

Sample Name	Genotype	Polymerase*	Surface
CAP 01564808	A	rtL157L/M rtA181V rtV207I rtN236T	sF83S sL173F sW199L

5

\* Nomenclature according to Stuyver *et al.*, 2001, *supra*.

\*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.



**TABLE 16** Summary of HBV mutations in Patient G treated with ADV

Sample Name	Genotype	Polymerase*	Surface
KAN 02510355	C	rtL80V rtP109S rtI163V rtM204I rtL229M rtN/H/A/S/Q238K	sI126T sK160R sS174N sA184V sW196L sS210N sF/C220L sY221C

\* Nomenclature according to Stuyver *et al.*, 2001, *supra*.

- 5 \*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.

**TABLE 17** Summary of HBV mutations in Patient H treated with ADV

Sample Name	Genotype	Polymerase*	Surface
LAV0303	D	rtS78S/T rtN118N/S rtN139N/K rtV142E rtA181A/T rtI204M rtQ/P/S/Stop215Q rtE218K/E rtN238N/H	sC69Stop/C sC76Y sI110V/I sY134N sW172Stop/W sW196Stop sS207R

10

\* Nomenclature according to Stuyver *et al.*, 2001, *supra*.

- \*\* Mutations in bold have not been detected in reference HBV genotypes, mutations not in bold are changes from the previous sample that are present in reference genotypes.

- 108 -

**TABLE 18** *In vitro* drug susceptibility of the HBV reference laboratory strain and patient-derived HBV isolate

Mutation	<i>In vitro</i> Susceptibility IC <sub>50</sub> (fold change from wild-type)		
	Real-time PCR	Southern Blot	
	Adefovir	Adefovir	Lamivudine
Wild-type (pPC)	1	1	1
rt N236T/N238D	23	NA <sup>1</sup>	NA <sup>1</sup>
rt A181T/N236T/N238D	5.1	7.3	>100
rt L180M/M204V <sup>2</sup>	NT <sup>5</sup>	0.9	>2500

5 <sup>1</sup> NA, not analyzed.

<sup>2</sup> Data from Delaney *et al.*, 2001, *supra*

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood  
 10 that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features

**BIBLIOGRAPHY**

- Allen *et al.*, *Hepatology* 27(6): 1670-1677, 1998
- Aye *et al.*, *J. Hepatol.* 26: 1148-1153, 1997
- Bartholomeusz *et al.*, *Intervirology* 40(5-6): 337-342 1997
- Benhamou *et al.*, *Lancet* 358: 718-723, 2001
- Benzaria *et al.*, *J Med Chem.* 39: 4958-4965, 1996
- Boyd *et al.*, *Antiviral Chem Chemother.* 32: 358-363, 1987
- Calio *et al.*, *Antiviral Res.* 23: 77-89, 1994
- Das *et al.*, *J. Virol.* 75(10): 4771-4779, 2001
- Delaney *et al.*, *Antimicrob Agents Chemother* 45(6): 1705-1013, 2001
- Dienstag *et al.*, *New England J Med* 333: 1657-1661, 1995
- Frick *et al.*, *Antimicrob. Agents Chemother.* 37: 2285-2292, 1993
- Gaillard *et al.*, *Antimicrob Agents Chemother.* 46(4): 1005-1013, 2002
- Gilson *et al.*, *J Viral Hepat* 6: 387-395, 1999
- Heathcote *et al.*, *Hepatology* 28: A620, 1998
- Hendricks *et al.*, *Am J Clin Pathol* 104: 537-46, 1995
- Kruger *et al.*, *Hepatology* 22: 219A, 1994
- Main *et al.*, *J. Viral Hepatitis* 3: 211-215, 1996
- Norder *et al.*, *(J. Gen. Virol.* 74: 341-1348, 1993
- Perrillo *et al.*, *Hepatology* 32: 129-134, 2000
- Peters *et al.*, *Transplantation* 68: 1912-1914, 1999
- Price *et al.*, *Proc. Natl. Acad. Sci. USA* 86(21): 8541-8544, 1989
- Ren and Nassal, *J. Virol.* 75(3): 1104-1116, 2001
- Severini *et al.*, *Antimicrobial Agents Chemother.* 39: 430-435, 1995
- Stuyver *et al.*, *Hepatology* 33: 751-757, 2001
- Summers and Mason, *Cell* 29: 403-415, 1982
- Suo *et al.*, *J Biol Chem.* 273(42): 27250-27258. 1998
- Vere Hodge, *Antiviral Chem Chemother* 4: 67-84, 1993
- Xiong *et al.*, *Hepatology.* 28(6): 1669-73, 1998
- Ying *et al.*, *J Viral Hepat.* 7(2): 161-165, 2000

- 110 -

Ying *et al.*, *J. Viral Hepat.* 7(1): 79-83, 2000

Ying *et al.*, *Viral Hepat.* 7(2): 161-165, 2000

- 111 -

**CLAIMS**

1. An isolated HBV variant wherein said variant comprises a nucleoside mutation in a gene encoding a DNA polymerase resulting in at least one amino acid addition, substitution and/or deletion to said DNA polymerase and wherein said variant exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combinations thereof
2. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV.
3. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to both LMV.
4. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to TFV.
5. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to FTC.
6. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and LMV.
7. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and TFV.
8. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to LMV and TFV.

- 112 -

9. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and FTC.
10. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to LMV and FTC.
11. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to TFV and FTC.
12. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and LMV and TFV.
13. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and LMV and FTC.
14. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to FTC and LMV and TFV.
15. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and FTC and TFV.
16. The isolated HBV variant of Claim 1 wherein said variant exhibits decreased sensitivity to ADV and LMV and TFV and FTC.
17. The isolated HBV variant of any one of Claims 1 to 16 wherein said variant exhibits reduced interactivity to an immunological reagent specific to HBsAg.
18. The isolated HBV variant of Claim 1 wherein said variant comprises a mutation in domain F of the HBV DNA polymerase thereby conferring an altered amino acid sequence to the sequence set forth in Formula I [SEQ ID NO:1]:

- 113 -

**FORMULA I**

L, X<sub>1</sub>, X<sub>2</sub>, D, W, G, P, C, X<sub>3</sub>, X<sub>4</sub>, H, G, X<sub>5</sub>, H, X<sub>6</sub>, I, R, X<sub>7</sub>, P, R, T, P, X<sub>8</sub>, R, V, X<sub>9</sub>, G, G, V, F, L, V, D, K, N, P, H, N, T, X<sub>10</sub>, E, S, X<sub>11</sub>, L, X<sub>12</sub>, V, D, F, S, Q, F, S, R, G, X<sub>13</sub>, X<sub>14</sub>, X<sub>15</sub>, V, S, W, P, K, F, A, V, P, N, L, X<sub>16</sub>, S, L, T, N, L, L, S\*

wherein:

- X<sub>1</sub> is L or R or I;
- X<sub>2</sub> is E or D;
- X<sub>3</sub> is T or D or A or N or Y;
- X<sub>4</sub> is E or D;
- X<sub>5</sub> is E or K or Q;
- X<sub>6</sub> is H or R or N;
- X<sub>7</sub> is I or T;
- X<sub>8</sub> is A or S;
- X<sub>9</sub> is T or R;
- X<sub>10</sub> is A or T or S;
- X<sub>11</sub> is R or T;
- X<sub>12</sub> is V or G;
- X<sub>13</sub> is S or I or T or N or V;
- X<sub>14</sub> is T or S or H or Y;
- X<sub>15</sub> is R or H or K or Q;
- X<sub>16</sub> is Q or P;

and wherein S\* is designated as amino acid 74.

19. The isolated HBV variant of Claim 1 wherein said variant comprises a mutation in any one of domains A through E thereby conferring an altered amino acid sequence to the sequence set forth in Formula II [SEQ ID NO:2]:

- 114 -

## FORMULA II

$SX_1LSWLSLDVSAAFYHX_2PLHPAAMPHELLX_3GSSGLX_4RYV$   
 $ARLSSX_5SX_6X_7XNX_8QX_9X_{10}XXXXX_{11}LHX_{12}X_{13}CSRX_{14}LYVSLX_{15}$   
 $LLYX_{16}TX_{17}GX_{18}KLHLX_{19}X_{20}HPIX_{21}LGFRKX_{22}PMGX_{23}GLSPFL$   
 $LAQFTSAIX_{24}X_{25}X_{26}X_{27}X_{28}RAFX_{29}HCX_{30}X_{31}FX_{32}YM^*DDX_{33}VLGA$   
 $X_{34}X_{35}X_{36}X_{37}HX_{38}EX_{39}LX_{40}X_{41}X_{42}X_{43}X_{44}X_{45}X_{46}LLX_{47}X_{48}GIHLNPX_{49}K$   
 $TKRWGYSLNFMGYX_{50}IG$

wherein:

- X is any amino acid
- $X_1$  is N or D;
- $X_2$  is I or P;
- $X_3$  is I or V;
- $X_4$  is S or D;
- $X_5$  is T or N;
- $X_6$  is R or N;
- $X_7$  is N or I;
- $X_8$  is N or Y or H;
- $X_9$  is H or Y;
- $X_{10}$  is G or R;
- $X_{11}$  is D or N;
- $X_{12}$  is D or N;
- $X_{13}$  is S or Y;
- $X_{14}$  is N or Q;
- $X_{15}$  is L or M;
- $X_{16}$  is K or Q;
- $X_{17}$  is Y or F;
- $X_{18}$  is R or W;



- 115 -

X<sub>19</sub> is Y or L;  
X<sub>20</sub> is S or A;  
X<sub>21</sub> is I or V;  
X<sub>22</sub> is I or L;  
X<sub>23</sub> is V or G;  
X<sub>24</sub> is C or L;  
X<sub>25</sub> is A or S;  
X<sub>26</sub> is V or M;  
X<sub>27</sub> is V or T;  
X<sub>28</sub> is R or C;  
X<sub>29</sub> is F or P;  
X<sub>30</sub> is L or V;  
X<sub>31</sub> is A or V;  
X<sub>32</sub> is S or A;  
X<sub>33</sub> is V or L or M;  
X<sub>34</sub> is K or R;  
X<sub>35</sub> is S or T;  
X<sub>36</sub> is V or G;  
X<sub>37</sub> is Q or E;  
X<sub>38</sub> is L or S or R;  
X<sub>39</sub> is S or F;  
X<sub>40</sub> is F or Y;  
X<sub>41</sub> is T or A;  
X<sub>42</sub> is A or S;  
X<sub>43</sub> is V or I;  
X<sub>44</sub> is T or C;  
X<sub>45</sub> is N or S;  
X<sub>46</sub> is F or V;  
X<sub>47</sub> is S or D;  
X<sub>48</sub> is L or V;  
X<sub>49</sub> is N or Q;

- 116 -

X<sub>50</sub> is V or I; and

M<sup>\*</sup> is amino acid 204;

and wherein the first S is designated as amino acid 75.

20. The isolated HBV variant of Claim 18 or 19 wherein said variant further comprises an altered HBsAg.

21. An isolated HBV variant comprising a mutation in the nucleotide sequence encoding HBsAg resulting in an amino acid addition, substitution and/or deletion in said HBsAg in a region corresponding to the amino acid sequence set forth in SEQ ID NO:1 or SEQ ID NO:2 and wherein said variant exhibits decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and/or optionally other nucleoside or nucleotide analogs or other anti-HBV agents or combinations thereof.

22. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV.

23. The isolated HBV variant of Claim 21 wherein said variants exhibits decreased sensitivity to LMV.

24. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to TFV.

25. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to FTC.

26. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased

- 117 -

sensitivity to ADV and LMV.

27. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to LMV and TFV.

28. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV and FTC.

29. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to LMV and FTC.

30. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to TFV and FTC.

31. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV and TFV.

32. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV and LMV and TFV.

33. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV and LMV and FTC.

34. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to FTC and LMV and TFV.

35. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV and FTC and TFV.

36. The isolated HBV variant of Claim 21 wherein said variant exhibits decreased sensitivity to ADV and LMV and TFV and FTC.

- 118 -

37. The isolated HBV variant of Claim 21 wherein an antibody specific for a wild-type HBsAg exhibits a reduced capacity to neutralize said HBV variant and wherein said HBV variant is selected by exposure of a subject to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, in single or combinations or sequential therapy.

38. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtS21A, rtL122F, rtN124H, rtT28N, rtP130Q, rtD131N and rtY135C or a combination thereof or an equivalent mutation.

39. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M or a combination thereof or an equivalent mutation.

40. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D or a combination thereof or an equivalent mutation.

41. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtH90D and rtL/F108L or a combination thereof or an equivalent mutation.

42. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtL157L/M, rtA181V and rtV207I or a combination thereof or an equivalent mutation.

43. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV

- 119 -

DNA polymerase selected from rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K or a combination thereof or an equivalent mutation.

44. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the surface antigen selected from sP120T, sM125F and sT127A or a combination thereof or an equivalent mutation.

45. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the surface antigen selected from sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C or a combination thereof or an equivalent mutation.

46. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the surface antigen selected from sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop or a combination thereof or an equivalent mutation.

47. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV surface antigen selected from sI81M and sP214Q or a combination thereof or an equivalent mutation.

48. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV surface antigen selected from sF83S, sL173F and sW199L or a combination thereof or an equivalent mutation.

49. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV surface antigen selected from sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C or a combination thereof or an equivalent mutation.

50. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV surface antigen selected from sC69Stop/C, sC76Y, sI110V/L, sY134N, sW172Stop/W, sW196Stop and sS207R or a combination thereof or an equivalent mutation.

- 120 -

51. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtK32, rtN33, rtP34, rtH35 and rtT37 or a combination thereof or an equivalent mutation.

52. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtP59, rtK60, rtF61, rtA62 and rtV63 or a combination thereof or an equivalent mutation.

53. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91 or a combination thereof or an equivalent mutation.

54. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184 or a combination thereof or an equivalent mutation.

55. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtM204 and rtY203 or a combination thereof or an equivalent mutation.

56. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rt235, rt236, rt237, rt238 and rt239 or a combination thereof or an equivalent mutation.

57. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rt247, rt248, rt249, rt250 and rt251 or a combination thereof or an equivalent mutation.

58. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;

- 121 -

P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;

- 122 -

A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion.

59. A method for determining the potential for an HBV to exhibit reduced sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV or optionally other nucleoside analogs or other anti-HBV agents, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV DNA polymerase resulting in at least one amino acid substitution, deletion and/or addition in any one or more of domains F and A through E or a region proximal thereto of said DNA polymerase and associated with resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, wherein the presence of such a mutation is an indication of the likelihood of resistance to said ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV.

60. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to



- 123 -

ADV.

61. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to LMV.
62. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to TFV.
63. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to FTC.
64. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV and LMV.
65. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV and TFV.
66. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to LMV and TFV.
67. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV and FTC.
68. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to TFV and FTC.
69. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to LMV and FTC.
70. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV and LMV and TFV.
71. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV

- 124 -

and LMV and FTC.

72. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to LMV and TFV and FTC.

73. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV and FTC and TFV.

74. The method of Claim 59 wherein the HBV exhibits reduced sensitivity to ADV and LMV and TFV and FTC.

75. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtS21A, rtL122F, rtN124H, rtT28N, rtP130Q, rtD131N and rtY135C or a combination thereof or an equivalent mutation.

76. The isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M or a combination thereof or an equivalent mutation.

77. The isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D or a combination thereof or an equivalent mutation.

78. The isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A or a combination thereof or an equivalent mutation.

- 125 -

79. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtH90D and rtL/F108L or a combination thereof or an equivalent mutation.

80. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected rtL157L/M, rtA181V and rtV207I or a combination thereof or an equivalent mutation.

81. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K or a combination thereof or an equivalent mutation.

82. The isolated HBV variant of Claim 1 or 37 comprising a mutation in the DNA polymerase selected from rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H or a combination thereof or an equivalent mutation.

83. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the surface antigen gene selected from sP120T, sM125F and sT127A or a combination thereof or an equivalent mutation.

84. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV Surface antigen gene selected from sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C or a combination thereof or an equivalent mutation.

85. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the surface antigen selected from sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop or a combination thereof or an equivalent mutation.

86. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA surface antigen selected from sN40S, sC69Stop, sM75I,

- 126 -

sL88P, sT118A, sW182Stop, sW196L, sY206H an sY225F or a combination thereof or an equivalent mutation.

87. An isolated DNA molecule from an HBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA surface antigen selected from s181M and sP214Q or a combination thereof or an equivalent mutation.

88. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA surface antigen selected from sF83S, sL173F and sW199L or a combination thereof or an equivalent mutation.

89. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA surface antigen selected from sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C or a combination thereof or an equivalent mutation.

90. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA surface antigen selected from sC69Stop/C, sC76Y, sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R or a combination thereof or an equivalent mutation.

91. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtK32, rtN33, rtP34, rtH35 and rtT37 or a combination thereof or an equivalent mutation.

92. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtP59, rtK60, rtF61, rtA62 and rtV63 or a combination thereof or an equivalent mutation.

93. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtD83, rtV84, rtS85, rtA86, rtY89,

- 127 -

rtH90 and rtI/L91 or a combination thereof or an equivalent mutation.

94. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184 or a combination thereof or an equivalent mutation.

95. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rtM204 and rtY203 or a combination thereof or an equivalent mutation.

96. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rt235, rt236, rt237, rt238 and rt239 or a combinations thereof or an equivalent mutation.

97. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from rt247, rt248, rt249, rt250 and rt251 or a combinations thereof or an equivalent mutation.

98. An isolated DNA molecule from anHBV variant of Claim 1 or 37 comprising a mutation in the HBV DNA polymerase selected from K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion; P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion; H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion; T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion; P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion; K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion; A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion; V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion; D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;

- 128 -

V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;

- 129 -

G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion.

99. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtS21A, rtL122F, rtN124H, rtT28N, rtP130Q, rtD131N and rtY135C or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

100. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

101. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and

- 130 -

LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

102. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and optionally other nucleoside analogs.

103. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtH90D and rtL/F108L or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

104. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtL157L/M, rtA181V and rtV207I or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and



- 131 -

optionally other nucleoside analogs.

105. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

106. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

107. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sP120T, sM125F and sT127A or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally

- 132 -

other nucleoside analogs.

108. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

109. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

110. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182Stop, sW196L, sY206H and sY225F or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

- 133 -

111. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of s181M and sP214Q or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

112. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sF83S, sL173F and sW199L or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

113. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

- 134 -

114. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of sC69Stop/C, sC76Y, sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

115. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtK32, rtN33, rtP34, rtH35 and rtT37 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

116. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtP59, rtK60, rtF61, rtA62 and rtV63 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

117. A method for determining whether an HBV strain exhibits reduced sensitivity

- 135 -

to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

118. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

119. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rtM204 and rtY203 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

120. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA

- 136 -

from said HBV and screening for a mutation wherein the presence of rt235, rt236, rt237, rt238 and rt239 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

121. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of rt247, rt248, rt249, rt250 and rt251 or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside analogs.

122. A method for determining whether an HBV strain exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation wherein the presence of K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;

- 137 -

D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/YV/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;

- 138 -

M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and optionally other nucleoside analogs.

123. A method for detecting an agent which exhibits inhibitory activity to an HBV which exhibits resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and optionally other nucleoside or nucleotide analogs or other anti-HBV agents, said method comprising:

generating a genetic construct comprising a replication competent-effective amount of the genome from said HBV contained in a plasmid vector and then transfecting said cells with said construct;

contacting said cells, before, during and/or after transfection, with the agent to be tested;

culturing said cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agent; and

subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of said agent.



- 139 -

124. A method for detecting an agent which exhibits inhibitory activity to an HBV which exhibits resistance or decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV, and optionally other nucleoside or nucleotide analogs or other anti-HBV agents, said method comprising:

generating a genetic construct comprising a replication competent-effective amount of the genome from said HBV contained in or fused to an amount of a baculovirus genome effective to infect cells and then infecting said cells with said construct;

contacting said cells, before, during and/or after infection, with the agent to be tested;

culturing said cells for a time and under conditions sufficient for the HBV to replicate, express genetic sequences and/or assemble and/or release virus or virus-like particles if resistant to said agent; and

subjecting the cells, cell lysates or culture supernatant fluid to viral- or viral-component-detection means to determine whether or not the virus has replicated, expressed genetic material and/or assembled and/or been released in the presence of said agent.

125. The method of Claim 123 or 124 wherein the HBV genome is stably integrated into the cells' genome.

126. An agent identified by the method of any one of Claims 123 to 124.

127. Use of an HBV variant according to any one of Claims 1 to 59 or a component thereof in the rational design of an anti-HBV agent.

- 140 -

128. Use according to Claim 126 wherein the rational design comprises microarray analysis.

129. Use according to Claim 126 wherein the rational design is based on the crystal structure or NMR structure of a viral component.

130. A vaccine comprising an antigenic component of the HBV variant of any one of Claims 1 to 59 or an antigenic component thereof or an antibody thereto.

131. The vaccine of Claim 130 wherein the antigenic component is an HBsAg or PreS1 or PreS2.

132. The vaccine of Claim 130 wherein the antigenic component is a defective HBV variant.

133. The vaccine of Claim 130 comprising an antibody to HBsAg or PreS1 or PreS2

134. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtS21A, rtL122F, rtN124H, rtT28N, rtP130Q, rtD131N and rtY135C.

135. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V and rtI235I/M.

136. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F.

- 141 -

137. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D.

138. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A.

139. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtH90D and rtL/F108L.

140. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtL157L/M, rtA181V and rtV207I.

141. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtL80V, rtP109S, rtI163V, rtL229M and rtN/H/A/S/Q238K.

142. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E and rtN238N/H.

143. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sP120T, sM125F and sT127A.

144. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sT118R, sM133T, sF134V, sI195M, sS207R and sY225Y/C.

- 142 -

145. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sS126T, sM133L/M, sS143S/T, sD144A, sG145A and sW172Stop.

146. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sN40S, sC69Stop, sM75I, sL88P, sT118A, sW182Stop, sW196L, sY206H and sY225F.

147. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sI81M and sP214Q.

148. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sF83S, sL173F and sW199L.

149. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C.

150. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from sC69Stop/C, sC76Y, sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R.

151. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtK32, rtN33, rtP34, rtH35 and rtT37.

152. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtP59, rtK60, rtF61, rtA62 and rtV63.

153. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 and rtI/L91.

- 143 -

154. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184.

155. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rtM204 and rtY203.

156. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rt235, rt236, rt237, rt238 and rt239.

157. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from rt247, rt248, rt249, rt250 and rt251.

158. The vaccine of Claim 130 wherein the HBV or its antigenic compound is from an HBV variant having a mutation selected from,

K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion;  
V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion;  
S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;

- 144 -

H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion;  
P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion.

- 145 -

159. A computer product for assessing the likely usefulness of a viral variant or biological sample comprising same for determining an appropriate therapeutic protocol in a subject, said product comprising:

- (1) code that receives as input index values ( $I_{vs}$ ) for at least two features associated with said viral agents or biological sample comprising same, wherein said features are selected from:
  - (a) the ability to exhibit resistance for reduced sensitivity to a particular compound or immunological agent;
  - (b) an altered DNA polymerase from wild-type HBV;
  - (c) an altered surface antigen from wild-type HBV;
  - (d) morbidity or recovery potential of a patient; or
  - (e) altered replication capacity (increased or decreased);
- (2) code that adds said  $I_{vs}$  to provide a sum corresponding to a potency value ( $P_v$ ) for said viral variants or biological samples; and
- (3) a computer readable medium that stores the codes.

160. A computer for assessing the likely usefulness of a viral variant or biological sample comprising same in a subject, wherein said computer comprises:

- (1) a machine-readable data storage medium comprising a data storage material encoded with machine-readable data, wherein said machine-readable data comprise  $I_{vs}$  for at least two features associated with said viral variant or biological sample; wherein said features are selected from:-
  - (a) the ability to exhibit resistance for reduced sensitivity to a particular compound or immunological agent;

- 146 -

- (b) an altered DNA polymerase from wild-type HBV;
  - (c) an altered surface antigen from wild-type HBV;
  - (d) morbidity or recovery potential of a patient; or
  - (e) altered replication capacity (increased or decreased);
- (2) a working memory for storing instructions for processing said machine-readable data;
- (3) a central-processing unit coupled to said working memory and to said machine-readable data storage medium, for processing said machine readable data to provide a sum of said  $I_{VS}$  corresponding to a  $P_V$  for said compound(s); and
- (4) an output hardware coupled to said central processing unit, for receiving said  $P_V$ .

161. A composition comprising an agent capable of directly or indirectly inhibiting an HBV variant as defined in any one of Claims 1 to 59, said composition further comprising one or more pharmaceutically acceptable carriers and/or diluents.

162. The composition of Claim 161 wherein the agent is a recombinant protein from said HBV variant.

163. The composition of Claim 161 wherein the recombinant protein is HBsAg or PreS1 or PreS2.

164. The composition of Claim 161 wherein the agent is capable of inhibiting an HBV variant polymerase.

165. The composition of Claim 161 wherein the agent is identified by natural product screening or rational drug design.



- 147 -

166. The composition of Claim 161 wherein the agent is a defective HBV variant.
167. The composition of Claim 161 wherein the agent is an antibody directed to an HBV compound.
168. The composition of Claim 161 wherein the agent is a ribozyme, antisense molecule or sense molecule relative to an HBV gene.
169. A method according to Claims 37 to 57 wherein a virus related to HBV from the family of hepadnaviruses such as WHV or DHBV exhibits reduced sensitivity to a nucleoside analog, said method comprising isolating DNA or corresponding mRNA from said HBV, or DHBV or WHV and screening for a mutation or combinations thereof or an equivalent one or more other mutation is indicative of a variant wherein said variant exhibits a decreased sensitivity to ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV.
170. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.
171. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA

- 148 -

or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sP120T, sM125T and sT127A selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

172. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sT118R, sM133T, sF134V sI195M, sS207R and sY225Y/C selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

173. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sS126T, sM133L/M, sS143S/T, sD144A sG145A and sW172Stop selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

- 149 -

174. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sN40S, and sC69 Stop, sM75L, sL88P, sT118A, sW182stop, sW196L, sY206H and sY225F selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

175. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sF83S, sL173F and sW199L selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

176. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and

- 150 -

LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

177. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from sC69Stop/C, sC76Y sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

178. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

179. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtS21A, rtL122F, rtN124H, rtH126R,

- 151 -

rtT128N, rtP130Q, rtD131N, rtY135C selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

180. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtN/S/T/I/V53D, rtY126Q, rtL180M, rtS202G, rtI204V, rtI235I/M selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

181. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtN53D, rtY54H, rtS57P, rtL91I, rtS116P, rtF122L, rtY124H, rtV134D, rtY141Y/F, rtL145M, rtF151F/Y, rtA181T, rtK212R, rtL217R, rtS219A, rtN236T and rtN238D selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

- 152 -

182. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtS78T, rtV84M, rtY126C, rtV191I, rtM204I and rtV214A selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

183. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtH90D and rtL/F108L selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

184. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtL157L/M, rtA181V, rtV207I; and rtN236T selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing

interactivity of said antibodies to said mutated surface antigen.

185. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtL80V, rtP109S, rtI163V, rtM204I, rtL229M and rtN/H/A/S/Q238K selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

186. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtS78S/T, rtN118N/S, rtN139N/K, rtV142E, rtA181A/T, rtI204M, rtQ/P/S/Stop215Q, rtE218K/E, and rtN238N/H selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

187. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtK32, rtN33, rtP34, rtH35 and rtT37 selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV,

- 154 -

LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

188. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtP59, rtK60, rtF61, rtA62 and rtV63 elected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

189. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtD83, rtV84, rtS85, rtA86, rtY89, rtH90 selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

190. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the



- 155 -

presence of such a mutation is selected from rtP177, rtF178, rtL179, rtL180, rtA181, rtQ182, rtF183 and rtT184 selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

191. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rtM204 and rtY203 selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

192. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rt235, rt236, rt237, rt238 and rt239 selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

193. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA

- 156 -

or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation is selected from rt247, rt248, rt249, rt250 and rt251 selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

194. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolating DNA or corresponding mRNA from said HBV and screening for a mutation in the nucleotide sequence encoding HBV surface antigen and/or HBV DNA polymerase genes wherein the presence of such a mutation, K32M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; N33D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion; P34S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion; H35I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion; T37W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion; P59S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion; K60M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion; F61P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion; A62R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion; V63A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion; D83C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/deletion; V84A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion; S85T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion; A86R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion; Y89V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion; H90I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion; I/L91K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/deletion; P177S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;

- 157 -

F178P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 L179K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 L180K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 A181R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 Q183E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 F183P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 T184W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 Y203V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/deletion;  
 M204F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 L235K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N236D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 T237W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/deletion;  
 P237S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/deletion;  
 N238D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H238I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 A238R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/deletion;  
 S239T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/deletion;  
 Q238E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/deletion;  
 K239M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/deletion;  
 L247K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/deletion;  
 N248D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/V/A/R/deletion;  
 H248I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/deletion;  
 F249P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/M/deletion;  
 M250F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/G/H/I/L/K/deletion;  
 G251H/I/L/K/M/F/P/S/T/W/Y/V/A/R/N/D/C/Q/E/deletion; and  
 V251A/R/N/D/C/Q/E/G/H/I/L/K/M/F/P/S/T/W/Y/deletion, is selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

- 158 -

195. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV and is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

196. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation is selected from sP120T, sM125T and sT127A selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

197. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected from sT118R, sM133T, sF134V sI195M, sS207R and sY225Y/C selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

- 159 -

198. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected from sS126T, sM133L/M, sS143S/T, sD144A sG145A and sW172Stop selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

199. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected from sN40S, and sC69 Stop, sM75I, sL88P, sT118A, sW182stop, sW196L, sY206H and sY225F selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

200. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected from sF83S, sL173F and sW199L selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen

- 160 -

201. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected from sI126T, sK160R, sS174N, sA184V, sW196L, sS210N, sF/C220L and sY221C selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

202. A method for determining the potential for an HBV to exhibit reduced interactivity to antibody to HBV surface antigen, said method comprising isolated protein from said HBV encoding HBV surface antigen wherein the presence of such a mutation was selected sC69Stop/C, sC76Y sI110V/I, sY134N, sW172Stop/W, sW196Stop and sS207R selected during treatment with ADV, LMV, TFV, or FTC, or ADV and LMV, ADV and TFV, LMV and TFV, FTC and ADV, FTC and TFV, FTC and LMV, or ADV and LMV and TFV, or ADV and FTC and TFV, TFV and FTC and LMV, ADV and LMV and FTC, or ADV and FTC and LMV and TFV is an indication of the likelihood of reducing interactivity of said antibodies to said mutated surface antigen.

1/32




Figure 1

2/32





Figure 2



3/32

**Figure 3**

[illegible]

## Figure 4

5/32

IL1A1 P, A-E	781	TGGCTCAGTTTACTAGTGGCCATTGTTTCAGTGGTTCGTAGGGCTTTCCCCACATGTTTGGCTTTTCAGTTATATGATGATGTGGGGGCCAAGTC	880	[SEQ ID NO:8]
IL1A2 P, A-E	789	TGGCTCAGTTTACTAGTGGCCATTGTTTCAGTGGTTCGTAGGGCTTTCCCCACATGTTTGGCTTTTCAGTTATATGATGATGTGGGGGCCAAGTC	888	[SEQ ID NO:9]
IL1A3 P, A-E	801	TGGCTCAGTTTACTAGTGGCCATTGTTTCAGTGGTTCGTAGGGCTTTCCCCACATGTTTGGCTTTTCAGTTATATGATGATGTGGGGGCCAAGTC	900	[SEQ ID NO:10]
IL1A4 P, A-E	774	TGGCTCAGTTTACTAGTGGCCATTGTTTCAGTGGTTCGTAGGGCTTTCCCCACATGTTTGGCTTTTCAGTTATATGATGATGTGGGGGCCAAGTC	873	[SEQ ID NO:11]
IL1A1 P, A-E	881	TGTAYAGCAVCTTGGAGTCCCTTTTACCGCTGTTACCAATTTCTTTTGTCTTTTGGGTATACATTTAAACCCCTAACAAAACTTAAAGATGGGGTTACTCT	980	
IL1A2 P, A-E	889	TGTACAGCATCTTGGAGTCCCTTTTACCGCTGTTACCAATTTCTTTTGTCTTTTGGGTATACATTTAAACCCCTAACAAAACTTAAAGATGGGGTTACTCT	988	
IL1A3 P, A-E	901	TGTACAGCATCTTGGAGTCCCTTTTACCGCTGTTACCAATTTCTTTTGTCTTTTGGGTATACATTTAAACCCCTAACAAAACTTAAAGATGGGGTTACTCT	1000	
IL1A4 P, A-E	874	TGTACAGCATCTTGGAGTCCCTTTTACCGCTGTTACCAATTTCTTTTGTCTTTTGGGTATACATTTAAACCCCTAACAAAACTTAAAGATGGGGTTACTCT	973	
IL1A1 P, A-E	981	TTACATTTTCANGGNTATGTCATGGATGTTATGGGTCAATGCCACAGATCACATCATACAGAAATCAAGATGGTTT	1060	
IL1A2 P, A-E	989	CTAAATTTTATGGGTTATGTCATGGATGTTATGGGTCTTGG	1030	
IL1A3 P, A-E	1001	CTAAATTTTATGGGTTATGTCATGGATGTTATGGGTCTTGGCACAAGACACATCATACAAAAATCAAGATG	1077	
IL1A4 P, A-E	974	TTAAATTTTCANGGNTATGTCATGGATGTTATGGGTCTTGG	1010	

Figure 4 (continued)

Patient A polymerase amino acid sequence alignment

Pol Trans Pre 1	KLAKSASSIXQSPV	RXAAYPAVSTFEKHSSSGHA	VEXXNL	PNSXBSQXERPV	PCWWLQ	FRNSKPCSDYCLSH	IVNLL	DWGPCAEHGEHH	93	[SEQ ID NO:12]													
Pol Trans 2	HTTNPASKASCLHQ	SPVKAAYPAVSTFEKHSSSGHA	VEFFNL	PNSARBSQSERPV	PCWWLQ	FRNSKPCSDYCLSH	IVNLL	DWGPCAEHGEHH	96	[SEQ ID NO:13]													
Pol Trans 3	LAQGIILQNFASKASCLHQ	SPVKAAYPAVSTFEKHSSSGHA	VEFFNL	PNSARBSQSERPV	PCWWLQ	FRNSKPCSDYCLSH	IVNLL	DWGPCAEHGEHH	100	[SEQ ID NO:14]													
Pol Trans 4	ASKSASSIYQSPVGTAAYPAVSTFEKHSSSGHA	VEFFNL	PNSARBSQSERPV	PCWWLQ	FRNSKPCSDYCLSH	IVNLL	DWGPCAEHGEHH	91	[SEQ ID NO:15]														
Pol Trans Pre 94	IRIPRTPXRVVTGGVFLVDK	KNPHNTAESRLVVD	FSQF	SRGNTRYRVSWPKFA	VNTLQSL	TNLLSSNL	SWLSLDV	SAAFYHLPLHPAAMP	PHLLAVGSSGLSR	YVA 193													
Pol Trans 2 97	IRIPRTPSKRVGTGGVFLVDK	KNPHNTAESRLVVD	FSQF	SRGNTRYRVSWPKFA	VNTLQSL	TNLLSSNL	SWLSLDV	SAAFYHLPLHPAAMP	PHLLAVGSSGLSR	YVA 196													
Pol Trans 3 101	IRIPRTPSKRVGTGGVFLVDK	KNPHNTAESRLVVD	FSQF	SRGNTRYRVSWPKFA	VNTLQSL	TNLLSSNL	SWLSLDV	SAAFYHLPLHPAAMP	PHLLAVGSSGLSR	YVA 200													
Pol Trans of 4 92	IRIPRTPARVVTGGVFLVDK	KNPHNTAESRLVVD	FSQF	SRGNTRYRVSWPKFA	VNTLQSL	TNLLSSNL	SWLSLDV	SAAFYHLPLHPAAMP	PHLLAVGSSGLSR	YVA 191													
Pol Trans Pre 194	RLSNSRLIFNHQBGXNQNL	HDYCSRNLYVSL	LLLYQTFGR	KLHLYSHPI	ILGFRKI	PMGVGLSP	PFLLAQ	FTSAICSVVRR	APPHCLAFSYMD	DVVLGAKS 293													
Pol Trans 2 197	RLSNSRLIANNQHGTM	PDLDYCSRNLYVSL	LLLYQTFGR	KLHLYSHPI	ILGFRKI	PMGVGLSP	PFLLAQ	FTSAICSVVRR	APPHCLAFSYMD	DVVLGAKS 296													
Pol Trans 3 201	RLSNSRLIANNQHGTM	PDLDYCSRNLYVSL	LLLYQTFGR	KLHLYSHPI	ILGFRKI	PMGVGLSP	PFLLAQ	FTSAICSVVRR	APPHCLAFSYMD	DVVLGAKS 300													
Pol Trans 4 192	RLSNSRLIFNHQBGXNQNL	HDYCSRNLYVSL	LLLYQTFGR	KLHLYSHPI	ILGFRKI	PMGVGLSP	PFLLAQ	FTSAICSVVRR	APPHCLAFSYMD	DVVLGAKS 291													
Pol Trans Pre 294	VXHLES	LF	TA	VN	TF	LL	SL	GI	HL	NP	NK	TR	MG	YS	L	HP	MG	Y	V	I	G	C	336
Pol Trans 2 297	VQHLES	LF	TA	VN	TF	LL	SL	GI	HL	NP	NK	TR	MG	YS	L	HP	MG	Y	V	I	G	C	340
Pol Trans 3 301	VQHLES	LF	TA	VN	TF	LL	SL	GI	HL	NP	NK	TR	MG	YS	L	HP	MG	Y	V	I	G	C	344
Pol Trans 4 292	VQHLES	LF	TA	VN	TF	LL	SL	GI	HL	NP	NK	TR	MG	YS	L	HP	MG	Y	V	I	G	C	336

Figure 5

1	Pre	HBAa9	Trans of	1	PPASTNRQSGRQPTPLSPPLRNTHPQAMQWNSTTFHQTLQDPRVRGLYFPAGGSSGTVNPTVTTASPLSSIFRIGDPALP	17	MEMITSGFLGFLVLVQA	[SEQ ID NO:16]
2	1	HBAa9	Trans of	2	PPASTNRQSGRQPTPLSPPLRNTHPQAMQWNSTTFHQTLQDPRVRGLYFPAGGSSGTVNPTVTTASPLSSIFRIGDPALP	100	MEMITSGFLGFLVLVQA	[SEQ ID NO:17]
3	1	HBAa9	Trans of	3	PPASTNRQSGRQPTPLSPPLRNTHPQAMQWNSTTFHQTLQDPRVRGLYFPAGGSSGTVNPTVTTASPLSSIFRIGDPALP	100	MEMITSGFLGFLVLVQA	[SEQ ID NO:18]
4	1	HBAa9	Trans of	4	PPASTNRQSGRQPTPLSPPLRNTHPQAMQWNSTTFHQTLQDPRVRGLYFPAGGSSGTVNPTVTTASPLSSIFRIGDPALP	100	MEMITSGFLGFLVLVQA	[SEQ ID NO:19]
18	Pre	HBAa9	Trans of	18	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	117	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	117
2	101	HBAa9	Trans of	2	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	200	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	200
3	101	HBAa9	Trans of	3	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	200	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	200
4	101	HBAa9	Trans of	4	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	200	GFPELLTRILLTPQSLDSMWTSLNFIAGTTCVLCQNSQPTSNHSPSCPTCPGYRWMCLRRRIIFLPIILMCLLFLVLLVQYQGM	200
118	Pre	HBAa9	Trans of	118	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	217	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	217
2	201	HBAa9	Trans of	2	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	300	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	300
3	201	HBAa9	Trans of	3	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	300	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	300
4	201	HBAa9	Trans of	4	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	300	AGACRGTCTTTAAGTSMYPSCCCTKPSDGNCTCIPIPSSWAFGKFLWNASARFSLSLVFPVQWFGVGLSPTVWLSVTWMMWYWG	300
218	Pre	HBAa9	Trans of	218	IFFCLMWYI 226		IFFCLMWYI 226	
2	301	HBAa9	Trans of	2	IFFCLMWYI 309		IFFCLMWYI 309	
3	301	HBAa9	Trans of	3	IFFCLMWYI 309		IFFCLMWYI 309	
4	301	HBAa9	Trans of	4	IFFCLMWYI 309		IFFCLMWYI 309	

## Figure 6

8/32

	10	20	30	40	50	
S0						[SEQ ID NO:20]
S6						[SEQ ID NO:21]
S8					T	[SEQ ID NO:22]
S12	TTTTGGGGAGCCCTCAGGCTCAGGGCATATTACAACTCTGCCAGCAAAT					[SEQ ID NO:23]
S15				TACAACTTTGCCAGCAAAT		[SEQ ID NO:24]

	60	70	80	90	100	
S0						
S6						
S8	GCCCCCTTCTGCCTCCACCAATCGCCAGTCAGGAAGGCAGCCTACCCCGCT					
S12	CCACCTCCTGCCTCCACCAATCGCCAGTCAGGAAGGCAGCCTACCCCGCT					
S15	CCACCTCCTGCCTCCACCAATCGCCAGTCAGGAAGGCAGCCTACCCCGCT					

	110	120	130	140	150	
S0						
S6						
S8	GTCTCCACCTTTGAGAGACACTCATCCTCAGGCCATGCAGTGGAACCTCAA					
S12	GTCTCCACCTTTGAGAGACACTCATCCTCAGGCCATGCAGTGGAACCTCAA					
S15	GTCTCCACCTTTGAGAGACACTCATCCTCAGGCCATGCAGTGGAACCTCAA					

	160	170	180	190	200	
S0						
S6						
S8	CAACCTTCCACCAAACCTCTGCAAGATCCCAGAGTGAAAGGCCTGTATTTTC					
S12	CAACCTTCCACCAAACCTCTGCAAGATCCCAGAGTGAAAGGCCTGTATTTTC					
S15	CAACCTTCCACCAAACCTCTGCAAGATCCCAGAGTGAAAGGCCTGTATTTTC					

	210	220	230	240	250	
S0						
S6						
S8	CCTGCTGGTGGCTCCAGTTCAGGAACAGTAAACCCTGTTCCGACTACTGC					
S12	CCTGCTGGTGGCTCCAGTTCAGGAACAGTAAACCCTGTTCCGACTACTGC					
S15	CCTGCTGGTGGCTCCAGTTCAGGAACAGTAAACCCTGTTCCGACTACTGC					

	260	270	280	290	300	
S0						
S6						
S8	CTCTCACTCATCGTCAATCTTCTCGAGGATTGGGGTCCCTGCGCTGAACA					
S12	CTCTCACTCATCGTCAATCTTCTCGAGGATTGGGGTCCCTGCGCTGAACA					
S15	CTCTCACTCATCGTCAATCTTCTCGAGGATTGGGGTCCCTGCGCTGAACA					

Figure 7

9/32

	310	320	330	340	350
S0					
S6					
S8	TGGAGAACATCACATCAGGACTCCTAGGACCCCTTCTCGTGTTACAGGCG				
S12	TGGAGAACATCACATCAGGACTCCTAGGACCCCTTCTCGTGTTACAGGCG				
S15	TGGAGAACATCACATCAGGACTCCTAGGACCCCTTCTCGTGTTACAGGCG				

	360	370	380	390	400
S0				CGCAGAGTCTAGACTC	
S6					
S8	GGGTTTTTCTTGTTGACAAGAATCCTCACAATACCGCAGAGTCTAGACTC				
S12	GGGTTTTTCTTGTTGACAAGAATCCTCACAATACCGCAGAGTCTAGACTC				
S15	GGGTTTTTNTTGTTGACAAGAATCCTCACAATACCGCAGAGTCTAGACTC				

	410	420	430	440	450
S0	GTGGTGGACTTCTCTCAATTTTCGAGGGGGGACTACCGTGTGTCTTGGCC				
S6					
S8	GTGGTGGACTTCTCTCAATTTTCGAGGGGGGACTACCGTGTGTCTTGGCC				
S12	GTGGTGGACTTCTCTCAATTTTCGAGGGGGGACTACCGTGTGTCTTGGCC				
S15	GTGGTGGACTTCTCTCAATTTTCGAGGGGGGACTACCGTGTGTCTTGGCC				

	460	470	480	490	500
S0	AAAATTGCGAGTCCCCAACCTCCAATCACTCACCAACCTCCTGTCCTCCA				
S6		TTACTCACCNACCTCCTGTCCTCCA			
S8	AAAATTGCGAGTCCCCAACCTCCAATCACTCACCAACCTCCTGTCCTCCA				
S12	AAAATTGCGAGTCCCCAACCTCCAATCACTCACCAACCTCCTGTCCTCCA				
S15	AAAATTGCGAGTCCCCAACCTCCAATCACTCACCAACCTCCTGTCCTCCA				

	510	520	530	540	550
S0	ACTTGTCTGCTGCTATGCTGGATGTGTCTGCGGCGTTTTATCATCTTCCT				
S6	ACTTGTCTGCTGCTATGCTGGATGTGTCTGCGGCGTTTTATCATCTTCCT				
S8	ACTTGTCTGCTGCTATGCTGGATGTGTCTGCGGCGTTTTATCATCTTCCT				
S12	ACTTGTCTGCTGCTATGCTGGATGTGTCTGCGGCGTTTTATCATCTTCCT				
S15	ACTTGTCTGCTGCTATGCTGGATGTGTCTGCGGCGTTTTATCATCTTCCT				

	560	570	580	590	600
S0	CTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCTGGACTGTC				
S6	CTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCTGGACTGTC				
S8	CTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCTGGACTGTC				
S12	CTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCTGGACTGTC				
S15	CTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCTGGACTGTC				

	610	620	630	640	650
S0	AAGGTATGTTGCCCGTTTGTCTCTAATTCCAGGATCCTCAACCACCAGC				
S6	AAGGTATGTTGCCCGTTTGTCTCTAATTCCAGGATCCTCAACCACCAGC				
S8	AAGGTATGTTGCCCGTTTGTCTCTAATTCCAGGATCCTCAACCACCAGC				
S12	AAGGTATGTTGCCCGTTTGTCTCTAATTCCAGGATCCTCAACCACCAGC				
S15	AAGGTATGTTGCCCGTTTGTCTCTAATTCCAGGATCCTCAACCACCAGC				

Figure 7 (continued)

10/32

	660	670	680	690	700
S0	ACGGGACCATGCCGAACCTGCACGACTCCTGCTCAAGGAACCTCTACGGT				
S6	AGGGGACCATGCCGAACCTGCACGACTCCTGCTCAAGGAACCTCTACGGT				
S8	AGGGGACCATGCCGAACCTGCACGACTCCTGCTCAAGGAACCTCTACGGT				
S12	AGGGGACCATGCCGAACCTGCACGACTCCTGCTCAAGGAACCTCTACGGT				
S15	AGGGGACCATGCCGAACCTGCACGACTCCTGCTCAAGGAACCTCTACGGT				
	710	720	730	740	750
S0	TCCCTCATGTTGCTGTACCAAACCTTCGGACGGAAATTGCACCTGTATTTC				
S6	TCCCTCATGTTGCTGTACCAAACCTTCGGACGGAAATTGCACCTGTATTTC				
S8	TCCCTCATGTTGCTGTACCAAACCTTCGGACGGAAATTGCACCTGTATTTC				
S12	TCCCTCATGTTGCTGTACCAAACCTTCGGACGGAAATTGCACCTGTATTTC				
S15	TCCCTCATGTTGCTGTACCAAACCTTCGGACGGAAATTGCACCTGTATTTC				
	760	770	780	790	800
S0	CCATCCCATCATCCTGGGCTTTTCGAAAATTCTATGGGAGTGGGCCTCA				
S6	CCATCCCATCATCCTGGGCTTTTCGAAAATTCTATGGGAGTGGGCCTCA				
S8	CCATCCCATCATCCTGGGCTTTTCGAAAATTCTATGGGAGTGGGCCTCA				
S12	CCATCCCATCATCCTGGGCTTTTCGAAAATTCTATGGGAGTGGGCCTCA				
S15	CCATCCCATCATCCTGGGCTTTTCGAAAATTCTATGGGAGTGGGCCTCA				
	810	820	830	840	850
S0	GCCCGTTTCTCCTGGCTCAGTTTACTAGTGCCATTTGTTTCAGTGGTTCGT				
S6	GCCCGTTTCTCCTGGCTCAGTTTACTAGTGCCATTTGTTTCAGTGGTTCGT				
S8	GCCCGTTTCTCCTGGCTCAGTTTACTAGTGCCATTTGTTTCAGTGGTTCGT				
S12	GCCCGTTTCTCCTGGCTCAGTTTACTAGTGCCATTTGTTTCAGTGGTTCGT				
S15	GCCCGTTTCTCCTGGCTCAGTTTACTAGTGCCATTTGTTTCAGTGGTTCGT				
	860	870	880	890	900
S0	AGGGCTTTCCCCCACTGTCTGGCTTTTAGTTATATGGATGATGTGGTATT				
S6	AGGGCTTTCCCCCACTGTCTGGCTTTTAGTTATATGGATGATGTGGTATT				
S8	AGGGCTTTCCCCCACTGTCTGGCTTTTAGTTATATGGATGATGTGGTATT				
S12	AGGGCTTTCCCCCACTGTCTGGCTTTTAGTTATATGGATGATGTGGTATT				
S15	AGGGCTTTCCCCCACTGTCTGGCTTTTAGTTATATGGATGATGTGGTATT				
	910	920	930	940	950
S0	GGGGGCCAAGTCTGTATCGCATCTTGAGTCCCTTTTACCGCTGTTACCA				
S6	GGGGGCCAAGTCTGTATCGCATCTTGAGTCCCTTTTACCGCTGTTACCA				
S8	GGGGGCCAAGTCTGTATCGCATCTTGAGTCCCTTTTACCGCTGTTACCA				
S12	GGGGGCCAAGTCTGTATCGCATCTTGAGTCCCTTTTACCGCTGTTACCA				
S15	GGGGGCCAAGTCTGTATCGCATCTTGAGTCCCTTTTACCGCTGTTACCA				
	960	970	980	990	1000
S0	ATTTTCCTTTGTCTTTGGGTATACATTTAAACCCTAACAAAACAAAAGA				
S6	ATTTTCCTTTGTCTTTGGGTATACATTTAAATCCTAACAAAACAAAAGA				
S8	ATTTTCCTTTGTCTTTGGGTATACATTTAAATCCTAACAAAACAAAAGA				
S12	ATTTTCCTTTGTCTTTGGGTATACATTTAAATCCTAACAAAACAAAAGA				
S15	ATTTTCCTTTGTCTTTGGGTATACATTTAAATCCTAACAAAACAAAAGA				
	1010	1020	1030	1040	1050

Figure 7 (continued)



11/32

S0 TGGGGTTACTCCCTACATTTTATGGGCTATGTCATTGGAT  
S6 TGGGGTTACTCCCTACATTTTATGGGCTATGTCATTGGATGTCATGGGTC  
S8 TGGGGTTACTCCCTACATTTTATGGGCTATGTCATTGGATGTCATGGGTC  
S12 TGGGGTTACTCCCTACATTTTATGGGCTATGTCATTGGATGTCATGGGTC  
S15 TGGGGTTACTCCCTACA

	1060	1070	1080	1090	1100
S0					
S6	CTTGCCACAAGAACACATCAGACAAAAAATCAAAGAATGTTTTAGAAAAC				
S8	CTTGCCACAAGAACACATCAGACAAAAAATCA				
S12	CTTGCCACAAGAACACATCAGACAAAAAATCAAAGAATGTTTTAGAAAAC				
S15					

**Figure 7 (continued)**

12/32

Patient B Am

	260	270	280	290	300	
S0	SGHTTNFASKSTSCLHQSPVRKAAYPAVSTFERHSSSGHAVELNNLPPNS					[SEQ ID NO:25]
S6						[SEQ ID NO:26]
S8	CPFCLHQSPVRKAAYPAVSTFERHSSSGHAVELNNLPPNS					[SEQ ID NO:27]
S12	SGHITNSASKSTSCLHQSPVRKAAYPAVSTFERHSSSGHAVELNNLPPNS					[SEQ ID NO:28]
S15	TNFASKSTSCLHQSPVRKAAYPAVSTFERHSSSGHAVELNNLPPNS					[SEQ ID NO:29]
	310	320	330	340	350	
S0	ARSQSERPVFPCWWLQFRNSKPCSDYCLSLIVNLLLEDWGPCAEGHEHHIR					
S6						
S8	ARSQSERPVFPCWWLQFRNSKPCSDYCLSLIVNLLLEDWGPCAEGHEHHIR					
S12	ARSQSERPVFPCWWLQFRNSKPCSDYCLSLIVNLLLEDWGPCAEGHEHHIR					
S15	ARSQSERPVFPCWWLQFRNSKPCSDYCLSLIVNLLLEDWGPCAEGHEHHIR					
	360	370	380	390	400	
S0	TPRTPSRVTGGVFLVDKNPHNTAESRLVVDQFSQFSGDYRVSWPKFAVPN					
S6						
S8	TPRTPSRVTGGVFLVDKNPHNTAESRLVVDQFSQFSGDYRVSWPKFAVPN					
S12	TPRTPSRVTGGVFLVDKNPHNTAESRLVVDQFSQFSGDYRVSWPKFAVPN					
S15	TPRTPSRVTGGVFXVDKNPHNTAESRLVVDQFSQFSGDYRVSWPKFAVPN					
	410	420	430	440	450	
S0	LQSLTNLLSSNLSWLSLDVSAAFYHLPLHPAAMPHLLVGSSGLSRYVARL					
S6	SNLSWLSLDVSAAFYHLPLHPAAMPHLLVGSSGLSRYVARL					
S8	LQSLTNLLSSNLSWLSLDVSAAFYHLPLHPAAMPHLLVGSSGLSRYVARL					
S12	LQSLTNLLSSNLSWLSLDVSAAFYHLPLHPAAMPHLLVGSSGLSRYVARL					
S15	LQSLTNLLSSNLSWLSLDVSAAFYHLPLHPAAMPHLLVGSTGLSRYVARL					
	460	470	480	490	500	
S0	SSNSRIILNHQGTMPNLHDSCSRNLYGSLMMLLYQTFGRKLHLYSHPIILG					
S6	SSNSRIILNHQGTMPNLHDSCSRNLYGSLMMLLYQTFGRKLHLYSHPIILG					
S8	SSNSRIILNHQGTMPNLHDSCSRNLYGSLMMLLYQTFGRKLHLYSHPIILG					
S12	SSNSRIILNHQGTMPNLHDSCSRNLYGSLMMLLYQTFGRKLHLYSHPIILG					
S15	SSNSRIILNHQGTMPNLHDSCSRNLYGSLMMLLYQTFGRKLHLYSHPIILG					
	510	520	530	540	550	
S0	FRKIPMGVGLSPFLMAQFTSAICSVVRRAPPHCLAFGYVDDVVLGAKSVS					
S6	FRKIPMGVGLSPFLMAQFTSAICSVVRRAPPHCLAFGYVDDVVLGAKSVS					
S8	FRKIPMGVGLSPFLMAQFTSAICSVVRRAPPHCLAFGYVDDVVLGAKSVS					
S12	FRKIPMGVGLSPFLMAQFTSAICSVVRRAPPHCLAFGYVDDVVLGAKSVS					
S15	FRKIPMGVGLSPFLMAQFTSAICSVVRRAPPHCLAFGYVDDVVLGAKSVS					
	560	570	580	590	600	
S0	HLESLFTAXTNFLLSLGIHLNPNKTKRWGYSLHFMGYVIGCHGSLPQEH					
S6	HLESLFTAVTNFLLSLGIHLNPNKTKRWGYSLHFMGYVIG					
S8	HLESLFTAXTNFLLSLGIHLNPNKTKRWGYSLHFMGYVIGCHGSLPQEH					
S12	HLESLFTAVTNFLLSLGIHLNPNKTKRWGYSLHFMGYVIG					
S15	HLESLFTAVTNFLLSLGXHLNPNKTKRWGYS					

Figure 8

13/32

	10	20	30	40	50	
S0						[SEQ ID NO:30]
S6						[SEQ ID NO:31]
S8						[SEQ ID NO:32]
S12	LGSPQAQGILQTL	PANPPPASTNRQSGRQPTPLSPPLRDTHPQAMQWNST				[SEQ ID NO:33]
S15		PPPASTNRQSGRQPTPLSPPLRDTHPQAMQWNST				[SEQ ID NO:34]

	60	70	80	90	100	
S0						
S6						
S8						
S12	TFHQTLQDPRVKGLYFPAGGSSSGTVNPVPTTASHSSSIFSRIGVPALNM					
S15	TFHQTLQDPRVKGLYFPAGGSSSGTVNPVPTTASHSSSIFSRIGVPALNM					

	110	120	130	140	150	
S0			QSLDSWWTSLNFRGGTTVCLGQ			
S6						
S8						
S12	ENITSGLLGPLLVLQAGFFLLTRILTI	QSLDSWWTSLNFRGGTTVCLGQ				
S15	ENITSGLLGPLLVLQAGFFLLTRILTI	QSLDSWWTSLNFRGGTTVCLGQ				

	160	170	180	190	200	
S0	NSQSPTS	NHSPTSCPPTCPGYRWMCLRRFII	FLFILLCLIFLLVLLDCQ			
S6		PPTCPGYRWMCLRRFII	FLFILLCLIFLLVLLDCQ			
S8		PTCPGYRWMCLRRFII	FLFILLCLIFLLVLLDCQ			
S12	NSQSPTS	NHSPTSCPPTCPGYRWMCLRRFII	FLFILLCLIFLLVLLDCQ			
S15	NSQSPTS	NHSPTSCPPTCPGYRWMCLRRFII	FLFILLCLIFLLVLLDCQ			

	210	220	230	240	250	
S0	GMLPVCPLIPGSSTTSRGPCRTCTTPAQGTSTVPSCCCTKPSDGNCTCIP					
S6	GMLPVCPLIPGSSTTSRGPCRTCTTPAQGTSTVPSCCCTKPSDGNCTCIP					
S8	GMLPVCPLIPGSSTTSRGPCRTCTTPAQGTSTVPSCCCTKPSDGNCTCIP					
S12	GMLPVCPLIPGSSTTSRGPCRTCTTPAQGTSTVPSCCCTKPSDGNCTCIP					
S15	GMLPVCPLIPGSSTTSRGPCRTCTTPAQGTSTVPSCCCTKPSDGNCTCIP					

	260	270	280	290	300	
S0	IPSSWAFGKFLWEWASARFSWLSLLVPFVQWVGLSPTVWLLVIMMMWYW					
S6	IPSSWAFGKFLWEWASARFSWLSLLVPFVQWVGLSPTVWLLVIMMMWYW					
S8	IPSSWAFGKFLWEWASARFSWLSLLVPFVQWVGLSPTVWLLVIMMMWYW					
S12	IPSSWAFGKFLWEWASARFSWLSLLVPFVQWVGLSPTVWLLVIMMMWYW					
S15	IPSSWAFGKFLWEWASARFSWLSLLVPFVQWVGLSPTVWLLVIMMMWYW					

	310	320	
S0	GPSLYRILSPFLPLXPIFFCLWVYI		
S6	GPSLYRILSPFLPLXPIFFCLWVYI		
S8	GPSLYRILSPFLPLXPIFFCLWVYI		
S12	GPSLYRILSPFLPLXPIFFCLWVYI		
S15	GPSLYRILSPFLPLXPIFFCLWVXI		

Figure 9

14/32

10 20 30 40 50 60 70 80 90 100  
TACTACAAACCTTGGCCAGCAAAATCCGGCCTCCTGCTCTTACCAATCGCAGTCAGGAAGGAGCCCTCTGACTCCACCTTTGAGAAACACTCATCC  
110 120 130 140 150 160 170 180 190 200  
TCAGGCCATGCAGTGAACCTCCACAAACTCCACCGAATCTACAGATCCCAAGATGAAAGGCTGTATCTCCCTGCTGGTGGCTCCAGTTCAGGAACA  
210 220 230 240 250 260 270 280 290 300  
GTAAACCCCTGTTCCGACTACTGTCTCTCACACATCGTCAATCTTATCGAGGATGGGGACCCCTGCACTGAACATGGAGAACATCACATCAGGATTCCTAG  
310 320 330 340 350 360 370 380 390 400  
GACCCCTGCTCGTGTACAGCGGGGTTTCTTGTGTGACAAGAAATCCTCACAATACCGCAGAGTCTAGACTCGTGTGGACTTCTCTCAATTTTCTTAGG  
410 420 430 440 450 460 470 480 490 500  
GGGGACCAACCGTGTCCCTTGGCCAAAATTCGCAGTCCCAACCTCCAATCACTCACCAACCTCCTGTCTCCTCCAACCTTGTCTCTGGTTATCGCTGGATGTGT  
510 520 530 540 550 560 570 580 590 600  
CTGCGGCGTTTATCATATTCTTCATCTCTGCTGTATGCTCATCTTCTTGTGGTTCTTCTGGACTATCAAGGTATGTTGCCGTTTGGCCCTCTAA  
610 620 630 640 650 660 670 680 690 700  
TTCCAGGATCCTCAACCACCAGCACGGGACCATGCAGAACCTTGCTCAGGAACCTCTGTGTATCCCTCATGTTGCTGTACCAAAACCTWC  
710 720 730 740 750 760 770 780 790 800  
GGMCGSAAATTGCACCTGTATTCCCATCCCATCATCTCTGGGCTTCGGAAATTCCTATGGGAGTGGGCCCTCAGCCCGTTTCTCTGACTCAGTTTACTA  
810 820 830 840 850 860 870 880 890 900  
GTGCCAATTTGTTTCAGTGGTTCTGAGGCTTTCCCCACATGTTTGGCTTTTCAGTTATATGGATGATGTGTGTTATGGGGGCCAGGTCTGTACAGCATCGTGA  
910 920 930 940 950 960 970 980 990 1000  
GGCCCTTTTACCGCTGTACCAATTTTCTTTTGTCTCTGGGTATACATTTTAACCCCGGACAAACAAAGATGGGGTTACTCTTTTACATTTTCATGGGC  
1010 1020 1030  
TATGTATGGATGTTATGGGTCAATGGCCAC

Figure 10

15/32

10 20 30 40 50 60 70 80 90 100  
TTNLASKSASCLYQSPVRKAAYPDSSTFEKHSSGHAVELHKLPPNSTRSQSERPVSPCWNLQFRNSKPCSDYCLSHIVNLIEDWGPCTEGEHHRIPR  
110 120 130 140 150 160 170 180 190 200  
TPARVTGGVFLVDKNPHTAESRLVVDPSQFSRGRHVPWPKFAVPNLQSLTNLLSSNLSWLSLDVSAAFYHIPLHPAAMPHELLVGSGLSRYVARLP  
210 220 230 240 250 260 270 280 290 300  
SRILNHQGTQNLHEDSCSRNLY/FVSIIMLLYQTF/TGRKLHLYSHPIILGFRKIPMGVGLSPFLLTQFTSAICSVVRRAPPHCLAFSYMDVVLGARSVQ  
310 320 330 340  
HREALFTAVTNFLLSLGIHLTPDKTKRWGYSLHFMGYVIGCYGSLP

Figure 11

16/32

10 20 30 40 50 60 70 80 90 100  
LQTLPANPPASTNRQSGRQPTPLTPPLENTHPQAMQWNSTNHFRTLQDPRVKGLYLPAGGSSSGTVNPVPTTVSHTSSILSRIGDPALNMENTSGFLG

110 120 130 140 150 160 170 180 190 200  
PLLVLQAGFFLLTRILTIPQSLDSWWTSLNFLGGTTVCLGQNSQSPTSNHSPSCPPTCPGYRWMCLRRFIIIFLILLCLIFLLVLLDYQGMFLPVCPLI

210 220 230 240 250 260 270 280 290 300  
PGSSTTSTGPCRCTCTTPAQGTSM/LYPSCCCTKPS/TAANCTCIPSPSSWAFGKFLMEWASARFS\*LSILLVPFVQWFGVGLSPTVWLSVIWMMWYWGPGLYS

310  
IVRPFLPLLPIFFCLWVYI

Figure 12

**17/32**

10	20	30	40	50	60	70	80	90	100
TGGTCA	CAGTGC	CAACAG	TTCTCT	CTCTG	CTCCAC	CAATCG	GAGG	CAGCCT	CTTA
110	120	130	140	150	160	170	180	190	200
GGCCAT	GGTGG	CTCAG	CCCTG	CTGCT	CCAG	TTCAG	GAAC	TCAAC	CTGTT
210	220	230	240	250	260	270	280	290	300
GACCC	TGCGC	CAACAT	GGAAC	ATCCT	AGGAC	CCCTG	CTGCT	TACAG	GCGGG
310	320	330	340	350	360	370	380	390	400
CGCAG	AGCTA	GACTCT	CTCAG	TTTCT	AGGGG	ATCAC	CCGTTG	TCTT	GCGGG
410	420	430	440	450	460	470	480	490	500
AAACC	TCTGT	CTCCA	TTTGAC	CTGGT	ATTCG	TGATAT	CTCG	TGCTG	TGCTG
510	520	530	540	550	560	570	580	590	600
GTTC	TTCG	ATAT	CAAG	TATG	TTGCC	CGTTT	GTCTC	TATCC	AGAT
610	620	630	640	650	660	670	680	690	700
AAAG	CACTC	TATG	TTTCC	CTCAT	GTGTG	CTGTAC	AAAA	CTTAG	CGAAT
710	720	730	740	750	760	770	780	790	800
ATGG	AGTGG	CGCTC	AGTCC	GTTC	CTG	GTCTC	AGTTT	ACTAG	TGCCA
810	820	830	840	850	860	870	880		
TTGAT	GATG	TGGT	ACTG	GGGG	CCAG	TCTG	CAACA	CACTT	GTAG

## Figure 13

10 20 30 40 50 60 70 80 90 100  
GHSANSSCLHQS AVREAAAYSHLSTSKRQSSSGHGGGSACWVLFQFNTQPCSQYCLSHLVNLLLEDWGPCAHEGHHIRIPRTPARVTGGVFLVDKNPHNT  
110 120 130 140 150 160 170 180 190 200  
AESRLVDFSQFSRGITRVSWPKFAVPNLQSLTNLLSSNLTWLSLDMSAAFYHPIPLHPAAMPHELLIGSSGLSRYVVARLSSNSRIHNNQCGTLQNLHDSCS  
210 220 230 240 250 260 270 280 290  
RQLYVSLMLLYKTYGWLHLYSHPIILGFRKIPMGVGLSPFLLAQFTSAICSVIRRAPPHCLAFSYIDDVVLGAKSAQHLESITYTAVTNFLLSLG

Figure 14



19/32

10 20 30 40 50 60 70 80 90 100  
VTVPVPPASTNRQSGRQPTPISPPLRDSDHPQAMVAQPAAGSSSGTLPVPNIASHISSISLRTGDPAPNMENITSGFLGELLVLQAGFFLLTRILLTIP  
110 120 130 140 150 160 170 180 190 200  
QSLDSWWTSLFLGSPVCLGQNSQSPTSNHSPTSCPPI\*PGYRWICLRRFIIFFILPLCLIFLLVLDDYQGMLPVCPLIPGSGTTTSAGPCKTCTTPAQ  
210 220 230 240 250 260 270 280 290  
GNSMFPSCCCTKPTDGNCTCIPIPSSWAFKYLWENASVRPSWLSLLVPFVQ\*FVGLSPTVWLSAILNMWYWGPSLHNILSPFIPLLPIFFCLWV

Figure 15

20/32

10 20 30 40 50  
TCCTGTCCTCCAATTTGTCCTGGTTATCGCTGGATGTGTCTGCGGCGTTT

60 70 80 90 100  
TATGATATTCCCTCTTCATCCTGCTGCTATGCCTCATCTTCTTATTGGTTC

110 120 130 140 150  
TTCTGGATTATCAAGGTATGTTGCCCGTCTGTCTCTAATTCCAGGATCA

160 170 180 190 200  
ACAACAACCAGTACGGGACCATGCAAAACCAAACCTGCACGACTCCTGC

210 220 230 240 250  
TCAAGGCAACTCTATGTTTCCCTCATGTTGCTGTACAAAACCTACGGATG

260 270 280 290 300  
GAAATTGCACCTGTATTCCCATCCCATCGTCCTGGGCTTTCGCAAAATTC

310 320 330 340 350  
CTATGGGAGTGGGCCTCAGTCCGTTTCTCTTGGCTCAGTTTACTAGTGCC

360 370 380 390 400  
ATTTGTTCAAGTGGTTCGTAGGGCTTCCCCACTGTTTGGCTTTCAGCTA

410 420 430 440 450  
TATGGATGATGTGGTATTGGGGGCCAAGTCTGTACAGCATCGTGAGGCCC

460 470 480 490 500  
TTTATACAGCTGTTACCAATTTTCTTTTGTCTCTGGGTATACATTTAAAC

510 520 530 540 550  
CCTAACAAAACAAAAGATGGGGTTATTCCCTAAACTTCATGGGTACAT

560 570 580 590  
AATTGGAAGTTGGGGAACATTGCCACAGGATCATATTGTAC

Figure 16

**21/32**

10 20 30 40 50  
SNLSWLSLDVSAAFYDIPLHPAAMPHELLIGSSGLSRYVARLSSNSRINNN  
60 70 80 90 100  
QYGTMQNQNLHDSCSRQLYVSLMLLYKTYGWKLHLYSHPIVLGFRKIPMG  
110 120 130 140 150  
VGLSPFLLAQFTSAICSVVRRAFPCLAFSYMDDVVLGAKSVQHREALYT  
160 170 180  
AVTNFLLSLGIHLNPNKTKRWGYSLNFMGYIIGSWG


**Figure 17**

**22/32**

10 20 30 40 50  
SCPPICPGYRWMCLRRFMIFLLCLIFLLVLLDYQGMLPVCPLIPGS  
60 70 80 90 100  
TTTSTGPCKTKTCTTPAQGNSMFPSCCCTKPTDGNCTCIPSSWAFKRF  
110 120 130 140 150  
LWEWASVRFSWLSLLVPFVQWFVGLSPTVWLSAIWMMWYWGPSLYSIVRP  
160  
FIQLLPFIFFCLWVYI

**Figure 18**

23/32

**Figure 19**

24/32

```
      10      20      30      40      50
AATCCTCACAATACCGCAGAGTCTAGACTTCGTGGTGA CTCTCTCAATT

      60      70      80      90     100
TTCTAGGGGACCACCCGTGTGTCTTGGCCAAAATTCGCAGTCCCCAACCT

     110     120     130     140     150
CCAATCACTCACCAACCTCTTGTCTCCAATTTGTCCTGGTTATCGCTGG

     160     170     180     190     200
ATGTGTCTGCGGCGTTTTATCATATCCCTCTTCATCCTGCTGCTATGCCT

     210     220     230     240     250
CATCTTCTTATTGGTTCTTCTGGATTATCAAGGTATGTTGCCCGTTTGTC

     260     270     280     290     300
CTCTAATTCAGGATCCACAACAACAGTACGGGACCCTGCAAAACCTGC

     310     320     330     340     350
ACGACTCCTGCTCAAGGCAACTCTATGTTTCCCTCATGTTGCTGTACAAA

     360     370     380     390     400
ACCTACGGATGGAAATTGCACMTGTATTCCCATCCCATCATCTTGGGCTT

     410     420     430     440     450
TCGCAAAATACCTATGGGAGTGGGCCTCAGTCCGTTTCTCTTGGTTCAGT

     460     470     480     490     500
TTACTAGTGCCATTTGTTTCAGTGGTTCGTAGGGCTTTCCCCCACTGTTTG

     510     520     530     540     550
GCTTTCAGCTATATGGATGATATTGTACTGGGGGCCAAGTCTGTACAACA

     560     570     580     590     600
TCTTGAGTCCCTTTATACCGCTGTTACCAATTTTCTTTGTCTTTGGGTA

     610     620     630     640     650
TACATTTAACCCCTAACAAAACAAAGAGATGGGGTTATTCCCTGAATTTT

     660
ATGGGTTATGTAATTGGAA
```

Figure 20

25/32

10 20 30 40 50  
SNLSWLSLDVSAAFYHIPLHPAAMPHLLIGSSGLSRYVARLSSNSRIHNN

60 70 80 90 100  
QYGTLONLHDSCSRQLYVSLMLLYKTYGWKLHXYSHPIILGFRKIPMGVG

110 120 130 140 150  
LSPFLLVQFTSAICSVVRRAPPHCLAFSYMDDIVLGAKSVQHLESlyTAV

160 170 180  
TNFLLSLGIHLTPNKTKRWGYSLNFMGYVIG

Figure 21

**26/32**

10 20 30 40 50  
PICPGYRWMCLRRFIISLFILLCLIFLLVLLDYQGMLPVCPLIPGSTTT  
60 70 80 90 100  
STGPCKTCTTPAQGNSMFPSCCCTKPTDGNCTCIPIPSSWAFAYLWEWA  
110 120 130 140 150  
SVRFSWFSLLVPFVQWFVGLSPTVWLSAIWMILYWGPSLYNILSPFIPLL  
160  
PIFFCLWVYI

**Figure 22**



27/32

10 20 30 40 50  
TCCAATTTGTCCTGGGTATCGCTGGATGTGTCTGCGGCGTTTTATCATAT

60 70 80 90 100  
TCCTCTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCTGGAC

110 120 130 140 150  
TATCAAGGTATGTTGCCCGTTTGTCTCTACTTCCAGGAACATCAACTAC

160 170 180 190 200  
CAGCACGGGACCATGCAAGACCTGCACGACTCCTGCTCAAGGAACCTCTA

210 220 230 240 250  
TGTTTCCCTCTTGTTGCTGTACAAAACCTTCGGACGGAAATTGCACTTGT

260 270 280 290 300  
ATTCCCATCCCATCGTCTTGGGCTTTCGCAAGATTCCTATGGGAGTGGGC

310 320 330 340 350  
CTCAGTCCGTTTTCTCTTGGCTCARTTTACTAGTGCCATTTGTTCACTGGT

360 370 380 390 400  
TCGTAGGGCTTTCCCCACTGTTTGGCTTTCAGTTATATTGATGATGTGG

410 420 430 440 450  
TATTGGGGGCCAAGTCTGTACAACATCTTGAATCCCTTTTACCTCTATT

460 470 480 490 500  
ACCAATTTTCTTATGTCTTTGGGTATACATTTAAACCCTAAGAAAACCAA

510 520 530 540 550  
ACGTTGGGGCTACTCCCTTAACTTCATGGGATATGTAATTGGAAGTTGGG

GTAC

Figure 23

28/32

10 20 30 40 50  
SNLSWVSLDVSAAFYHIPLHPAAMPHELLVGSSGLSRYVARLSSTSRNINY

60 70 80 90 100  
QHGTMOQLHDSCSRNLYVSLLLLYKTFGRKLHLYSHPIVLGFRKIPMGVG

110 120 130 140 150  
LSPFLLAQFTSAICSVVRRAPHCIAFSYIDDVVLGAKSVQHLESLFTSI

160 170 180  
TNFLMSLGIHLNPKKTKRWGYSLNFMGYVIGSWG

Figure 24

29/32

10	20	30	40	50
PICPGYRWMCLRRFIIFLFILLLLCLIFLLVLLDYQGMLPVCPLLPGTSTT				
60	70	80	90	100
STGPCKTCTTPAQGTSMFPSCCCTKPSDGNCTCIPSSWAFARFLWEWA				
110	120	130	140	150
SVRFSWLXLLVPFVQWFVGLSPTVWLSVILMMWYWGPSLYNILNPFLPLL				
160				
PIFLCLWVYI				

Figure 25

30/32

```

      10      20      30      40      50
CAGCAAATCCGCCTCCTGCCTCTACCAATCGCCAGTCAGGAAGGCAGCCT
      60      70      80      90     100
ACCCCTCTGTCTCCACCTTTGRGAAACACTCATCCTCAGGCCATGCAGTG
      110     120     130     140     150
GAACTCCACAACCTTCCACCAAACCTCTGCWAGATCCCAGAGTGAGAGGCC
      160     170     180     190     200
TGTATTTCCCTGCTGGTGGCTCCAGTTCAGGAACAGTAAACCTGTTCCG
      210     220     230     240     250
ACTTCTGTCTCTCACACATCGTCAATCTTCTCGAGGATTGGGGWCCCTGC
      260     270     280     290     300
GCTGAACATGGAGAACATCACATCAGGATTCCCTAGGACCCCTGCTCGTGT
      310     320     330     340     350
TACAGGCGGGGTTTTTCTTGTTGACAAGAATCCTCACAATACCGCAGAGT
      360     370     380     390     400
CTAGACTCGTGGTGGACTTCTCTCAATTTCTAGGGGGAACCTACCGTGTG
      410     420     430     440     450
TCTTGGCCAAAATTTCGCAGTTCCCAACCTCCAATCACTACCAACCTCCT
      460     470     480     490     500
GTCCTCCAACCTTGWCTGGTTATCGCTGGATGTRTCTGCGGCGTTTTATC
      510     520     530     540     550
ATCTTCCTCTTCATCCTGCTGCTATGCCTCATCTTCTTGTTGGTTCTTCT
      560     570     580     590     600
GGACTATCAAGGTATGTTGCCCCTTTGTCCTCTARTTCCAGGATCTTCAA
      610     620     630     640     650
CCACCAGCACGGGACCATGCAGAACCTGCACGACTCCTGCTCAAGGAAMC
      660     670     680     690     700
TCTATGAATCCCTCCTGTTGCTGTACCAAACCTTCGGACGGAAATTGCAC
      710     720     730     740     750
CTGTATTCCCATCCCATCATCCTGGGCTTTCGGAAAATTCCTATGGGAGT
      760     770     780     790     800
GGGCCTCAGCCCGTTTCTCCTGRCTCAGTTTACTAGTGCCATTTGTTTCAG
      810     820     830     840     850
TGGTTCGTAGGGCTTTCCCCCACTGTTTGGCTTTCAGTTATATGGATGAT
      860     870     880     890     900
GTGGTATTGGGGGCAAGTCTGTAYMGCATCTTRAGTCCCTTTTTACCGC
      910     920     930     940     950
TGTTACCAATTTTCTTTTGTCTYTGGGTATACATTAAACCTMACAAAA
      960     970     980     990    1000
CAAAAAGATGGGGTTACTCTTTACATTTTCATGGGCTATGTCATTGGATGT
      1010    1020    1030    1040
TATGGGTCATTGCCACAAGATCACATCAGACAGAAAATCAAAGAA

```

Figure 26

**31/32**

10 20 30 40 50  
SKSASCLYQSPVRKAAYPSVSTFXKHSSSGHAVELHNLPPNSARSQSERP  
60 70 80 90 100  
VFPCWWLQFRNSKPCSDFLSHIVNLLLEDWGPCAEGEHHIRIPRTPARV  
110 120 130 140 150  
TGGVFLVDKNPHNTAESRLVVDIFSQFSRGNYRVSWPKFAVPNLQSLTNLL  
160 170 180 190 200  
SSNLXWLSLDVSAAFYHLPLHPAAMPHELLVGSSGLSRYVARLSSXSRIFN  
210 220 230 240 250  
HQHGTMQNLHDSCSRXLYESLLLLLYQTFGRKLHLYSHPIILGFRKIPMGV  
260 270 280 290 300  
GLSPFLLXQFTSAICSVVRRAPPHCLAFSYMDDVVLGAKSVXHLXSLFTA  
310 320 330 340  
VTNFLLSLGIHLNPKTKRWGYSLHFMGYVIGCYGSLPQDHIROKIKE

**Figure 27**

**32/32**

10 20 30 40 50  
ANPPPPASTNRQSGRQPTPLSPPLXNTHPQAMQWNSTTFHQTLXDPRVRGL

60 70 80 90 100  
YFPAGGSSSGTVNVPVPTSVSHTSSIFSRIGXPALNMENITSGFLGPLLLVL

110 120 130 140 150  
QAGFFLLTRILTIPQSLDSWWTSLNFLGGTTVCLGQNSQFPTSNHSPTSC

160 170 180 190 200  
PPTXPGYRWMXLRRFIIIFLLCLIFLLVLLDYQGMLPVCPLXPGSST

210 220 230 240 250  
TSTGPCRTCTTPAQGXSMNPSCCCTKPSDGNCTCIPSSWAFGKFLWEW

260 270 280 290 300  
ASARFSXLSLLVPFVQWVGLSPTVWLSVIWMMWYWGPSLYXILSPFLPL

310  
LPIFFCLWVYI

**Figure 28**

- 1 -

## SEQUENCE LISTING

<110> Melbourne Health (for all States except the US)  
Austin and Repatriation Medical Centre (for all States except the US)  
Southern Health (for all States except the US)  
Bartholomeusz, Angeline (US only)  
Locarnini, Stephen (US only)  
Ayres, Anna (US only)  
Colledge, Danielle (US only)  
Sasadeusz, Joseph (US only)  
Tillmann, Hans (US only)  
Angus, Peter (US only)  
Sievert, William (US only)

<120> Viral variants, detection and use

<130> 12175980/EJH

<140> International

<141> 2003-04-11

<150> AU PS1710

<151> 2002-04-12

<150> AU PS3224

<151> 2002-06-26

<160> 59

<170> PatentIn version 3.1

<210> 1

<211> 76

<212> PRT

<213> synthetic

<220>

<221> MISC\_FEATURE

<222> (2)..(2)

<223> X = L or R or I

<220>

<221> MISC\_FEATURE

<222> (3)..(3)

<223> X = E or D

<220>

<221> MISC\_FEATURE

<222> (9)..(9)

<223> X = T or D or A or N or Y

<220>

<221> MISC\_FEATURE

- 2 -

<222> (10)..(10)  
<223> X = E or D

<220>  
<221> MISC\_FEATURE  
<222> (13)..(13)  
<223> X = E or K or Q

<220>  
<221> MISC\_FEATURE  
<222> (15)..(15)  
<223> X = H or R or N

<220>  
<221> MISC\_FEATURE  
<222> (18)..(18)  
<223> X = I or T

<220>  
<221> MISC\_FEATURE  
<222> (23)..(23)  
<223> X = A or S

<220>  
<221> MISC\_FEATURE  
<222> (26)..(26)  
<223> X = T or R

<220>  
<221> MISC\_FEATURE  
<222> (40)..(40)  
<223> X = A or T or S

<220>  
<221> MISC\_FEATURE  
<222> (43)..(43)  
<223> X = R or T

<220>  
<221> MISC\_FEATURE  
<222> (45)..(45)  
<223> X = V or G

<220>  
<221> MISC\_FEATURE  
<222> (55)..(55)  
<223> X = S or I or T or N or V



- 3 -

<220>  
 <221> MISC\_FEATURE  
 <222> (56)..(56)  
 <223> X = T or S or H or Y

<220>  
 <221> MISC\_FEATURE  
 <222> (57)..(57)  
 <223> X = R or H or K or Q

<220>  
 <221> MISC\_FEATURE  
 <222> (69)..(69)  
 <223> X = Q or P

<220>  
 <221> MISC\_FEATURE  
 <222> (76)..(76)  
 <223> S is designated amino acid 74

<400> 1

Leu Xaa Xaa Asp Trp Gly Pro Cys Xaa Xaa His Gly Xaa His Xaa Ile  
 1 5 10 15

Arg Xaa Pro Arg Thr Pro Xaa Arg Val Xaa Gly Gly Val Phe Leu Val  
 20 25 30

Asp Lys Asn Pro His Asn Thr Xaa Glu Ser Xaa Leu Xaa Val Asp Phe  
 35 40 45

Ser Gln Phe Ser Arg Gly Xaa Xaa Xaa Val Ser Trp Pro Lys Phe Ala  
 50 55 60

Val Pro Asn Leu Xaa Ser Leu Thr Asn Leu Leu Ser  
 65 70 75

<210> 2  
 <211> 181  
 <212> PRT  
 <213> synthetic

<220>  
 <221> MISC\_FEATURE  
 <222> (1)..(1)  
 <223> S = amino acid 75

- 4 -

<220>  
<221> MISC\_FEATURE  
<222> (2) .. (2)  
<223> X = N or D

<220>  
<221> MISC\_FEATURE  
<222> (17) .. (17)  
<223> X = I or P

<220>  
<221> MISC\_FEATURE  
<222> (29) .. (29)  
<223> X = I or v

<220>  
<221> MISC\_FEATURE  
<222> (35) .. (35)  
<223> X = S or D

<220>  
<221> MISC\_FEATURE  
<222> (44) .. (44)  
<223> X = T or N

<220>  
<221> MISC\_FEATURE  
<222> (46) .. (46)  
<223> X = R or N

<220>  
<221> MISC\_FEATURE  
<222> (47) .. (47)  
<223> X = N or I

<220>  
<221> MISC\_FEATURE  
<222> (48) .. (48)  
<223> X = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (50) .. (50)  
<223> X = N or Y or H

<220>  
<221> MISC\_FEATURE

- 5 -

<222> (52) .. (52)  
<223> X = H or Y

<220>  
<221> MISC\_FEATURE  
<222> (53) .. (53)  
<223> X = G or R

<220>  
<221> MISC\_FEATURE  
<222> (54) .. (56)  
<223> X = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (57) .. (57)  
<223> X = D or N

<220>  
<221> MISC\_FEATURE  
<222> (60) .. (60)  
<223> X = D or N

<220>  
<221> MISC\_FEATURE  
<222> (61) .. (61)  
<223> X = S or Y

<220>  
<221> MISC\_FEATURE  
<222> (65) .. (65)  
<223> X = N or Q

<220>  
<221> MISC\_FEATURE  
<222> (71) .. (71)  
<223> X = L or M

<220>  
<221> MISC\_FEATURE  
<222> (75) .. (75)  
<223> X = K or Q

<220>  
<221> MISC\_FEATURE  
<222> (77) .. (77)  
<223> X = Y or F

- 6 -

<220>  
<221> MISC\_FEATURE  
<222> (79)..(79)  
<223> X = R or W

<220>  
<221> MISC\_FEATURE  
<222> (84)..(84)  
<223> X = Y or L

<220>  
<221> MISC\_FEATURE  
<222> (85)..(85)  
<223> X = S or A

<220>  
<221> MISC\_FEATURE  
<222> (89)..(89)  
<223> X = I or V

<220>  
<221> MISC\_FEATURE  
<222> (95)..(95)  
<223> X = I or L

<220>  
<221> MISC\_FEATURE  
<222> (99)..(99)  
<223> X = V or G

<220>  
<221> MISC\_FEATURE  
<222> (114)..(114)  
<223> X = C or L

<220>  
<221> MISC\_FEATURE  
<222> (115)..(115)  
<223> X = A or S

<220>  
<221> MISC\_FEATURE  
<222> (116)..(116)  
<223> X = V or M

<220>  
<221> MISC\_FEATURE

- 7 -

<222> (117)..(117)  
<223> X = V or T

<220>  
<221> MISC\_FEATURE  
<222> (118)..(118)  
<223> X = R or C

<220>  
<221> MISC\_FEATURE  
<222> (122)..(122)  
<223> X = F or P

<220>  
<221> MISC\_FEATURE  
<222> (125)..(125)  
<223> X = L or V

<220>  
<221> MISC\_FEATURE  
<222> (126)..(126)  
<223> X = A or V

<220>  
<221> MISC\_FEATURE  
<222> (128)..(128)  
<223> X = S or A

<220>  
<221> MISC\_FEATURE  
<222> (131)..(131)  
<223> M = amino acid 204

<220>  
<221> MISC\_FEATURE  
<222> (133)..(133)  
<223> X = V or L or M

<220>  
<221> MISC\_FEATURE  
<222> (138)..(138)  
<223> X = K or R

<220>  
<221> MISC\_FEATURE  
<222> (139)..(139)  
<223> X = S or T

- 8 -

<220>  
<221> MISC\_FEATURE  
<222> (140)..(140)  
<223> X = V or G

<220>  
<221> MISC\_FEATURE  
<222> (141)..(141)  
<223> X = Q or E

<220>  
<221> MISC\_FEATURE  
<222> (143)..(143)  
<223> X = L or S or R

<220>  
<221> MISC\_FEATURE  
<222> (145)..(145)  
<223> X = S or F

<220>  
<221> MISC\_FEATURE  
<222> (147)..(147)  
<223> X = F or Y

<220>  
<221> MISC\_FEATURE  
<222> (148)..(148)  
<223> X = T or A

<220>  
<221> MISC\_FEATURE  
<222> (149)..(149)  
<223> X = A or S

<220>  
<221> MISC\_FEATURE  
<222> (150)..(150)  
<223> X = V or I

<220>  
<221> MISC\_FEATURE  
<222> (151)..(151)  
<223> X = T or C

<220>  
<221> MISC\_FEATURE

- 9 -

&lt;222&gt; (152)..(152)

&lt;223&gt; X = N or S

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (153)..(153)

&lt;223&gt; X = F or V

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (156)..(156)

&lt;223&gt; X = S or D

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (157)..(157)

&lt;223&gt; X = L or V

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (164)..(164)

&lt;223&gt; X = N or Q

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (179)..(179)

&lt;223&gt; X = V or I

&lt;400&gt; 2

Ser	Xaa	Leu	Ser	Trp	Leu	Ser	Leu	Asp	Val	Ser	Ala	Ala	Phe	Tyr	His
1			5						10					15	

Xaa	Pro	Leu	His	Pro	Ala	Ala	Met	Pro	His	Leu	Leu	Xaa	Gly	Ser	Ser
		20						25					30		

Gly	Leu	Xaa	Arg	Tyr	Val	Ala	Arg	Leu	Ser	Ser	Xaa	Ser	Xaa	Xaa	Xaa
	35						40					45			

Asn	Xaa	Gln	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Leu	His	Xaa	Xaa	Cys	Ser	Arg
50							55				60				

Xaa	Leu	Tyr	Val	Ser	Leu	Xaa	Leu	Leu	Tyr	Xaa	Thr	Xaa	Gly	Xaa	Lys
65					70					75					80

Leu	His	Leu	Xaa	Xaa	His	Pro	Ile	Xaa	Leu	Gly	Phe	Arg	Lys	Xaa	Pro
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

- 10 -

	85		90		95
Met Gly Xaa Gly Leu Ser Pro Phe Leu Leu Ala Gln Phe Thr Ser Ala					
	100		105		110
Ile Xaa Xaa Xaa Xaa Xaa Arg Ala Phe Xaa His Cys Xaa Xaa Phe Xaa					
	115		120		125
Tyr Met Asp Asp Xaa Val Leu Gly Ala Xaa Xaa Xaa Xaa His Xaa Glu					
	130		135		140
Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Leu Xaa Xaa Gly Ile His					
	145		150		155
Leu Asn Pro Xaa Lys Thr Lys Arg Trp Gly Tyr Ser Leu Asn Phe Met					
	165		170		175
Gly Tyr Xaa Ile Gly					
	180				

<210> 3  
 <211> 23  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> OS1 primer

<400> 3  
 gcctcatttt gtgggtcacc ata

23

<210> 4  
 <211> 18  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> TTA3 primer

<400> 4  
 aaattcgcag tccccaaa

18

<210> 5  
 <211> 21  
 <212> DNA  
 <213> artificial sequence

<220>



- 11 -

&lt;223&gt; JM primer

&lt;400&gt; 5

ttgggggtgga gccctcaggc t

21

&lt;210&gt; 6

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; TTA4 primer

&lt;400&gt; 6

gaaaattggt aacagcgg

18

&lt;210&gt; 7

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; OS2 primer

&lt;400&gt; 7

tctctgacat actttccaat

20

&lt;210&gt; 8

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; sense primer

&lt;400&gt; 8

gcctcatttt gtgggtcacc ata

23

&lt;210&gt; 9

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; antisense primer

&lt;400&gt; 9

tctctgacat actttccaat

20

&lt;210&gt; 10

&lt;211&gt; 18

&lt;212&gt; DNA

- 12 -

<213> artificial sequence

<220>

<223> internal regions primer

<400> 10

tgcacgattc ctgctcaa

18

<210> 11

<211> 20

<212> DNA

<213> artificial sequence

<220>

<223> internal regions primer

<400> 11

tttctcaaag gtggagacag

20

<210> 12

<211> 18

<212> DNA

<213> artificial sequence

<220>

<223> PC1 forward primer

<400> 12

gggaggagat taggttaa

18

<210> 13

<211> 20

<212> DNA

<213> artificial sequence

<220>

<223> PC2 reverse primer

<400> 13

ggcaaaaacg agagtaactc

20

<210> 14

<211> 42

<212> DNA

<213> artificial sequence

<220>

<223> HBV-specific molecular beacon primer

<220>

<221> misc\_feature

<222> (42)..(42)

<223> n = L

- 13 -

<400> 14  
cgcgctcctac tgttcaagcc tccaagctgt gacgcgdabc yn 42

<210> 15  
<211> 280  
<212> DNA  
<213> artificial sequence

<220>  
<223> ILA 1 F, A-E

<220>  
<221> misc\_feature  
<222> (215)..(215)  
<223> n = any nucleotide

<400> 15  
tggctcagtt tactagtgcc atttggtcag tgggtcgtag ggctttcccc cactggttgg 60  
ctttcagtta tatggatgat gtggtattgg gggccaagtc tgtayagcay cttgagtccc 120  
tttttaccgc tgttaccaat tttcttttgt ctttgggtat acatttaaac cctaacaaaa 180  
ctaaaagatg gggttactct ttacatttca tgggntatgt cattggatgt tatgggtcat 240  
tgccacaaga tcacatcata cagaaaatca aagatggttt 280

<210> 16  
<211> 242  
<212> DNA  
<213> artificial sequence

<220>  
<223> ILA 2 F, A-E

<400> 16  
tggctcagtt tactagtgcc atttggtcag tgggtcgtag ggctttcccc cactggttgg 60  
ctttcagtta tatggatgat gtggtattgg gggccaagtc tgtacagcat cttgagtccc 120  
tttttaccgc tgttaccaat tttcttttgt ctttgggtat acatttaaac cctaacaaaa 180  
caaagagatg gggttactct ctaaatttta tgggttatgt cattggatgt tatgggtcct 240  
tg 242

<210> 17  
<211> 277  
<212> DNA  
<213> artificial sequence

- 14 -

&lt;220&gt;

&lt;223&gt; ILA 3 F, A-E

&lt;400&gt; 17

tggtcagtt tactagtgcc atttgttcag tggttcgtag ggctttcccc cactggttgg 60  
ctttcagtta tatggatgat gtggtattgg gggccaagtc tgtacagcat cttgagtc 120  
ttttaccgc tgttaccaat tttcttttgt ctttgggtat acatttaa 180  
caaagagatg ggggtactct cttaaatttta tgggttatgt cattggatgt tatgggtcct 240  
tgccacaaga acacatcata caaaaaatca aagaatg 277

&lt;210&gt; 18

&lt;211&gt; 237

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; ILA 4 F, A-E

&lt;400&gt; 18

tggtcagtt tactagtgcc atttgttcag tggttcgtag ggctttcccc cactggttgg 60  
ctttcagtta tatggatgat gtggtattgg gggccaagtc tgtacagcat cttgagtc 120  
ttttaccgc tgttaccaat tttcttttgt ctttgggcat acatttaa 180  
ctaaaagatg ggggtactct ttaaatttca tgggatatgt cattggatgg tatgggg 237

&lt;210&gt; 19

&lt;211&gt; 336

&lt;212&gt; PRT

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; Pol Trans Pre 1

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (11)..(11)

&lt;223&gt; x = any amino acid

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (17)..(17)

&lt;223&gt; x = any amino acid

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (38)..(38)

&lt;223&gt; x = any amino acid

- 15 -

<220>  
 <221> MISC\_FEATURE  
 <222> (46)..(46)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (50)..(50)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (101)..(101)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (208)..(208)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (295)..(295)  
 <223> x = any amino acid

<400> 19

Lys Leu Ala Ser Lys Ser Ala Ser Ser Ile Xaa Gln Ser Pro Val Arg  
 1 5 10 15

Xaa Ala Ala Tyr Pro Ala Val Ser Thr Phe Glu Lys His Ser Ser Ser  
 20 25 30

Gly His Ala Val Glu Xaa His Asn Leu Pro Pro Asn Ser Xaa Arg Ser  
 35 40 45

Gln Xaa Glu Arg Pro Val Phe Pro Cys Trp Trp Leu Gln Phe Arg Asn  
 50 55 60

Ser Lys Pro Cys Ser Asp Tyr Cys Leu Ser His Ile Val Asn Leu Leu  
 65 70 75 80

Glu Asp Trp Gly Pro Cys Ala Glu His Gly Glu His His Ile Arg Ile  
 85 90 95

- 16 -

Pro Arg Thr Pro Xaa Arg Val Thr Gly Gly Val Phe Leu Val Asp Lys  
 100 105 110

Asn Pro His Asn Thr Ala Glu Ser Arg Leu Val Val Asp Phe Ser Gln  
 115 120 125

Phe Ser Arg Gly Asn Tyr Arg Val Ser Trp Pro Lys Phe Ala Val Pro  
 130 135 140

Asn Leu Gln Ser Leu Thr Asn Leu Leu Ser Ser Asn Leu Ser Trp Leu  
 145 150 155 160

Ser Leu Asp Val Ser Ala Ala Phe Tyr His Leu Pro Leu His Pro Ala  
 165 170 175

Ala Met Pro His Leu Leu Val Gly Ser Ser Gly Leu Ser Arg Tyr Val  
 180 185 190

Ala Arg Leu Ser Ser Asn Ser Arg Ile Phe Asn His Gln Arg Gly Xaa  
 195 200 205

Met Gln Asn Leu His Asp Tyr Cys Ser Arg Asn Leu Tyr Val Ser Leu  
 210 215 220

Leu Leu Leu Tyr Gln Thr Phe Gly Arg Lys Leu His Leu Tyr Ser His  
 225 230 235 240

Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro Met Gly Val Gly Leu Ser  
 245 250 255

Pro Phe Leu Leu Ala Gln Phe Thr Ser Ala Ile Cys Ser Val Val Arg  
 260 265 270

Arg Ala Phe Pro His Cys Leu Ala Phe Ser Tyr Met Asp Asp Val Val  
 275 280 285

Leu Gly Ala Lys Ser Val Xaa His Leu Glu Ser Leu Phe Thr Ala Val  
 290 295 300

Thr Asn Phe Leu Leu Ser Leu Gly Ile His Leu Asn Pro Asn Lys Thr  
 305 310 315 320

Lys Arg Trp Gly Tyr Ser Leu His Phe Met Gly Tyr Val Ile Gly Cys  
325 330 335

<400> 20

Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His Leu Pro Leu  
165 170 175

- 18 -

His Pro Ala Ala Met Pro His Leu Leu Val Gly Ser Ser Gly Leu Ser  
 180 185 190

Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile Leu Asn Asn Gln  
 195 200 205

His Gly Thr Met Pro Asp Leu His Asp Tyr Cys Ser Arg Asn Leu Tyr  
 210 215 220

Val Ser Leu Leu Leu Leu Tyr Gln Thr Phe Gly Arg Lys Leu His Leu  
 225 230 235 240

Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro Met Gly Val  
 245 250 255

Gly Leu Ser Pro Phe Leu Leu Ala Gln Phe Thr Ser Ala Ile Cys Ser  
 260 265 270

Val Val Arg Arg Ala Phe Pro His Cys Leu Ala Phe Ser Tyr Met Asp  
 275 280 285

Asp Val Val Leu Gly Ala Lys Ser Val Gln His Leu Glu Ser Leu Phe  
 290 295 300

Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly Ile His Leu Asn Pro  
 305 310 315 320

Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu Asn Phe Met Gly Tyr Val  
 325 330 335

Ile Gly Cys Tyr  
 340

<210> 21  
 <211> 344  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> Pol Trans 3

<400> 21

Leu Ala Gln Gly Ile Leu Gln Asn Phe Ala Ser Lys Ser Ala Ser Cys  
 1 5 10 15



- 19 -

Leu His Gln Ser Pro Val Arg Lys Ala Ala Tyr Pro Ala Val Ser Thr  
                   20                  25                  30

Phe Glu Lys His Ser Ser Ser Gly His Ala Val Glu Phe His Asn Leu  
           35                  40                  45

Pro Pro Asn Ser Ala Arg Ser Gln Ser Glu Arg Pro Val Phe Pro Cys  
       50                  55                  60

Trp Trp Leu Gln Phe Arg Asn Ser Lys Pro Cys Ser Asp Tyr Cys Leu  
   65                  70                  75                  80

Ser Leu Ile Val Asn Leu Leu Glu Asp Trp Gly Pro Cys Ala Glu His  
                   85                  90                  95

Gly Glu His His Ile Arg Ile Pro Arg Thr Pro Ser Arg Val Thr Gly  
                   100                  105                  110

Gly Val Phe Leu Val Asp Lys Asn Pro His Asn Thr Ala Glu Ser Arg  
       115                  120                  125

Leu Val Val Asp Phe Ser Gln Phe Ser Arg Gly Asn Tyr Arg Val Ser  
   130                  135                  140

Trp Pro Lys Phe Ala Val Pro Asn Leu Gln Ser Leu Thr Asn Leu Leu  
   145                  150                  155                  160

Ser Ser Asn Leu Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr  
                   165                  170                  175

His Leu Pro Leu His Pro Ala Ala Met Pro His Leu Leu Val Gly Ser  
                   180                  185                  190

Ser Gly Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile  
       195                  200                  205

Leu Asn Asn Gln His Gly Thr Met Pro Asp Leu His Asp Tyr Cys Ser  
       210                  215                  220

Arg Asn Leu Tyr Val Ser Leu Leu Leu Leu Tyr Gln Thr Phe Gly Arg  
   225                  230                  235                  240

- 20 -

Lys Leu His Leu Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile  
245 250 255

Pro Met Gly Val Gly Leu Ser Pro Phe Leu Leu Ala Gln Phe Thr Ser  
260 265 270

Ala Ile Cys Ser Val Val Arg Arg Ala Phe Pro His Cys Leu Ala Phe  
275 280 285

Ser Tyr Met Asp Asp Val Val Leu Gly Ala Lys Ser Val Gln His Leu  
290 295 300

Glu Ser Leu Phe Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly Ile  
305 310 315 320

His Leu Asn Pro Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu Asn Phe  
325 330 335

Met Gly Tyr Val Ile Gly Cys Tyr  
340

```
<210> 22
<211> 336
<212> PRT
<213> artificial sequence
```

<220>  
<223> Pol Trans 4

```

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> x =any amino acid

```

<400> 22

Ala Ser Lys Ser Ala Ser Ser Ile Tyr Gln Ser Pro Val Gly Thr Ala  
1 5 10 15

Ala Tyr Pro Ala Val Ser Thr Xaa Glu Lys His Ser Ser Ser Gly His  
20 25 30

Ala Val Glu Leu His Asn Leu Pro Pro Asn Ser Glu Arg Ser Gln Gly  
35 40 45

- 21 -

Glu Arg Pro Val Phe Pro Cys Trp Trp Leu Gln Phe Arg Asn Ser Lys  
 50 55 60

Pro Cys Ser Asp Tyr Cys Leu Ser His Ile Val Asn Leu Leu Glu Asp  
 65 70 75 80

Trp Gly Pro Cys Ala Glu His Gly Glu His His Ile Arg Ile Pro Arg  
 85 90 95

Thr Pro Ala Arg Val Thr Gly Gly Val Phe Leu Val Asp Lys Asn Pro  
 100 105 110

His Asn Thr Ala Glu Ser Arg Leu Val Val Asp Phe Ser Gln Phe Ser  
 115 120 125

Arg Gly Asn Tyr Arg Val Ser Trp Pro Lys Phe Ala Val Pro Asn Leu  
 130 135 140

Gln Ser Leu Thr Asn Leu Leu Ser Ser Asn Leu Ser Trp Leu Ser Leu  
 145 150 155 160

Asp Val Ser Ala Ala Phe Tyr His Leu Pro Leu His Pro Ala Ala Met  
 165 170 175

Pro His Leu Leu Val Gly Ser Ser Gly Leu Ser Arg Tyr Val Ala Arg  
 180 185 190

Leu Ser Ser Asn Ser Arg Ile Phe Asn His Gln Arg Gly Asn Met Gln  
 195 200 205

Asn Leu His Asp Cys Cys Ser Arg Asn Leu Tyr Val Ser Leu Leu Leu  
 210 215 220

Leu Tyr Gln Thr Phe Gly Arg Lys Leu His Leu Tyr Ser His Pro Ile  
 225 230 235 240

Ile Leu Gly Phe Arg Lys Ile Pro Met Gly Val Gly Leu Ser Pro Phe  
 245 250 255

Leu Leu Ala Gln Phe Thr Ser Ala Ile Cys Ser Val Val Arg Arg Ala  
 260 265 270

Phe Pro His Cys Leu Ala Phe Ser Tyr Met Asp Asp Val Val Leu Gly

- 22 -

275

280

285

Ala Lys Ser Val Gln His Leu Glu Ser Leu Phe Thr Ala Val Thr Asn  
 290 295 300

Phe Leu Leu Ser Leu Gly Ile His Leu Asn Pro Asn Lys Thr Lys Arg  
 305 310 315 320

Trp Gly Tyr Ser Leu Asn Phe Met Gly Tyr Val Ile Gly Trp Tyr Gly  
 325 330 335

<210> 23  
 <211> 226  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of Pre 1

<220>  
 <221> MISC\_FEATURE  
 <222> (120)..(120)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (208)..(208)  
 <223> x = any amino acid

<400> 23

Met Glu Asn Ile Thr Ser Gly Phe Leu Gly Pro Leu Leu Val Leu Gln  
 1 5 10 15

Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro Gln Ser Leu  
 20 25- 30

Asp Ser Trp Trp Thr Ser Leu Asn Phe Leu Gly Gly Thr Thr Val Cys  
 35 40 45

Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser Pro Thr Ser  
 50 55 60

Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu Arg Arg Phe  
 65 70 75 80

- 23 -

Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe Leu Leu Val  
85 90 95

Leu Leu Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu Ile Pro Gly  
100 105 110

Ser Ser Thr Thr Ser Ala Gly Xaa Cys Arg Thr Cys Thr Thr Thr Ala  
115 120 125

Gln Gly Thr Ser Met Tyr Pro Ser Cys Cys Cys Thr Lys Pro Ser Asp  
130 135 140

Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala Phe Gly Lys  
145 150 155 160

Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu Ser Leu Leu  
165 170 175

Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr Val Trp Leu  
180 185 190

Ser Val Ile Trp Met Met Trp Tyr Trp Gly Pro Ser Leu Tyr Ser Xaa  
195 200 205

Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys Leu Trp Val  
210 215 220

Tyr Ile  
225

<210> 24  
<211> 309  
<212> PRT  
<213> artificial sequence

<220>  
<223> HBsAg Trans of 2  
<400> 24

Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro Thr Pro  
1 5 10 15

Leu Ser Pro Pro Leu Arg Asn Thr His Pro Gln Ala Met Gln Trp Asn  
20 25 30

- 24 -

Ser Thr Thr Phe His Gln Thr Leu Gln Asp Pro Arg Val Arg Gly Leu  
 35 40 45

Tyr Phe Pro Ala Gly Gly Ser Ser Ser Gly Thr Val Asn Pro Val Leu  
 50 55 60

Thr Thr Ala Ser Pro Leu Ser Ser Ile Phe Ser Arg Ile Gly Asp Pro  
 65 70 75 80

Ala Leu Asn Met Glu Asn Ile Thr Ser Gly Phe Leu Gly Pro Leu Leu  
 85 90 95

Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro  
 100 105 110

Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Leu Gly Gly Thr  
 115 120 125

Thr Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser  
 130 135 140

Pro Thr Ser Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu  
 145 150 155 160

Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe  
 165 170 175

Leu Leu Val Leu Leu Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu  
 180 185 190

Ile Pro Gly Ser Ser Thr Thr Ser Thr Gly Pro Cys Arg Thr Cys Met  
 195 200 205

Thr Thr Ala Gln Gly Thr Ser Met Tyr Pro Ser Cys Cys Cys Thr Lys  
 210 215 220

Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 225 230 235 240

Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu  
 245 250 255

- 25 -

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
                   260                  265                  270

Val Trp Leu Ser Val Ile Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
                   275                  280                  285

Tyr Ser Ile Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys  
                   290                  295                  300

Leu Trp Val Tyr Ile  
                   305

<210> 25  
 <211> 309  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of 3

<400> 25

Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro Thr Pro  
   1                  5                  10                  15

Leu Ser Pro Pro Leu Arg Asn Thr His Pro Gln Ala Met Gln Trp Asn  
                   20                  25                  30

Ser Thr Thr Phe His Gln Thr Leu Gln Asp Pro Arg Val Arg Gly Leu  
                   35                  40                  45

Tyr Phe Pro Ala Gly Gly Ser Ser Ser Gly Thr Val Asn Pro Val Leu  
                   50                  55                  60

Thr Thr Ala Ser Pro Leu Ser Ser Ile Phe Ser Arg Ile Gly Asp Pro  
   65                  70                  75                  80

Ala Leu Asn Met Glu Asn Ile Thr Ser Gly Phe Leu Gly Pro Leu Leu  
                   85                  90                  95

Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro  
                   100                  105                  110

Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Leu Gly Gly Thr  
                   115                  120                  125

- 26 -

Thr Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser  
 130 135 140

Pro Thr Ser Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu  
 145 150 155 160

Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe  
 165 170 175

Leu Leu Val Leu Leu Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu  
 180 185 190

Ile Pro Gly Ser Ser Thr Thr Ser Thr Gly Pro Cys Arg Thr Cys Met  
 195 200 205

Thr Thr Ala Gln Gly Thr Ser Met Tyr Pro Ser Cys Cys Cys Thr Lys  
 210 215 220

Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 225 230 235 240

Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu  
 245 250 255

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
 260 265 270

Val Trp Leu Ser Val Ile Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
 275 280 285

Tyr Ser Ile Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys  
 290 295 300

Leu Trp Val Tyr Ile  
 305

<210> 26  
 <211> 309  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of 4



- 27 -

<220>  
 <221> MISC\_FEATURE  
 <222> (21)..(21)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (46)..(46)  
 <223> x = any amino acid

<400> 26

Pro Pro Pro Pro Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro Thr Pro  
 1 5 10 15

Leu Ser Pro Pro Xaa Arg Asn Thr His Pro Gln Ala Met Gln Trp Asn  
 20 25 30

Ser Thr Thr Phe His Gln Thr Leu Lys Asp Pro Arg Val Xaa Gly Leu  
 35 40 45

Tyr Phe Pro Ala Gly Gly Ser Ser Ser Gly Thr Val Asn Pro Val Pro  
 50 55 60

Thr Thr Ala Ser Pro Ile Ser Ser Ile Phe Ser Arg Ile Gly Asp Pro  
 65 70 75 80

Ala Leu Asn Met Glu Asn Ile Thr Ser Gly Phe Leu Gly Pro Leu Leu  
 85 90 95

Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro  
 100 105 110

Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Leu Gly Gly Thr  
 115 120 125

Thr Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser  
 130 135 140

Pro Thr Ser Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu  
 145 150 155 160

Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe  
 165 170 175

- 28 -

Leu Leu Val Leu Leu Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu  
 180 185 190

Ile Pro Gly Ser Ser Thr Thr Ser Ala Gly Thr Cys Arg Thr Cys Thr  
 195 200 205

Thr Ala Ala Gln Gly Thr Ser Met Tyr Pro Ser Cys Cys Cys Thr Lys  
 210 215 220

Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 225 230 235 240

Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu  
 245 250 255

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
 260 265 270

Val Trp Leu Ser Val Ile Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
 275 280 285

Tyr Ser Ile Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys  
 290 295 300

Leu Trp Ala Tyr Ile  
 305

<210> 27  
 <211> 656  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> S0

<220>  
 <221> misc\_feature  
 <222> (561)..(561)  
 <223> n = any nucleotide

<400> 27  
 cgcagagtct agactcgtgg tggacttctc tcaattttcg aggggggact accgtgtgtc 60  
 ttggccaaaa ttgcagtc ccaacctcca atcactcacc aacctcctgt cctccaactt 120  
 gtcttggtta tcgctggatg tgtctgcggc gttttatcat cttcctcttc atcctgtgc 180

- 29 -

```

tatgcctcat cttcttggtg gttcttctgg actgtcaagg tatgttgccc gtttgcctc   240
taattccagg atcctcaacc accagcacgg gaccatgccg aacctgcacg actcctgctc   300
aaggaaacct tacggttccc tcatgttgct gtaccaaacc ttcggacgga aattgcacct   360
gtattcccat cccatcatcc tgggctttcg gaaaattcct atgggagtg gctcagccc   420
gtttctcctg gctcagttta ctagtgccat ttgttcagtg gttagtaggg ctttccccc   480
ctgtctggct tttagttata tggatgatgt ggtattgggg gccagtctg tatcgcatct   540
tgagtccttt tttaccgctg ntaccaatct tcttttctct ttgggtatac atttaaacc   600
taacaaaaca aaaagatggg gttactccct acattttatg ggctatgtca ttggat   656

```

```

<210> 28
<211> 625
<212> DNA
<213> artificial sequence

```

```

<220>
<223> S6

```

```

<220>
<221> misc_feature
<222> (10)..(10)
<223> n = any nucleotide

```

```

<400> 28
ttactcaccn acctcctgtc ctccaacttg tcttggttat cgctggatgt gtctgcgggc   60
ttttatcatc ttctcttca tctgtctgt atgcctcatc ttcttggttg ttcttctgga   120
ctgtcaaggt atgttgcccg tttgtcctct aattccagga tcttcaacca ccagcagggg   180
accatgccga acctgcacga ctctgtctca aggaacctct acggttccct catgttgctg   240
taccaaacct tcggacggaa attgcacctg tattcccatc ccatcatcct gggctttcgg   300
aaaattccta tgggagtggg cctcagcccg tttctcatgg cttagtttac tagtgccatt   360
tgttcagtgg ttcttagggc tttccccac tgtctggctt ttggttatgt ggatgatgtg   420
gtattggggg ccaagtctgt atcgcatctt gagtcccttt ttaccgctgt taccaatttt   480
cttttgtctt tgggtataca tttaaatcct aacaaaacaa aaagatgggg ttactoccta   540
cattttatgg gctatgtcat tggatgtcat gggtccttgc cacaagaaca catcagacaa   600
aaaatcaaag aatgttttag aaaac   625

```

```

<210> 29

```

- 30 -

&lt;211&gt; 1033

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; S8

&lt;400&gt; 29

```

tgcccccttct gcctccacca atcgccagtc aggaaggcag cctaccccg cgtctccacc      60
tttgagagac actcatcctc aggccatgca gtggaactca acaaccttcc accaaactct      120
gcaagatccc agagtgaaag gcctgtatct ccctgctggg ggctccagtt caggaacagt      180
aaacctgttt ccgactactg cctctcactc atcgtcaatc ttctcgagga ttgggggtccc      240
tgcgctgaac atggagaaca tcacatcagg actcctagga cccctttctg tggttacaggc      300
gggggtttttc ttgttgacaa gaatcctcac aataccgcag agtctagact cgtgggtggac      360
ttctctcaat ttctgagggg ggactaccgt gtgtcttggc caaaattcgc agtccccaac      420
ctccaatcac tcaccaacct cctgtcctcc aacttgcctc gggtatcgct ggatgtgtct      480
gcggcgtttt atcatcttcc tcttcactct gctgctatgc ctcatcttct tgttggttct      540
tctggactgt caaggatatg tgcccgtttg tcctctaatt ccaggatcct caaccaccag      600
caggggacca tgccgaacct gcacgactcc tgctcaagga acctctacgg ttccctcatg      660
ttgtgttacc aaaccttcgg acggaaattg cacctgtatt cccatcccat catcctgggc      720
tttcggaaaaa ttcctatggg agtgggcctc agcccgtttc tcatggctca gtttactagt      780
gccatttggt cagtgggttc tagggcttcc cccactgtc tggcttttgg ttatgtggat      840
gatgtggtat tgggggccaa gtctgtatcg catcttgagt ccccttttac cgtgtgtacc      900
aattttcttt tgtcttttgg tatacattta aatcctaaca aaacaaaaag atgggggttac      960
tccttacatt ttatgggcta tgcattgga tgcattgggt ccttgccaca agaacacatc     1020
agacaaaaaa tca                                     1033

```

&lt;210&gt; 30

&lt;211&gt; 1100

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; S12

&lt;400&gt; 30

```

ttttggggag ccttcaggct cagggcatat tacaaactct gccagcaaat ccacctctg      60
cctccaccaa tcgccagtca ggaaggcagc ctaccccgct gtctccacct ttgagagaca     120

```

- 31 -

```

ctcatcctca ggccatgcag tggaactcaa caaccttcca ccaaactctg caagatccca 180
gagtgaaagg cctgtatttc cctgctggtg gctccagttc aggaacagta aaccctgttc 240
cgactactgc ctctcactca tcgtcaatct tctcgaggat tgggggtccct gcgctgaaca 300
tgggagaacat cacatcagga ctcttaggac cccttctcgt gttacaggcg gggtttttct 360
tgttgacaag aatcctcaca ataccgcaga gtctagactc gtgggtggact tctctcaatt 420
ttcgaggggg gactaccgtg tgtcttggcc aaaattcgca gtccccaacc tccaatcact 480
caccaacctc ctgtcctcca acttgtcctg gttatcgctg gatgtgtctg cggcgtttta 540
tcattcttct cttcatcctg ctgctatgcc tcatottctt gttggttctt ctggactgtc 600
aaggatatgt gcccgtttgt cctctaattc caggatcctc aaccaccagc aggggaccat 660
gccgaacctg cagcactcct gctcaaggaa cctctacggt tccctcatgt tgctgtacca 720
aaccttcgga cggaattgc acctgtatcc ccattccatc atcctgggct ttcggaaaat 780
tcctatggga gtgggcctca gcccgtttct catggctcag tttactagtg ccatttgttc 840
agtgggtcgt agggctttcc cccactgtct ggcttttgggt tatgtggatg atgtgggtatt 900
gggggccaag tctgtatcgc atcttgagtc cctttttacc gctgttacca attttctttt 960
gtctttgggt atacatttaa atcctaacaa aacaaaaaga tgggggttact ccctacattt 1020
tatgggctat gtcattggat gtcattgggtc cttgccacaa gaacacatca gacaaaaaat 1080
caaagaatgt ttagaaaaac 1100

```

```

<210> 31
<211> 987
<212> DNA
<213> artificial sequence

```

```

<220>
<223> S15

```

```

<220>
<221> misc_feature
<222> (329)..(329)
<223> y = any nucleotide

```

```

<220>
<221> misc_feature
<222> (943)..(943)
<223> y = any nucleotide

```

```

<400> 31

```

- 32 -

```

tacaaaacttt gccagcaaat ccacctcctg cctccaccaa tcgccagtca ggaaggcagc      60
ctaccccgct gtetccacct ttgagagaca ctcatcctca ggccatgcag tggaactcaa      120
caaccttcca ccaaactctg caagatccca gagtgaaagg cctgtatttc cctgctggtg      180
gtccagttc aggaacagta aacctgttc cgactactgc ctctcactca tcgtcaatct      240
tctcgaggat tggggtccct gcgctgaaca tggagaacat cacatcagga ctctaggagc      300
cccttctcgt gttacaggcg gggtttttyt tgttgacaag aatcctcaca ataccgcaga      360
gtctagactc gtggtggact tctctcaatt ttcgaggggg gactaccgtg tgtcttggcc      420
aaaattegca gtccccaacc tccaatcact caccaacctc ctgtcctcca acttgtcctg      480
gttatcgctg gatgtgtctg cggcgtttta tcatcttctc cttcatcctg ctgctatgcc      540
tcatcttctt gttggctcta ctggactgtc aaggatgtt gcccgtttgt cctctaattc      600
caggatctc aaccaccagc aggggaccat gccgaacctg cagcactcct gctcaaggaa      660
cctctacggg tccctcatgt tgctgtacca aaccttcgga cggaaattgc acctgtattc      720
ccatcccatc atcctgggct ttcggaaaat tctatggga gtgggcctca gcccgtttct      780
catggctcag tttactagtg ccatttgttc agtgggtcgt agggctttcc cccactgtct      840
ggcttttggt tatgtggatg atgtggtatt gggggccaag tctgtatcgc atcttgagtc      900
cctttttacc gctgttacca attttctttt gtctttgggt atrcatttaa atcctaacia      960
aacaaaaaga tgggggttact ccctaca                                     987

```

```

<210> 32
<211> 350
<212> PRT
<213> artificial sequence

```

```

<220>
<223> Pol Trans S0

```

```

<220>
<221> MISC_FEATURE
<222> (309)..(309)
<223> x = any amino acid

```

```

<220>
<221> MISC_FEATURE
<222> (345)..(345)
<223> x = any amino acid

```

```

<400> 32

```

- 33 -

Ser Gly His Thr Thr Asn Phe Ala Ser Lys Ser Thr Ser Cys Leu His  
 1 5 10 15

Gln Ser Pro Val Arg Lys Ala Ala Tyr Pro Ala Val Ser Thr Phe Glu  
 20 25 30

Arg His Ser Ser Ser Gly His Ala Val Glu Leu Asn Asn Leu Pro Pro  
 35 40 45

Asn Ser Ala Arg Ser Gln Ser Glu Arg Pro Val Phe Pro Cys Trp Trp  
 50 55 60

Leu Gln Phe Arg Asn Ser Lys Pro Cys Ser Asp Tyr Cys Leu Ser Leu  
 65 70 75 80

Ile Val Asn Leu Leu Glu Asp Trp Gly Pro Cys Ala Glu His Gly Glu  
 85 90 95

His His Ile Arg Thr Pro Arg Thr Pro Ser Arg Val Thr Gly Gly Val  
 100 105 110

Phe Leu Val Asp Lys Asn Pro His Asn Thr Ala Glu Ser Arg Leu Val  
 115 120 125

Val Asp Phe Ser Gln Phe Ser Arg Gly Asp Tyr Arg Val Ser Trp Pro  
 130 135 140

Lys Phe Ala Val Pro Asn Leu Gln Ser Leu Thr Asn Leu Leu Ser Ser  
 145 150 155 160

Asn Leu Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His Leu  
 165 170 175

Pro Leu His Pro Ala Ala Met Pro His Leu Leu Val Gly Ser Ser Gly  
 180 185 190

Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile Leu Asn  
 195 200 205

His Gln His Gly Thr Met Pro Asn Leu His Asp Ser Cys Ser Arg Asn  
 210 215 220

Leu Tyr Gly Ser Leu Met Leu Leu Tyr Gln Thr Phe Gly Arg Lys Leu

- 34 -

225                      230                      235                      240

His Leu Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro Met  
                        245                      250                      255

Gly Val Gly Leu Ser Pro Phe Leu Leu Ala Gln Phe Thr Ser Ala Ile  
                        260                      265                      270

Cys Ser Val Val Arg Arg Ala Phe Pro His Cys Leu Ala Phe Ser Tyr  
                        275                      280                      285

Met Asp Asp Val Val Leu Gly Ala Lys Ser Val Ser His Leu Glu Ser  
                        290                      295                      300

Leu Phe Thr Ala Xaa Thr Asn Phe Leu Leu Ser Leu Gly Ile His Leu  
305                      310                      315                      320

Asn Pro Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu His Phe Met Gly  
                        325                      330                      335

Tyr Val Ile Gly Cys His Gly Ser Xaa Pro Gln Glu His Ile  
                        340                      345                      350

<210> 33  
<211> 181  
<212> PRT  
<213> artificial sequence

<220>  
<223> Pol Trans S6

<400> 33

Ser Asn Leu Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His  
1                      5                      10                      15

Leu Pro Leu His Pro Ala Ala Met Pro His Leu Leu Val Gly Ser Ser  
                        20                      25                      30

Gly Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile Leu  
                        35                      40                      45

Asn His Gln Gln Gly Thr Met Pro Asn Leu His Asp Ser Cys Ser Arg  
50                      55                      60



- 35 -

Asn Leu Tyr Gly Ser Leu Met Leu Leu Tyr Gln Thr Phe Gly Arg Lys  
 65 70 75 80

Leu His Leu Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro  
 85 90 95

Met Gly Val Gly Leu Ser Pro Phe Leu Met Ala Gln Phe Thr Ser Ala  
 100 105 110

Ile Cys Ser Val Val Arg Arg Ala Phe Pro His Cys Leu Ala Phe Gly  
 115 120 125

Tyr Val Asp Asp Val Val Leu Gly Ala Lys Ser Val Ser His Leu Glu  
 130 135 140

Ser Leu Phe Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly Ile His  
 145 150 155 160

Leu Asn Pro Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu His Phe Met  
 165 170 175

Gly Tyr Val Ile Gly  
 180

<210> 34

<211> 340

<212> PRT

<213> artificial sequence

<220>

<223> Pol Trans S8

<400> 34

Cys Pro Phe Cys Leu His Gln Ser Pro Val Arg Lys Ala Ala Tyr Pro  
 1 5 10 15

Ala Val Ser Thr Phe Glu Arg His Ser Ser Ser Gly His Ala Val Glu  
 20 25 30

Leu Asn Asn Leu Pro Pro Asn Ser Ala Arg Ser Gln Ser Glu Arg Pro  
 35 40 45

Val Phe Pro Cys Trp Trp Leu Gln Phe Arg Asn Ser Lys Pro Cys Ser  
 50 55 60

- 36 -

Asp Tyr Cys Leu Ser Leu Ile Val Asn Leu Leu Glu Asp Trp Gly Pro  
 65 70 75 80

Cys Ala Glu His Gly Glu His His Ile Arg Thr Pro Arg Thr Pro Ser  
 85 90 95

Arg Val Thr Gly Gly Val Phe Leu Val Asp Lys Asn Pro His Asn Thr  
 100 105 110

Ala Glu Ser Arg Leu Val Val Asp Phe Ser Gln Phe Ser Arg Gly Asp  
 115 120 125

Tyr Arg Val Ser Trp Pro Lys Phe Ala Val Pro Asn Leu Gln Ser Leu  
 130 135 140

Thr Asn Leu Leu Ser Ser Asn Leu Ser Trp Leu Ser Leu Asp Val Ser  
 145 150 155 160

Ala Ala Phe Tyr His Leu Pro Leu His Pro Ala Ala Met Pro His Leu  
 165 170 175

Leu Val Gly Ser Ser Gly Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser  
 180 185 190

Asn Ser Arg Ile Leu Asn His Gln Gln Gly Thr Met Pro Asn Leu His  
 195 200 205

Asp Ser Cys Ser Arg Asn Leu Tyr Gly Ser Leu Met Leu Leu Tyr Gln  
 210 215 220

Thr Phe Gly Arg Lys Leu His Leu Tyr Ser His Pro Ile Ile Leu Gly  
 225 230 235 240

Phe Arg Lys Ile Pro Met Gly Val Gly Leu Ser Pro Phe Leu Met Ala  
 245 250 255

Gln Phe Thr Ser Ala Ile Cys Ser Val Val Arg Arg Ala Phe Pro His  
 260 265 270

Cys Leu Ala Phe Gly Tyr Val Asp Asp Val Val Leu Gly Ala Lys Ser  
 275 280 285

- 37 -

Val Ser His Leu Glu Ser Leu Phe Thr Ala Val Thr Asn Phe Leu Leu  
 290 295 300

Ser Leu Gly Ile His Leu Asn Pro Asn Lys Thr Lys Arg Trp Gly Tyr  
 305 310 315 320

Ser Leu His Phe Met Gly Tyr Val Ile Gly Cys His Gly Ser Leu Pro  
 325 330 335

Gln Glu His Ile  
 340

<210> 35  
 <211> 340  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> Pol Trans S12

<400> 35

Ser Gly His Ile Thr Asn Ser Ala Ser Lys Ser Thr Ser Cys Leu His  
 1 5 10 15

Gln Ser Pro Val Arg Lys Ala Ala Tyr Pro Ala Val Ser Thr Phe Glu  
 20 25 30

Arg His Ser Ser Ser Gly His Ala Val Glu Leu Asn Asn Leu Pro Pro  
 35 40 45

Asn Ser Ala Arg Ser Gln Ser Glu Arg Pro Val Phe Pro Cys Trp Trp  
 50 55 60

Leu Gln Phe Arg Asn Ser Lys Pro Cys Ser Asp Tyr Cys Leu Ser Leu  
 65 70 75 80

Ile Val Asn Leu Leu Glu Asp Trp Gly Pro Cys Ala Glu His Gly Glu  
 85 90 95

His His Ile Arg Thr Pro Arg Thr Pro Ser Arg Val Thr Gly Gly Val  
 100 105 110

Phe Leu Val Asp Lys Asn Pro His Asn Thr Ala Glu Ser Arg Leu Val  
 115 120 125

- 38 -

Val Asp Phe Ser Gln Phe Ser Arg Gly Asp Tyr Arg Val Ser Trp Pro  
 130 135 140

Lys Phe Ala Val Pro Asn Leu Gln Ser Leu Thr Asn Leu Leu Ser Ser  
 145 150 155 160

Asn Leu Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His Leu  
 165 170 175

Pro Leu His Pro Ala Ala Met Pro His Leu Leu Val Gly Ser Ser Gly  
 180 185 190

Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile Leu Asn  
 195 200 205

His Gln Gln Gly Thr Met Pro Asn Leu His Asp Ser Cys Ser Arg Asn  
 210 215 220

Leu Tyr Gly Ser Leu Met Leu Leu Tyr Gln Thr Phe Gly Arg Lys Leu  
 225 230 235 240

His Leu Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro Met  
 245 250 255

Gly Val Gly Leu Ser Pro Phe Leu Met Ala Gln Phe Thr Ser Ala Ile  
 260 265 270

Cys Ser Val Val Arg Arg Ala Phe Pro-His Cys Leu Ala Phe Gly Tyr  
 275 280 285

Val Asp Asp Val Val Leu Gly Ala Lys Ser Val Ser His Leu Glu Ser  
 290 295 300

Leu Phe Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly Ile His Leu  
 305 310 315 320

Asn Pro Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu His Phe Met Gly  
 325 330 335

Tyr Val Ile Gly  
 340

- 39 -

<210> 36  
 <211> 328  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> Pol Trans S15

<220>  
 <221> MISC\_FEATURE  
 <222> (110)..(110)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (314)..(314)  
 <223> x = any amino acid

<400> 36

Thr Asn Phe Ala Ser Lys Ser Thr Ser Cys Leu His Gln Ser Pro Val  
 1 5 10 15

Arg Lys Ala Ala Tyr Pro Ala Val Ser Thr Phe Glu Arg His Ser Ser  
 20 25 30

Ser Gly His Ala Val Glu Leu Asn Asn Leu Pro Pro Asn Ser Ala Arg  
 35 40 45

Ser Gln Ser Glu Arg Pro Val Phe Pro Cys Trp Trp Leu Gln Phe Arg  
 50 55 60

Asn Ser Lys Pro Cys Ser Asp Tyr Cys Leu Ser Leu Ile Val Asn Leu  
 65 70 75 80

Leu Glu Asp Trp Gly Pro Cys Ala Glu His Gly Glu His His Ile Arg  
 85 90 95

Thr Pro Arg Thr Pro Ser Arg Val Thr Gly Gly Val Phe Xaa Val Asp  
 100 105 110

Lys Asn Pro His Asn Thr Ala Glu Ser Arg Leu Val Val Asp Phe Ser  
 115 120 125

Gln Phe Ser Arg Gly Asp Tyr Arg Val Ser Trp Pro Lys Phe Ala Val  
 130 135 140

- 40 -

Pro Asn Leu Gln Ser Leu Thr Asn Leu Leu Ser Ser Asn Leu Ser Trp  
 145 150 155 160

Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His Leu Pro Leu His Pro  
 165 170 175

Ala Ala Met Pro His Leu Leu Val Gly Ser Thr Gly Leu Ser Arg Tyr  
 180 185 190

Val Ala Arg Leu Ser Ser Asn Ser Arg Ile Leu Asn His Gln Gln Gly  
 195 200 205

Thr Met Pro Asn Leu His Asp Ser Cys Ser Arg Asn Leu Tyr Gly Ser  
 210 215 220

Leu Met Leu Leu Tyr Gln Thr Phe Gly Arg Lys Leu His Leu Tyr Ser  
 225 230 235 240

His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro Met Gly Val Gly Leu  
 245 250 255

Ser Pro Phe Leu Met Ala Gln Phe Thr Ser Ala Ile Cys Ser Val Val  
 260 265 270

Arg Arg Ala Phe Pro His Cys Leu Ala Phe Gly Tyr Val Asp Asp Val  
 275 280 285

Val Leu Gly Ala Lys Ser Val Ser His Leu Glu Ser Leu Phe Thr Ala  
 290 295 300

Val Thr Asn Phe Leu Leu Ser Leu Gly Xaa His Leu Asn Pro Asn Lys  
 305 310 315 320

Thr Lys Arg Trp Gly Tyr Ser Leu  
 325

<210> 37  
 <211> 197  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of S0

- 41 -

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (187)..(187)

&lt;223&gt; x = any amino acid

&lt;400&gt; 37

Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Arg Gly Gly Thr  
 1 5 10 15

Thr Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser  
 20 25 30

Pro Thr Ser Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu  
 35 40 45

Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Cys Leu Ile Phe  
 50 55 60

Leu Leu Val Leu Leu Asp Cys Gln Gly Met Leu Pro Val Cys Pro Leu  
 65 70 75 80

Ile Pro Gly Ser Ser Thr Thr Ser Thr Gly Pro Cys Arg Thr Cys Thr  
 85 90 95

Thr Pro Ala Gln Gly Thr Ser Thr Val Pro Ser Cys Cys Cys Thr Lys  
 100 105 110

Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 115 120 125

Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu  
 130 135 140

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
 145 150 155 160

Val Trp Leu Leu Val Ile Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
 165 170 175

Tyr Arg Ile Leu Ser Pro Phe Leu Pro Leu Xaa Pro Ile Phe Phe Cys  
 180 185 190

Leu Trp Val Tyr Ile

- 42 -

195

<210> 38  
 <211> 161  
 <212> PRT  
 <213> artificial sequence

&lt;220&gt;

&lt;223&gt; HBsAg Trans of S6

&lt;400&gt; 38

Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu Arg Arg Phe Ile  
 1 5 10 15

Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe Leu Leu Val Leu  
 20 25 30

Leu Asp Cys Gln Gly Met Leu Pro Val Cys Pro Leu Ile Pro Gly Ser  
 35 40 45

Ser Thr Thr Ser Arg Gly Pro Cys Arg Thr Cys Thr Thr Pro Ala Gln  
 50 55 60

Gly Thr Ser Thr Val Pro Ser Cys Cys Cys Thr Lys Pro Ser Asp Gly  
 65 70 75 80

Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala Phe Gly Lys Phe  
 85 90 95

Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu Ser Leu Leu Val  
 100 105 110

Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr Val Trp Leu Leu  
 115 120 125

Val Met Trp Met Met Trp Tyr Trp Gly Pro Ser Leu Tyr Arg Ile Leu  
 130 135 140

Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys Leu Trp Val Tyr  
 145 150 155 160

Ile



- 43 -

<210> 39  
 <211> 160  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of S8

<400> 39

Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu Arg Arg Phe Ile Ile  
 1 5 10 15

Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe Leu Leu Val Leu Leu  
 20 25 30

Asp Cys Gln Gly Met Leu Pro Val Cys Pro Leu Ile Pro Gly Ser Ser  
 35 40 45

Thr Thr Ser Arg Gly Pro Cys Arg Thr Cys Thr Thr Pro Ala Gln Gly  
 50 55 60

Thr Ser Thr Val Pro Ser Cys Cys Cys Thr Lys Pro Ser Asp Gly Asn  
 65 70 75 80

Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala Phe Gly Lys Phe Leu  
 85 90 95

Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu Ser Leu Leu Val Pro  
 100 105 110

Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr Val Trp Leu Leu Val  
 115 120 125

Met Trp Met Met Trp Tyr Trp Gly Pro Ser Leu Tyr Arg Ile Leu Ser  
 130 135 140

Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys Leu Trp Val Tyr Ile  
 145 150 155 160

<210> 40  
 <211> 325  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of S12

- 44 -

&lt;400&gt; 40

Leu Gly Ser Pro Gln Ala Gln Gly Ile Leu Gln Thr Leu Pro Ala Asn  
 1 5 10 15

Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro Thr Pro  
 20 25 30

Leu Ser Pro Pro Leu Arg Asp Thr His Pro Gln Ala Met Gln Trp Asn  
 35 40 45

Ser Thr Thr Phe His Gln Thr Leu Gln Asp Pro Arg Val Lys Gly Leu  
 50 55 60

Tyr Phe Pro Ala Gly Gly Ser Ser Ser Gly Thr Val Asn Pro Val Pro  
 65 70 75 80

Thr Thr Ala Ser His Ser Ser Ser Ile Phe Ser Arg Ile Gly Val Pro  
 85 90 95

Ala Leu Asn Met Glu Asn Ile Thr Ser Gly Leu Leu Gly Pro Leu Leu  
 100 105 110

Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro  
 115 120 125

Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Arg Gly Gly Thr  
 130 135 140

Thr Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser  
 145 150 155 160

Pro Thr Ser Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu  
 165 170 175

Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe  
 180 185 190

Leu Leu Val Leu Leu Asp Cys Gln Gly Met Leu Pro Val Cys Pro Leu  
 195 200 205

Ile Pro Gly Ser Ser Thr Thr Ser Arg Gly Pro Cys Arg Thr Cys Thr  
 210 215 220

- 45 -

Thr Pro Ala Gln Gly Thr Ser Thr Val Pro Ser Cys Cys Cys Thr Lys  
 225 230 235 240

Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 245 250 255

Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu  
 260 265 270

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
 275 280 285

Val Trp Leu Leu Val Met Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
 290 295 300

Tyr Arg Ile Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys  
 305 310 315 320

Leu Trp Val Tyr Ile  
 325

<210> 41  
 <211> 309  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of S15

<220>  
 <221> MISC\_FEATURE  
 <222> (308)..(308)  
 <223> x = any amino acid

<400> 41

Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro Thr Pro  
 1 5 10 15

Leu Ser Pro Pro Leu Arg Asp Thr His Pro Gln Ala Met Gln Trp Asn  
 20 25 30

Ser Thr Thr Phe His Gln Thr Leu Gln Asp Pro Arg Val Lys Gly Leu  
 35 40 45

- 46 -

Tyr Phe Pro Ala Gly Gly Ser Ser Ser Gly Thr Val Asn Pro Val Pro  
 50 55 60

Thr Thr Ala Ser His Ser Ser Ser Ile Phe Ser Arg Ile Gly Val Pro  
 65 70 75 80

Ala Leu Asn Met Glu Asn Ile Thr Ser Gly Leu Leu Gly Pro Leu Leu  
 85 90 95

Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro  
 100 105 110

Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Arg Gly Gly Thr  
 115 120 125

Thr Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr Ser Asn His Ser  
 130 135 140

Pro Thr Ser Cys Pro Pro Thr Cys Pro Gly Tyr Arg Trp Met Cys Leu  
 145 150 155 160

Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe  
 165 170 175

Leu Leu Ala Leu Leu Asp Cys Gln Gly Met Leu Pro Val Cys Pro Leu  
 180 185 190

Ile Pro Gly Ser Ser Thr Thr Ser Arg Gly Pro Cys Arg Thr Cys Thr  
 195 200 205

Thr Pro Ala Gln Gly Thr Ser Thr Val Pro Ser Cys Cys Cys Thr Lys  
 210 215 220

Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 225 230 235 240

Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser Trp Leu  
 245 250 255

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
 260 265 270

- 47 -

Val Trp Leu Leu Val Met Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
 275 280 285

Tyr Arg Ile Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe Phe Cys  
 290 295 300

Leu Trp Val Xaa Ile  
 305

<210> 42  
 <211> 1031  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> Nucleotide sequence of Patient C

<400> 42  
 tactacaaac cttgccagca aatccgcctc ctgcctctac caatcgccag tcaggaaggg 60  
 agcctacccc tctgaactcca cctttgagaa acactcatcc tcaggccatg cagtggaaact 120  
 ccacaaactt ccaccgaact ctacaagatc ccagagtgaaggcctgtat ctccctgctg 180  
 gtggctccag ttcaggaaca gtaaaccctg ttccgactac tgtctctcac acatcgtcaa 240  
 tcttatcgag gattggggac cctgcactga acatggagaa catcacatca ggattcctag 300  
 gaccctgct cgtgttacag gcggggtttt tcttgttgac aagaatcctc acaataccgc 360  
 agagtctaga ctctgtgttg acttctctca attttctagg ggggaccacc gtgtgccttg 420  
 gccaaaattc gcagtcccca acctccaatc actcaccaac ctctgtcct ccaactgtc 480  
 ctgggttatcg ctggatgtgt ctgcccgtt ttatcatatt cctcttcac ctgctgetat 540  
 gcctcatctt cttgttggtt cttctggact atcaaggat gttgcccgtt tgccctctaa 600  
 ttccaggatc ctcaaccacc agcacgggac catgcagaac ctgcacgact cctgctcaag 660  
 gaacctctwt gtatccctca tgttgctgta ccaaacctwc gmcgsaaat tgcacctgta 720  
 ttcccatccc atcatcctgg gctttcgga aattcctatg ggagtgggccc tcagcccgtt 780  
 tctcctgact cagtttacta gtgccatttg ttcagtggtt cgtagggctt tccccactg 840  
 tttggctttc agttatatgg atgatgtggt attggggggc aggtctgtac agcatcgtga 900  
 ggcccttttt accgctgtta ccaattttct tttgtctctg ggtatacatt taaccccga 960  
 caaaacaaaa agatgggggtt actctttaca tttcatgggc tatgtcattg gatgttatgg 1020  
 gtcattgcca c 1031

- 48 -

<210> 43  
 <211> 316  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> POL Trans of Patient

<400> 43

Thr Thr Asn Leu Ala Ser Lys Ser Ala Ser Cys Leu Tyr Gln Ser Pro  
 1 5 10 15

Val Arg Lys Ala Ala Tyr Pro Ser Asp-Ser Thr Phe Glu Lys His Ser  
 20 25 30

Ser Ser Gly His Ala Val Glu Leu His Lys Leu Pro Pro Asn Ser Thr  
 35 40 45

Arg Ser Gln Ser Glu Arg Pro Val Ser Pro Cys Trp Trp Leu Gln Phe  
 50 55 60

Arg Asn Ser Lys Pro Cys Ser Asp Tyr Cys Leu Ser His Ile Val Asn  
 65 70 75 80

Leu Ile Glu Asp Trp Gly Pro Cys Thr Glu His Gly Glu His His Ile  
 85 90 95

Arg Ile Pro Arg Thr Pro Ala Arg Val Thr Gly Gly Val Phe Leu Val  
 100 105 110

Asp Lys Asn Pro His Asn Thr Ala Glu Ser Arg Leu Val Val Asp Phe  
 115 120 125

Ser Gln Phe Ser Arg Gly Asp His Arg Val Pro Trp Pro Lys Phe Ala  
 130 135 140

Val Pro Asn Leu Gln Ser Leu Thr Asn Leu Leu Ser Ser Asn Leu Ser  
 145 150 155 160

Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His Ile Pro Leu His  
 165 170 175

Pro Ala Ala Met Pro His Leu Leu Val Gly Ser Ser Gly Leu Ser Arg  
 180 185 190

- 49 -

Tyr Val Ala Arg Leu Pro Ser Asn Ser Arg Ile Leu Asn His Gln His  
 195 200 205

Gly Thr Met Gln Asn Leu His Asp Ser Cys Ser Arg Asn Leu Tyr Phe  
 210 215 220

Val Ser Leu Met Leu Leu Tyr Gln Thr Phe Thr Gly Arg Lys Leu His  
 225 230 235 240

Leu Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro Met Gly  
 245 250 255

Val Gly Leu Ser Pro Phe Leu Leu Thr Gln Phe Thr Ser Ala Ile Cys  
 260 265 270

Ser Ala Leu Phe Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly Ile  
 275 280 285

His Leu Thr Pro Asp Lys Thr Lys Arg Trp Gly Tyr Ser Leu His Phe  
 290 295 300

Met Gly Tyr Val Ile Gly Cys Tyr Gly Ser Leu Pro  
 305 310 315

<210> 44

<211> 301

<212> PRT

<213> artificial sequence

<220>

<223> HBsAg Trans of Patient C

<400> 44

Leu Gln Thr Leu Pro Ala Asn Pro Pro Pro Ala Ser Thr Asn Arg Gln  
 1 5 10 15

Ser Gly Arg Gln Pro Thr Pro Leu Thr Pro Pro Leu Arg Asn Thr His  
 20 25 30

Pro Gln Ala Met Gln Trp Asn Ser Thr Asn Phe His Arg Thr Leu Gln  
 35 40 45

Asp Pro Arg Val Lys Gly Leu Tyr Leu Pro Ala Gly Gly Ser Ser Ser

- 50 -

50	55	60
Gly Thr Val Asn Pro Val Pro Thr Thr Val Ser His Thr Ser Ser Ile		
65	70	75 80
Leu Ser Arg Ile Gly Asp Pro Ala Leu Asn Met Glu Asn Ile Thr Ser		
	85	90 95
Gly Phe Leu Gly Pro Leu Leu Val Leu Gln Ala Gly Phe Phe Leu Leu		
	100	105 110
Thr Arg Ile Leu Thr Ile Pro Gln Ser Leu Asp Ser Trp Trp Thr Ser		
	115	120 125
Leu Asn Phe Leu Gly Gly Thr Thr Val Cys Leu Gly Gln Asn Ser Gln		
	130	135 140
Ser Pro Thr Ser Asn His Ser Pro Thr Ser Cys Pro Pro Thr Cys Pro		
	145	150 155 160
Gly Tyr Arg Trp Met Cys Leu Arg Arg Phe Ile Ile Phe Leu Phe Ile		
	165	170 175
Leu Leu Leu Cys Leu Ile Phe Leu Leu Val Leu Leu Asp Tyr Gln Gly		
	180	185 190
Met Leu Pro Val Cys Pro Leu Ile Pro Gly Ser Ser Thr Thr Ser Thr		
	195	200 205
Gly Pro Cys Arg Thr Cys Thr Thr Pro Ala Gln Gly Thr Ser Met Leu		
	210	215 220
Tyr Pro Ser Cys Cys Cys Thr Lys Pro Ser Thr Ala Ala Asn Cys Thr		
	225	230 235 240
Cys Ile Pro Ile Pro Ser Ser Trp Ala Phe Gly Lys Phe Leu Trp Glu		
	245	250 255
Trp Ala Ser Ala Arg Phe Ser Leu Ser Leu Leu Val Pro Phe Val Gln		
	260	265 270
Trp Phe Val Gly Leu Ser Pro Thr Val Trp Leu Ser Val Ile Trp Met		
	275	280 285



- 51 -

Met Trp Tyr Trp Gly Pro Gly Leu Tyr Ser Ile Val Arg  
 290 295 300

<210> 45  
 <211> 888  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> Nucleotide sequence of Patient D

<400> 45  
 tggtcacagt gccaacagtt cctcctcctg cctccaccaa tcggcagtc gggaggcagc 60  
 ctactcccat ctctccacct ctaagagaca gtcacctca ggccatgggt gctcagcctg 120  
 ctggtggctc cagttcagga aactcaacc ctgttccaa tattgcctct cacatctcgt 180  
 caatctcctt gaggactggg gaccctgccc cgaacatgga gaacatcaca tcaggattcc 240  
 taggaccctt gctcgtgta caggcggggg ttttcttggt gacaagaatc ctcaacaatac 300  
 cgcagagtct agactcgtgg tggacttctc tcagttttct agggggatca cccgtgtgtc 360  
 ttggccaaaa ttgcagtc ccaacctcca atcactcacc aacctcctgt cctccaattt 420  
 gacctgggta tcgctggata tgtctgcggc gttttatcat attcctcttc atcctgccgc 480  
 tatgcctcat cttcttattg gttcttctgg attatcaagg tatgttgccc gtttgcctc 540  
 taattccagg atccacaaca accagtgcgg gaccctgcaa aacctgcacg actcctgtc 600  
 aaggcaactc tatgtttccc tcatgttgct gtacaaaacc tacggatgga aattgcacct 660  
 gtattcccat cccatcatct tgggctttcg caaaatacct atgggagtgg gcctcagtcc 720  
 gtttctcttg gtcagttta ctagtccat ttgttcagtg attcgtaggg ctttccccc 780  
 ctggttggtt ttcagctata ttgatgatgt ggtactgggg gccaaagtctg cacaacatct 840  
 tgagtccctt tataccgctg ttaccaattt tcttttgtct ttgggtat 888

<210> 46  
 <211> 295  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> Pol Trans of Patient D

<400> 46

Gly His Ser Ala Asn Ser Ser Ser Ser Cys Leu His Gln Ser Ala Val

- 52 -

1 5 10 15

Arg Glu Ala Ala Tyr Ser His Leu Ser Thr Ser Lys Arg Gln Ser Ser  
20 25 30

Ser Gly His Gly Gly Ser Ala Cys Trp Trp Leu Gln Phe Arg Asn Thr  
35 40 45

Gln Pro Cys Ser Gln Tyr Cys Leu Ser His Leu Val Asn Leu Leu Glu  
50 55 60

Asp Trp Gly Pro Cys Ala Glu His Gly Glu His His Ile Arg Ile Pro  
65 70 75 80

Arg Thr Pro Ala Arg Val Thr Gly Gly Val Phe Leu Val Asp Lys Asn  
85 90 95

Pro His Asn Thr Ala Glu Ser Arg Leu Val Val Asp Phe Ser Gln Phe  
100 105 110

Ser Arg Gly Ile Thr Arg Val Ser Trp Pro Lys Phe Ala Val Pro Asn  
115 120 125

Leu Gln Ser Leu Thr Asn Leu Leu Ser Ser Asn Leu Thr Trp Leu Ser  
130 135 140

Leu Asp Met Ser Ala Ala Phe Tyr His Ile Pro Leu His Pro Ala Ala  
145 150 155 160

Met Pro His Leu Leu Ile Gly Ser Ser Gly Leu Ser Arg Tyr Val Ala  
165 170 175

Arg Leu Ser Ser Asn Ser Arg Ile His Asn Asn Gln Cys Gly Thr Leu  
180 185 190

Gln Asn Leu His Asp Ser Cys Ser Arg Gln Leu Tyr Val Ser Leu Met  
195 200 205

Leu Leu Tyr Lys Thr Tyr Gly Trp Lys Leu His Leu Tyr Ser His Pro  
210 215 220

Ile Ile Leu Gly Phe Arg Lys Ile Pro Met Gly Val Gly Leu Ser Pro  
225 230 235 240

Phe Leu Leu Ala Gln Phe Thr Ser Ala Ile Cys Ser Val Ile Arg Arg  
245 250 255

Gly Ala Lys Ser Ala Gln His Leu Glu Ser Leu Tyr Thr Ala Val Thr  
275 280 285

```
<210> 47
<211> 293
<212> PRT
<213> artificial sequence
```

<220>  
<223> HBsAg Trans of Patient D

<400> 47

Val	Thr	Val	Pro	Thr	Val	Pro	Pro	Pro	Ala	Ser	Thr	Asn	Arg	Gln	Ser
1				5					10					15	

Gly Arg Gln Pro Thr Pro Ile Ser Pro Pro Leu Arg Asp Ser His Pro  
20 25 30

Gln Ala Met Val Ala Gln Pro Ala Gly Gly Ser Ser Ser Gly Thr Leu  
35 40 45

Asn Pro Val Pro Asn Ile Ala Ser His Ile Ser Ser Ile Ser Leu Arg  
50 55 60

Thr Gly Asp Pro Ala Pro Asn Met Glu Asn Ile Thr Ser Gly Phe Leu  
65 70 75 80

Gly Pro Leu Leu Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile  
85 90 95

Leu Thr Ile Pro Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Ser Phe  
100 105 110

Leu Gly Gly Ser Pro Val Cys Leu Gly Gln Asn Ser Gln Ser Pro Thr

- 54 -

115                      120                      125  
 Ser Asn His Ser Pro Thr Ser Cys Pro Pro Ile Pro Gly Tyr Arg Trp  
 130                      135                      140  
 Ile Cys Leu Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Pro Leu Cys  
 145                      150                      155                      160  
 Leu Ile Phe Leu Leu Val Leu Leu Asp Tyr Gln Gly Met Leu Pro Val  
 165                      170                      175  
 Cys Pro Leu Ile Pro Gly Ser Thr Thr Thr Ser Ala Gly Pro Cys Lys  
 180                      185                      190  
 Thr Cys Thr Thr Pro Ala Gln Gly Asn Ser Met Phe Pro Ser Cys Cys  
 195                      200                      205  
 Cys Thr Lys Pro Thr Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser  
 210                      215                      220  
 Ser Trp Ala Phe Ala Lys Tyr Leu Trp Glu Trp Ala Ser Val Arg Phe  
 225                      230                      235                      240  
 Ser Trp Leu Ser Leu Leu Val Pro Phe Val Gln Phe Val Gly Leu Ser  
 245                      250                      255  
 Pro Thr Val Trp Leu Ser Ala Ile Leu Met Met Trp Tyr Trp Gly Pro  
 260                      265                      270  
 Ser Leu His Asn Ile Leu Ser Pro Phe Ile Pro Leu Leu Pro Ile Phe  
 275                      280                      285  
 Phe Cys Leu Trp Val  
 290

&lt;210&gt; 48

&lt;211&gt; 591

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; Nucleotide sequence of Patient E

&lt;400&gt; 48

tcctgtcctc caattgtcc tggttatcgc tggatgtgtc tgcggcggtt tatgatattc 60

- 55 -

```

ctcttcatcc tgctgctatg cctcatcttc ttattgggtc ttctggatta tcaaggtatg      120
ttgcccgtct gtectctaatt tccaggatca acaacaacca gtacgggacc atgcaaaacc      180
aaaacctgca cgactcctgc tcaaggcaac tctatgtttc cctcatgttg ctgtacaaaa      240
cctacgggatg gaaattgcac ctgtattccc atcccatcgt cctgggcttt cgcaaaattc      300
ctatgggagt gggcctcagt ccgtttctct tggctcagtt tactagtgcc atttgttcag      360
tggttcgtag ggctttcccc cactgtttgg ctctcagcta tatggatgat gtggtattgg      420
gggccaagtc tgtacagcat cgtgaggccc ttatacagc tgttaccaat tttcttttgt      480
ctctgggtat acatttaaac cctaacaaaa caaaaagatg gggttattcc ctaaacttca      540
tgggttacat aattggaagt tggggaacat tgccacagga tcatattgta c                591

```

<210> 49  
 <211> 186  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> Pol Trans of Patient B

<400> 49

```

Ser Asn Leu Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr Asp
1              5              10              15

```

```

Ile Pro Leu His Pro Ala Ala Met Pro-His Leu Leu Ile Gly Ser Ser
              20              25              30

```

```

Gly Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile Asn
35              40              45

```

```

Asn Asn Gln Tyr Gly Thr Met Gln Asn Gln Asn Leu His Asp Ser Cys
50              55              60

```

```

Ser Arg Gln Leu Tyr Val Ser Leu Met Leu Leu Tyr Lys Thr Tyr Gly
65              70              75              80

```

```

Trp Lys Leu His Leu Tyr Ser His Pro Ile Val Leu Gly Phe Arg Lys
85              90              95

```

```

Ile Pro Met Gly Val Gly Leu Ser Pro Phe Leu Leu Ala Gln Phe Thr
100              105              110

```

- 56 -

Ser Ala Ile Cys Ser Val Val Arg Arg Ala Phe Pro His Cys Leu Ala  
 115 120 125

Phe Ser Tyr Met Asp Asp Val Val Leu Gly Ala Lys Ser Val Gln His  
 130 135 140

Arg Glu Ala Leu Tyr Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly  
 145 150 155 160

Ile His Leu Asn Pro Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu Asn  
 165 170 175

Phe Met Gly Tyr Ile Ile Gly Ser Trp Gly  
 180 185

<210> 50

<211> 165

<212> PRT

<213> artificial sequence

<220>

<223> HBsAg Trans of Patient E

<400> 50

Ser Cys Pro Pro Ile Cys Pro Gly Tyr Arg Trp Met Cys Leu Arg Arg  
 1 5 10 15

Phe Met Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe Leu Leu  
 20 25 30

Val Leu Leu Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu Ile Pro  
 35 40 45

Gly Ser Thr Thr Thr Ser Thr Gly Pro Cys Lys Thr Lys Thr Cys Thr  
 50 55 60

Thr Pro Ala Gln Gly Asn Ser Met Phe Pro Ser Cys Cys Cys Thr Lys  
 65 70 75 80

Pro Thr Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala  
 85 90 95

Phe Ala Lys Phe Leu Trp Glu Trp Ala Ser Val Arg Phe Ser Trp Leu  
 100 105 110

- 57 -

Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr  
 115 120 125

Val Trp Leu Ser Ala Ile Trp Met Met Trp Tyr Trp Gly Pro Ser Leu  
 130 135 140

Tyr Ser Ile Val Arg Pro Phe Ile Gln Leu Leu Pro Ile Phe Phe Cys  
 145 150 155 160

Leu Trp Val Tyr Ile  
 165

<210> 51  
 <211> 669  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> Nucleotide sequence of Patient

<400> 51  
 aatcctcaca ataccgcaga gtctagactt cgtggtgact tototcaatt ttctagggga 60  
 ccacccgtgt gtcttggcca aaattcgcag tccccaacct ccaatcactc accaaccctct 120  
 tgtcctccaa tttgtcctgg ttatcgctgg atgtgtctgc ggcgttttat catatccctc 180  
 ttcatcctgc tgctatgcct catcttctta ttggttcttc tggattatca aggtatgttg 240  
 cccgtttgtc ctctaattcc aggatccaca acaaccagta cgggaccctg caaaacctgc 300  
 acgactcctg ctcaaggcaa ctctatgttt ccctcatggt gctgtacaaa acctacggat 360  
 ggaaattgca cmtgtattcc catcccatca tcttgggctt tcgcaaaata cctatgggag 420  
 tgggcctcag tccgtttctc ttggttcagt ttactagtgc catttgttca gtggttcgta 480  
 gggctttccc ccaactgtttg gctttcagct atatggatga tattgtactg ggggccaagt 540  
 ctgtacaaca tcttgagtcc ctttataccg ctgttaccaa ttttcttttg tctttgggta 600  
 tacatttaac ccctaacaaa acaaagagat ggggttatte cctgaatttc atgggttatg 660  
 taattggaa 669

<210> 52  
 <211> 181  
 <212> PRT  
 <213> artificial sequence

- 58 -

&lt;220&gt;

&lt;223&gt; Deduced sequence of DNA polymerase of Patient F

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (83)..(83)

&lt;223&gt; x = any amino acid

&lt;400&gt; 52

Ser Asn Leu Ser Trp Leu Ser Leu Asp Val Ser Ala Ala Phe Tyr His  
 1 5 10 15

Ile Pro Leu His Pro Ala Ala Met Pro His Leu Leu Ile Gly Ser Ser  
 20 25 30

Gly Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Asn Ser Arg Ile His  
 35 40 45

Asn Asn Gln Tyr Gly Thr Leu Gln Asn Leu His Asp Ser Cys Ser Arg  
 50 55 60

Gln Leu Tyr Val Ser Leu Met Leu Leu Tyr Lys Thr Tyr Gly Trp Lys  
 65 70 75 80

Leu His Xaa Tyr Ser His Pro Ile Ile Leu Gly Phe Arg Lys Ile Pro  
 85 90 95

Met Gly Val Gly Leu Ser Pro Phe Leu Leu Val Gln Phe Thr Ser Ala  
 100 105 110

Ile Cys Ser Val Val Arg Arg Ala Phe Pro His Cys Leu Ala Phe Ser  
 115 120 125

Tyr Met Asp Asp Ile Val Leu Gly Ala Lys Ser Val Gln His Leu Glu  
 130 135 140

Ser Leu Tyr Thr Ala Val Thr Asn Phe Leu Leu Ser Leu Gly Ile His  
 145 150 155 160

Leu Thr Pro Asn Lys Thr Lys Arg Trp Gly Tyr Ser Leu Asn Phe Met  
 165 170 175

Gly Tyr Val Ile Gly  
 180



- 59 -

--

<210> 53  
 <211> 160  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of Patient F

<400> 53

Pro Ile Cys Pro Gly Tyr Arg Trp Met Cys Leu Arg Arg Phe Ile Ile  
 1 5 10 15

Ser Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe Leu Leu Val Leu Leu  
 20 25 30

Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu Ile Pro Gly Ser Thr  
 35 40 45

Thr Thr Ser Thr Gly Pro Cys Lys Thr Cys Thr Thr Pro Ala Gln Gly  
 50 55 60

Asn Ser Met Phe Pro Ser Cys Cys Cys Thr Lys Pro Thr Asp Gly Asn  
 65 70 75 80

Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala Phe Ala Lys Tyr Leu  
 85 90 95

Trp Glu Trp Ala Ser Val Arg Phe Ser Trp Phe Ser Leu Leu Val Pro  
 100 105 110

Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr Val Trp Leu Ser Ala  
 115 120 125

Ile Trp Met Ile Leu Tyr Trp Gly Pro Ser Leu Tyr Asn Ile Leu Ser  
 130 135 140

Pro Phe Ile Pro Leu Leu Pro Ile Phe Phe Cys Leu Trp Val Tyr Ile  
 145 150 155 160

<210> 54  
 <211> 554  
 <212> DNA  
 <213> artificial sequence

- 60 -

&lt;220&gt;

&lt;223&gt; Nucleotide sequence of Patient G

&lt;400&gt; 54

```

tccaatttgt cctgggtatc gctggatgtg tctgcggtt tttatcatat tcctcttcat      60
cctgctgcta tgcctcatct tcttggttgt tcttctggac tatcaaggta tgttgccgt      120
ttgtcctcta cttccaggaa catcaactac cagcacggga ccatgcaaga cctgcacgac      180
tcctgctcaa ggaacctcta tgtttccctc ttgttgctgt acaaaacctt cggacggaaa      240
ttgcacttgt attcccatcc catcgtcttg ggctttcgca agattcctat gggagtgggc      300
ctcagtcctg ttctcttggc tcartttact agtgccattt gttcagtggg tcgtagggct      360
ttccccact gtttggtttt cagttatatt gatgatgtgg tattgggggc caagtctgta      420
caacatcttg aatccctttt tacctctatt accaattttc ttatgtcttt gggatacat      480
ttaaaccta agaaaaccaa acgttggggc tactccotta acttcatggg atatgtaatt      540
ggaagttggg gtac                                                              554

```

&lt;210&gt; 55

&lt;211&gt; 184

&lt;212&gt; PRT

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; Deduced sequence of DNA polymerase of Patient G

&lt;400&gt; 55

```

Ser Asn Leu Ser Trp Val Ser Leu Asp Val Ser Ala Ala Phe Tyr His
1           5           10          15
Ile Pro Leu His Pro Ala Ala Met Pro His Leu Leu Val Gly Ser Ser
          20          25          30
Gly Leu Ser Arg Tyr Val Ala Arg Leu Ser Ser Thr Ser Arg Asn Ile
          35          40          45
Asn Tyr Gln His Gly Thr Met Gln Asp Leu His Asp Ser Cys Ser Arg
          50          55          60
Asn Leu Tyr Val Ser Leu Leu Leu Tyr Lys Thr Phe Gly Arg Lys
65          70          75          80
Leu His Leu Tyr Ser His Pro Ile Val Leu Gly Phe Arg Lys Ile Pro
          85          90          95

```

- 61 -

Met Gly Val Gly Leu Ser Pro Phe Leu Leu Ala Gln Phe Thr Ser Ala  
 100 105 110

Ile Cys Ser Val Val Arg Arg Ala Phe Pro His Cys Leu Ala Phe Ser  
 115 120 125

Tyr Ile Asp Asp Val Val Leu Gly Ala Lys Ser Val Gln His Leu Glu  
 130 135 140

Ser Leu Phe Thr Ser Ile Thr Asn Phe Leu Met Ser Leu Gly Ile His  
 145 150 155 160

Leu Asn Pro Lys Lys Thr Lys Arg Trp Gly Tyr Ser Leu Asn Phe Met  
 165 170 175

Gly Tyr Val Ile Gly Ser Trp Gly  
 180

<210> 56  
 <211> 160  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of Patient G

<220>  
 <221> MISC\_FEATURE  
 <222> (108)..(108)  
 <223> x = any amino acid

<400> 56

Pro Ile Cys Pro Gly Tyr Arg Trp Met Cys Leu Arg Arg Phe Ile Ile  
 1 5 10 15

Phe Leu Phe Ile Leu Leu Leu Cys Leu Ile Phe Leu Leu Val Leu Leu  
 20 25 30

Asp Tyr Gln Gly Met Leu Pro Val Cys Pro Leu Leu Pro Gly Thr Ser  
 35 40 45

Thr Thr Ser Thr Gly Pro Cys Lys Thr Cys Thr Thr Pro Ala Gln Gly  
 50 55 60

- 62 -

Thr Ser Met Phe Pro Ser Cys Cys Cys Thr Lys Pro Ser Asp Gly Asn  
65 70 75 80

Cys Thr Cys Ile Pro Ile Pro Ser Ser Trp Ala Phe Ala Arg Phe Leu  
85 90 95

Trp Glu Trp Ala Ser Val Arg Phe Ser Trp Leu Xaa Leu Leu Val Pro  
100 105 110

Phe Val Gln Trp Phe Val Gly Leu Ser Pro Thr Val Trp Leu Ser Val  
115 120 125

Ile Leu Met Met Trp Tyr Trp Gly Pro Ser Leu Tyr Asn Ile Leu Asn  
130 135 140

Pro Phe Leu Pro Leu Leu Pro Ile Phe Leu Cys Leu Trp Val Tyr Ile  
145 150 155 160

<210> 57

<211> 1045

<212> DNA

<213> artificial sequence

<220>

<223> Nucleotide sequence of Patient H

<400> 57

cagcaaattcc gcctcctgcc tctaccaatc gccagtcagg aaggcagcct acccctctgt 60  
ctccaccttt grgaaacact catcctcagg ccatgcagtg gaactccaca acctccacc 120  
aaactctgcw agatcccaga gtgagaggcc tgtatttccc tgctgggtggc tccagttcag 180  
gaacagtaaa ccctgttccg acttctgtct ctcacacatc gtcaatcttc tcgaggattg 240  
gggwcctgc gctgaacatg gagaacatca catcaggatt cctaggaccc ctgctcgtgt 300  
tacaggcggg gtttttcttg ttgacaagaa tcctcacaat accgcagagt ctagactcgt 360  
ggtggacttc totcaatttt ctagggggaa ctaccgtgtg tcttggccaa aattcgcagt 420  
tcccaacctc caatcactca ccaacctcct gtctccaac ttgwcctggg tatcgctgga 480  
tgtrtctgcg gcgttttata atcttctctt tcctcctgct gctatgcctc atcttcttgt 540  
tggttcttct ggactatcaa ggtatgttgc ccgtttgtcc tctarttcca ggatcttcaa 600  
ccaccagcac gggaccatgc agaacctgca cgactcctgc tcaaggaamc tctatgaatc 660  
cctcctgttg ctgtacaaa ccttcggacg gaaattgcac ctgtattccc atcccatcat 720

- 63 -

cctgggcttt cggaaaattc ctatgggagt gggcctcagc ccgtttctcc tgrctcagtt 780  
tactagtgcc atttggtcag tgggtcgtag ggctttcccc cactgtttgg ctttcagtta 840  
tatggatgat gtggtattgg gggccaagtc tgtaymgcat cttragtccc tttttaccgc 900  
tgttaccaat tttcttttgt ctytgggtat acatttaaac cctmacaaaa caaaaagatg 960  
gggttactct ttacatttca tgggctatgt cattggatgt tatgggtcat tgccacaaga 1020  
tcacatcaga cagaaaatca aagaa 1045

<210> 58  
<211> 348  
<212> PRT  
<213> artificial sequence

<220>  
<223> Deduced sequence of DNA polymerase of Patient H

<220>  
<221> MISC\_FEATURE  
<222> (24)..(24)  
<223> x = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (155)..(155)  
<223> x = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (195)..(195)  
<223> x = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (216)..(216)  
<223> x = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (258)..(258)  
<223> x = any amino acid

<220>  
<221> MISC\_FEATURE  
<222> (292)..(292)  
<223> x = any amino acid

- 64 -

<220>  
 <221> MISC\_FEATURE  
 <222> (295)..(295)  
 <223> x = any amino acid

<220>  
 <221> MISC\_FEATURE  
 <222> (315)..(315)  
 <223> x = any amino acid

<400> 58

Ser Lys Ser Ala Ser Cys Leu Tyr Gln Ser Pro Val Arg Lys Ala Ala  
 1 5 10 15

Tyr Pro Ser Val Ser Thr Phe Xaa Lys His Ser Ser Ser Gly His Ala  
 20 25 30

Val Glu Leu His Asn Leu Pro Pro Asn Ser Ala Arg Ser Gln Ser Glu  
 35 40 45

Arg Pro Val Phe Pro Cys Trp Trp Leu Gln Phe Arg Asn Ser Lys Pro  
 50 55 60

Cys Ser Asp Phe Cys Leu Ser His Ile Val Asn Leu Leu Glu Asp Trp  
 65 70 75 80

Gly Pro Cys Ala Glu His Gly Glu His His Ile Arg Ile Pro Arg Thr  
 85 90 95

Pro Ala Arg Val Thr Gly Gly Val Phe Leu Val Asp Lys Asn Pro His  
 100 105 110

Asn Thr Ala Glu Ser Arg Leu Val Val Asp Phe Ser Gln Phe Ser Arg  
 115 120 125

Gly Asn Tyr Arg Val Ser Trp Pro Lys Phe Ala Val Pro Asn Leu Gln  
 130 135 140

Ser Leu Thr Asn Leu Leu Ser Ser Asn Leu Xaa Trp Leu Ser Leu Asp  
 145 150 155 160

Val Ser Ala Ala Phe Tyr His Leu Pro Leu His Pro Ala Ala Met Pro  
 165 170 175

- 65 -

His Leu Leu Val Gly Ser Ser Gly Leu Ser Arg Tyr Val Ala Arg Leu  
 180 185 190

Ser Ser Xaa Ser Arg Ile Phe Asn His Gln His Gly Thr Met Gln Asn  
 195 200 205

Leu His Asp Ser Cys Ser Arg Xaa Leu Tyr Glu Ser Leu Leu Leu Leu  
 210 215 220

Tyr Gln Thr Phe Gly Arg Lys Leu His Leu Tyr Ser His Pro Ile Ile  
 225 230 235 240

Leu Gly Phe Arg Lys Ile Pro Met Gly Val Gly Leu Ser Pro Phe Leu  
 245 250 255

Leu Xaa Gln Phe Thr Ser Ala Ile Cys Ser Val Val Arg Arg Ala Phe  
 260 265 270

Pro His Cys Leu Ala Phe Ser Tyr Met Asp Asp Val Val Leu Gly Ala  
 275 280 285

Lys Ser Val Xaa His Leu Xaa Ser Leu Phe Thr Ala Val Thr Asn Phe  
 290 295 300

Leu Leu Ser Leu Gly Ile His Leu Asn Pro Xaa Lys Thr Lys Arg Trp  
 305 310 315 320

Gly Tyr Ser Leu His Phe Met Gly Tyr Val Ile Gly Cys Tyr Gly Ser  
 325 330 335

Leu Pro Gln Asp His Ile Arg Gln Lys Ile Lys Glu  
 340 345

<210> 59  
 <211> 311  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> HBsAg Trans of Patient H

<220>  
 <221> MISC\_FEATURE  
 <222> (24)..(24)

- 66 -

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (43)..(43)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (81)..(81)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (154)..(154)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (161)..(161)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (195)..(195)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (216)..(216)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (257)..(257)

<223> x = any amino acid

<220>

<221> MISC\_FEATURE

<222> (292)..(292)

<223> x = any amino acid

<400> 59

Ala Asn Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro  
1 5 10 15



- 67 -

Thr Pro Leu Ser Pro Pro Leu Xaa Asn Thr His Pro Gln Ala Met Gln  
 20 25 30

Trp Asn Ser Thr Thr Phe His Gln Thr Leu Xaa Asp Pro Arg Val Arg  
 35 40 45

Gly Leu Tyr Phe Pro Ala Gly Gly Ser Ser Ser Gly Thr Val Asn Pro  
 50 55 60

Val Pro Thr Ser Val Ser His Thr Ser Ser Ile Phe Ser Arg Ile Gly  
 65 70 75 80

Xaa Pro Ala Leu Asn Met Glu Asn Ile Thr Ser Gly Phe Leu Gly Pro  
 85 90 95

Leu Leu Val Leu Gln Ala Gly Phe Phe Leu Leu Thr Arg Ile Leu Thr  
 100 105 110

Ile Pro Gln Ser Leu Asp Ser Trp Trp Thr Ser Leu Asn Phe Leu Gly  
 115 120 125

Gly Thr Thr Val Cys Leu Gly Gln Asn Ser Gln Phe Pro Thr Ser Asn  
 130 135 140

His Ser Pro Thr Ser Cys Pro Pro Thr Xaa Pro Gly Tyr Arg Trp Met  
 145 150 155 160

Xaa Leu Arg Arg Phe Ile Ile Phe Leu Phe Ile Leu Leu Leu Cys Leu  
 165 170 175

Ile Phe Leu Leu Val Leu Leu Asp Tyr Gln Gly Met Leu Pro Val Cys  
 180 185 190

Pro Leu Xaa Pro Gly Ser Ser Thr Thr Ser Thr Gly Pro Cys Arg Thr  
 195 200 205

Cys Thr Thr Pro Ala Gln Gly Xaa Ser Met Asn Pro Ser Cys Cys Cys  
 210 215 220

Thr Lys Pro Ser Asp Gly Asn Cys Thr Cys Ile Pro Ile Pro Ser Ser  
 225 230 235 240

Trp Ala Phe Gly Lys Phe Leu Trp Glu Trp Ala Ser Ala Arg Phe Ser

- 68 -

245

250

255

Xaa Leu Ser Leu Leu Val Pro Phe Val Gln Trp Phe Val Gly Leu Ser  
260 265 270

Pro Thr Val Trp Leu Ser Val Ile Trp Met Met Trp Tyr Trp Gly Pro  
275 280 285

Ser Leu Tyr Xaa Ile Leu Ser Pro Phe Leu Pro Leu Leu Pro Ile Phe  
290 295 300

Phe Cys Leu Trp Val Tyr Ile  
305 310

a

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU03/00432

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>												
Int. Cl. <sup>7</sup> : C12N7/00, 15/01, 15/36, 15/51, C12Q1/70												
According to International Patent Classification (IPC) or to both national classification and IPC												
<b>B. FIELDS SEARCHED</b>												
Minimum documentation searched (classification system followed by classification symbols) SEE BELOW												
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched SEE BELOW												
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPIDS, MEDLINE, CA. KEYWORDS: Hepatitis B/Nucleotide/Nucleoside/Resistan?/Muta?/Adefovir/ Lamivudine/Tenofovir/Thiacydidine/HBsAg/Polymerase and similar terms												
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>												
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.										
X	WO 01/94559 (Melbourne Health) 13 December 2001  Claims, p.12-19, p.22 line 25- p.24 line 25, and p. 25 line 30 - p. 26 line 9	1-125, 169-202										
X	WO 01/57244 (Melbourne Health) 9 August 2001  Claims, Table 1, p. 17 line 7 - p. 21 line 12 and p. 25 line 16 - p. 33 line 2	1-17, 19-74, 80, 86, 95, 106, 119, 123-127										
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex												
<p>* Special categories of cited documents:</p> <table border="0"> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E" earlier application or patent but published on or after the international filing date</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td>"&amp;" document member of the same patent family</td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	"P" document published prior to the international filing date but later than the priority date claimed	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention											
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone											
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art											
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family											
"P" document published prior to the international filing date but later than the priority date claimed												
Date of the actual completion of the international search 27 May 2003		Date of mailing of the international search report 06 JUN 2003										
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929		Authorized officer  <b>ALISTAIR BESTOW</b> Telephone No : (02) 6283 2450										

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU03/00432

**Box I** Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos :  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos : 126, 159, 160 (completely), 161-168 (partially)  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
The claims are not restricted to the technical features of the invention, namely hepatitis B variants, methods of using these variants and products of these methods.  
Claims 161-168 have only been searched in so far as they relate to HBV nucleic acid or peptides sequences, antibodies, ribozymes and antisense that are capable of inhibiting the variant HBVs of the invention
3. ☐ Claims Nos :  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

**Box II** Observations where unity of invention is lacking (Continuation of Item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See Supplemental Box for summary inventions.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU03/00432

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/21317 (Western Health Care Network) 22 May 1998  Claims, p. 4 line 5 - p. 7 line 12 and p. 8 line 28 - p. 11 line 20,	1-17, 19-37, 39, 40, 42-50, 54, 55, 58-74, 76-78, 80-90, 94, 95, 98, 100-114, 118, 119, 122, 170- 178, 180-182, 184-186, 190, 194-202
X	Yeh, C. et al., 2000, <i>Clearance of the original hepatitis B virus YMDD-motif mutants with emergence of distinct lamivudine-resistant mutants during prolonged lamivudine therapy</i> , Hepatology, 31:1318-1325  See whole document	1-17, 19-37, 39, 40, 49, 50, 54, 55, 58-74, 76, 77, 78, 82, 85, 86, 89, 90, 94, 95, 98, 100-102, 106, 109, 113, 114, 118, 119, 122, 127-129
X	Seignerres, B. et al., 2000, <i>Evolution of Hepatitis B virus polymerase gene sequence during famciclovir therapy for chronic hepatitis B</i> , Journal of Infectious Diseases, 181:1221-33  See whole document	1-17, 19-38, 42, 50, 54, 57, 58-74, 80, 85, 86, 90, 97, 104, 109, 110, 114, 118, 121, 122, 127-129
X	Ogata, N. et al., 1999, <i>Novel patterns of amino acid mutations in the hepatitis b virus polymerase in association with resistance to lamivudine therapy in japanese patients with chronic hepatitis B</i> , Journal of Medical Virology, 59:270-276  See whole document	1-17, 19-37, 43, 59-74, 81, 105
X	Bock, C. et al., 2002(February), <i>Selection of hepatitis B virus polymerase mutants with enhanced replication by lamivudine treatment after liver transplantation</i> , Gastroenterology, 122:264-273  See whole document	1-17, 19-37, 39, 44, 54, 55, 58-74, 83, 94, 95, 98, 100, 107, 118, 119, 122

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU03/00432

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>Cane, P. et al., 1999, <i>Analysis of hepatitis B virus quasispecies changes during the emergence and reversion of lamivudine resistance in liver transplantation</i>, Antiviral Therapy, 4:7-14</p> <p>See whole document</p>	1-17, 19-37, 39, 43, 45, 49, 50, 55, 58-74, 81, 84, 86, 89, 90, 94, 95, 98, 100, 102, 108, 110, 113, 114, 118, 119, 122

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU03/00432

**Supplemental Box**

(To be used when the space in any of Boxes I to VIII is not sufficient)

**Continuation of Box No II:**

Invention 1: Claims 1- 20, 38, 59-75, 99, 123-125, 127, 130-134, 170, 178, 179, 195 (all partially) relating to an isolated Hepatitis B virus (HBV) variant comprising a mutation in a gene encoding DNA polymerase resulting in an amino acid addition, substitution and/or deletion at amino acid position rt21 of the HBV DNA polymerase, the use of these variants, and methods for determining whether a HBV variant exhibits decrease sensitivity to anti-viral agents or reduced interactivity to an antibody to HBV surface antigen.

Invention 2: 1- 20, 38, 59-75, 77, 99, 101, 123-125, 127, 130-134, 136, 170, 178, 179, 181, 195 (all partially) relating to an isolated HBV variant comprising a mutation in a gene encoding DNA polymerase resulting in an amino acid addition, substitution and/or deletion at amino acid position rt122 of the HBV DNA polymerase, the use of these variants, and methods for determining whether a HBV variant exhibits decrease sensitivity to anti-viral agents or reduced interactivity to an antibody to HBV surface antigen.

Inventions 3-37: relating to HBV variants comprising a mutation in a gene encoding DNA polymerase and/or HBsAg resulting in an amino acid addition, substitution and/or deletion at amino acid positions rt124, rt28, rt130, etc, rt251 or the equivalent position in the overlapping HBsAg, the use of these variants, and methods for determining whether a HBV variant exhibits decrease sensitivity to anti-viral agents or reduced interactivity to an antibody to HBV surface antigen.

The common feature linking the group of inventions resides in the elucidation of the link between reduced sensitivity to anti-HBV agents such as ADV, LMV, FTC and TFV and mutation of the nucleotide sequence encoding HBV DNA polymerase or the overlapping S gene (HBsAg). However, this is already known in the art (see Chen et al., 1999, Human hepatitis B virus mutants: significance of molecular changes, FEBS Letters, 453:237-242, and citations below), as such the claims relate to multiple inventions, *a posteriori*.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU03/00432

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
WO	200194559	AU	20008109	AU	200163672	EP	1297109
WO	200157244	AU	200131415	EP	1257661		
WO	9821317	AU	37628/97	EP	964916	US	6555311
							END OF ANNEX